

## Investigator's Brochure

<b>SPONSOR</b>	Multidisciplinary Association for Psychedelic Studies (MAPS)
<b>PRODUCT</b>	3,4-methylenedioxymethamphetamine (MDMA)
<b>IND #</b>	063384
<b>DATA CUT-OFF DATE</b>	01 October 2015
<b>EFFECTIVE DATE</b>	30 March 2016
<b>EDITION</b>	8 <sup>th</sup> Edition
<b>REPLACES</b>	7 <sup>th</sup> Edition (dated 01 August 2013)

## Table of Contents

<b>List of Figures and Tables</b> .....	<b>4</b>
<b>List of Abbreviations</b> .....	<b>6</b>
<b>1.0 Summary</b> .....	<b>8</b>
<b>2.0 Introduction</b> .....	<b>10</b>
<b>3.0 Physical, Chemical, and Pharmaceutical Properties and Formulation</b> .....	<b>12</b>
<b>4.0 Nonclinical Studies</b> .....	<b>13</b>
<b>4.1 Nonclinical Pharmacology</b> .....	<b>14</b>
<b>4.2 Pharmacology in Animals</b> .....	<b>14</b>
4.2.1 Pharmacokinetics in Animals.....	14
4.2.2 Pharmacodynamics in Animals .....	15
4.2.2.1 Stable Effects on Gene Expression in Animals .....	17
4.2.2.2 Immunological Effects in Animals .....	18
4.2.2.3 Thermoregulatory Effects in Animals.....	18
4.2.2.4 Cardiovascular Effects in Animals .....	19
4.2.2.5 Osmoregulatory Effects in Animals.....	19
4.2.2.6 Neurobiological Effects in Animals.....	20
4.2.2.7 Neuropsychological Effects in Animals .....	20
<b>4.3 Physiological Effects in Epidemiological Settings</b> .....	<b>21</b>
4.3.1 Immunological Effects .....	22
4.3.2 Thermoregulatory Effects .....	22
4.3.3 Cardiovascular Effects .....	23
4.3.4 Osmoregulatory Effects.....	23
4.3.5 Neurobiological Effects.....	23
4.3.6 Neuropsychological Effects .....	25
<b>4.4 Toxicology in Animals and Epidemiological Settings</b> .....	<b>26</b>
4.4.1 Single Dose Studies in Animals .....	26
4.4.2 Repeated Dose Studies in Animals .....	27
4.4.3 Genotoxicity .....	27
4.4.4 Carcinogenicity .....	27
4.4.5 Reproductive and Developmental Toxicity.....	27
4.4.6 Hyperthermia.....	30
4.4.7 Cardiovascular Toxicity .....	30
4.4.8 Hyponatremia .....	31
4.4.9 Hepatotoxicity .....	31
4.4.10 Neurotoxicity.....	32
<b>4.5 Serious Reports, Mortality, and Morbidity in Animals and Epidemiological Settings</b> .....	<b>34</b>
<b>4.6 Abuse Potential in Nonclinical Studies</b> .....	<b>38</b>
<b>5.0 Effects in Humans in Clinical Settings</b> .....	<b>39</b>
<b>5.1 History of Use in Clinical Settings</b> .....	<b>39</b>
<b>5.2 Pharmacology in Humans</b> .....	<b>41</b>
5.2.1 Pharmacokinetics .....	41
5.2.2 Pharmacodynamics.....	44
<b>5.3 Safety of MDMA in Humans</b> .....	<b>46</b>
5.3.1 Reproductive and Developmental .....	47
5.3.2 Immunological Effects .....	48
5.3.3 Thermoregulatory Effects .....	48
5.3.4 Cardiovascular Effects .....	50
5.3.5 Osmoregulatory Effects.....	56

5.3.6 Hepatic Effects .....	56
5.3.7 Neurobiological Effects.....	57
5.3.8 Neuropsychological Effects .....	58
5.3.8.1 Cognitive Function.....	59
5.3.8.2 Perceptual Effects .....	61
5.3.8.3 Social Effects .....	62
5.3.8.4 Emotional Effects.....	63
5.3.8.5 Suicidal Ideation, Behavior, and Depression .....	64
5.3.9 Adverse Events.....	73
5.3.9.1 Commonly Reported Adverse Events.....	73
5.3.9.2 Adverse Events .....	79
5.3.9.3 Serious Adverse Events .....	87
5.3.10 Abuse Potential .....	88
<b>5.4 Efficacy of MDMA Across Populations.....</b>	<b>89</b>
5.4.1 PTSD .....	89
5.4.2 Social Anxiety in Autistic Adults.....	91
5.4.3 Anxiety Associated with Life-Threatening Illness.....	91
<b>6.0 Summary of Data and Guidance for the Investigator .....</b>	<b>92</b>
<b>6.1 Pharmacology .....</b>	<b>92</b>
<b>6.2 Toxicology .....</b>	<b>93</b>
<b>6.3 Physiological Effects.....</b>	<b>95</b>
6.3.1 Immunological Effects .....	96
6.3.2 Hepatic Effects .....	96
<b>6.4 Suicidal Ideation, Behavior, and Depression .....</b>	<b>97</b>
<b>6.5 Adverse Events.....</b>	<b>97</b>
<b>6.6 Risk Mitigation in MDMA-Assisted Clinical Trials.....</b>	<b>98</b>
<b>6.7 Abuse Potential .....</b>	<b>100</b>
<b>7.0 Conclusion .....</b>	<b>101</b>
<b>8.0 References.....</b>	<b>102</b>
<b>9.0 Appendix.....</b>	<b>134</b>

## List of Figures and Tables

Table 1: Pharmacokinetic Constants for Plasma MDMA After Various Routes of Administration to Humans or Animals .....	15
Table 2: Summary of Published Morbidity and Mortality Reports .....	35
Figure 1: Metabolism of MDMA in Humans .....	43
Table 3: Pre-Drug, Peak, and Final Body Temperature During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations.....	50
Table 4: Pre-drug, Peak, and Final Systolic Blood Pressure During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations.....	52
Table 5: Pre-drug, Peak, and Final Systolic Blood Pressure During Experimental Sessions in Controlled Hypertension Subjects in MAPS-Sponsored PTSD Study MP-8.....	53
Table 6: Pre-drug, Peak, and Final Diastolic Blood Pressure During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations.....	54
Table 7: Pre-drug, Peak, and Final Diastolic Blood Pressure During Experimental Sessions in Controlled Hypertension Subjects in MAPS-Sponsored PTSD Study MP-8.....	54
Table 8: Pre-drug, Peak, and Final Heart Rate During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations .....	55
Table 9: List of All Clinically Significant Changes in Laboratory Values in Two Subjects from MP-2 .....	56
Table 10: Average ALT Values at Baseline and 2-Month Follow-up After Two Experimental Sessions in Subjects from MP-1 .....	57
Table 11: Neurocognitive Function - RBANS Mean Total Scores at Baseline, Primary Endpoint, End of Stage 1, and End of Stage 2 for MP-1, MP-4, and MP-12 as of 01 October 2015 .....	60
Table 12: Neurocognitive Function - PASAT Trial 1 and Trial 2 Mean Raw Total Scores at Baseline, Primary Endpoint, End of Stage 1, and End of Stage 2 for MP-1, MP-4, and MP-12 as of 01 October 2015.....	61
Table 13: Summary of Baseline Positive and Serious Responses on C-SSRS for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015 .....	65
Table 14: C-SSRS Positive and Serious Responses During Experimental Sessions and 1-Day Post-Drug for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015 .....	66
Table 15: C-SSRS Positive and Serious Responses During Experimental Sessions and 1-Day Post-Drug for Studies MAA-1 and MDA-1 as of 01 October 2015 .....	68
Table 16: C-SSRS Positive Responses During Telephone Contact Following Experimental Sessions for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015 .....	69
Table 17: C-SSRS Positive Responses at Endpoints After Treatment for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015.....	70
Table 18: Mean BDI-II Scores at Baseline, Primary Endpoint, and End of Stage 1 by Dose for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015.....	71
Table 19: Mean BDI-II Scores at Secondary Endpoint, End of Stage 2, and 12-month Follow-up for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015 .....	72
Table 20: Mean BDI-II Scores After MDMA or Placebo in MAA-1 as of 01 October 2015 .....	72
Table 21: Mean Percentage of Subjects Reporting Commonly Reported Reactions During MDMA or Placebo Treatment Collected from 12 Phase 1 Studies Conducted Outside of Sponsor Support .....	74
Table 22: Percentage of Observations of Most Commonly Reported Spontaneously Reported Reactions During Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015 .....	75
Table 23: Percentage of Observations of Most Commonly Reported Spontaneously Reported Reactions During Telephone Contact on Day 1-7 After Experimental Sessions in	

Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015 ..... 77

Table 24: Overview of All Adverse Events Post-Drug by Severity and Relationship in MAPS-Sponsored Studies Across Populations as of 01 October 2015 ..... 79

Table 25: Body Systems of All Adverse Events Post-Drug Reported by 2% or More of Subjects in MAPS-Sponsored Studies Across Populations ..... 81

Table 26: Related Adverse Events in Sponsor Supported Studies of MDMA-Assisted Psychotherapy Across Populations Organized by Body System as of 01 October 2015 ..... 84

Table 27: Severe Related Adverse Events in Sponsor Supported Studies of MDMA-Assisted Psychotherapy Across Populations as of 01 October 2015 ..... 87

Table 28: Serious Adverse Events in Sponsor-Supported Studies of MDMA-Assisted Psychotherapy Across Populations as of 01 October 2015 ..... 88

Table 29: Mean Global CAPS Scores in Stage 1 of Sponsor-Supported Studies of MDMA-Assisted Psychotherapy for PTSD as of 01 October 2015 ..... 90

Table 30: Mean Global CAPS Scores in Stage 2 and Long-term Follow-up of Sponsor-Supported Studies of MDMA-Assisted Psychotherapy for PTSD as of 01 October 2015 ..... 91

Table 31: Percentage of Observations of Spontaneously Reported Reactions During Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015 ..... 134

Table 32: Spontaneously Reported Reactions on Day 1-7 After All Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015 ..... 139

## List of Abbreviations

Ach	Acetylcholine
AE(s)	Adverse Event(s)
ALT	Alanine Aminotransferase
API	Active Pharmaceutical Ingredient
ARF	Acute Renal Failure
AVP	Arginine Vasopressin
BDI	Beck Depression Inventory
BDI-II	Beck Depression Inventory II
BDNF	Brain Derived Neurotrophic Factor
BOLD	Blood Oxygen Level Dependent
C	Celsius
CAPS	Clinician Administered PTSD Scale
CBF	Cerebral Blood Flow
cGMP	Current Good Manufacturing Practice
CNS	Central Nervous System
COMT	Catechol-O-methyltransferase
CPK	Creatine Phosphokinase
CRA	Clinical Research Associate
C-SSRS	Columbia Suicide Severity Rating Scale
CTproAVP	Stimulating Secretion of Copeptin
DAT	Dopamine Transporters
DEA	Drug Enforcement Administration
DBP	Diastolic Blood Pressure
DIC	Disseminated Intravascular Coagulation
DMF	Drug Master File
DNA	Deoxyribonucleic Acid
DSM-IV	Diagnostic and Statistical Manual of Mental Disorders IV
E	Embryonic Days
EEG	Electroencephalography
EKG	Electrocardiogram
ESR	Erythrocyte Sedimentation Rate
FDA	Food and Drug Administration
fMRI	Functional Magnetic Resonance Imaging
G-CSF	Granulocyte-colony Stimulating Factor
GD	Gestational Days
HHMA	3,4-Dihydroxymethamphetamine
HMA	4-Hydroxy-3-methoxy-amphetamine
HMMA	4-Hydroxy-3-methoxy-methamphetamine
HPA	Hypothalamus-pituitary-adrenal
HR	Heart Rate
IB	Investigator's Brochure
IL	Interleukin
IND	Investigational New Drug
LD50	Lethal Dose in 50% of Cases
LSD	d-Lysergic Acid Diethylamide
MAA-1	Phase 2 clinical trial of MDMA-assisted therapy for social anxiety in people on the autism spectrum
MAO	Monoamine Oxidase
MAO-A	Monoamine Oxidase A
MAOI	Monoamine Oxidase Inhibitor

MAPS	Multidisciplinary Association for Psychedelic Studies
MDA	3,4-Methylenedioxyamphetamine
MDA-1	Phase 2 clinical trial of MDMA-assisted psychotherapy for anxiety in relation to a life-threatening illness
MDE	Methylenedioxyethylamphetamine
MDMA	3,4-Methylenedioxymethamphetamine
MP-1	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Charleston, South Carolina
MP1-E2	Relapse study Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Charleston, South Carolina
MP-2	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Switzerland
MP-3	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Israel
MP-4	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Canada
MP-8	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Canada
MP-9	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Canada
MP-12	Phase 2 clinical trial of MDMA-assisted psychotherapy for PTSD in Canada
MRI	Magnetic Resonance Imaging
mRNA	Messenger Ribonucleic Acid
MT-1	Phase 1 clinical trial of MDMA-assisted psychotherapy for PTSD in healthy volunteers in Charleston, South Carolina
NET	Norepinephrine Transporter
NK	Natural Killer
NLP	Natural Language Processing
PASAT	Paced Auditory Serial Addition Task
PET	Positron Emission Tomography
PFC	Prefrontal Cortex
PMA	Paramethoxyamphetamine
PMMA	Paramethoxymethamphetamine
PND	Postnatal Day
PTSD	Posttraumatic Stress Disorder
RBANS	Repeatable Battery for the Assessment of Neuropsychological Status
rCBF	Regional Cerebral Blood Flow
SAE(s)	Serious Adverse Event(s)
SBP	Systolic Blood Pressure
SERT	Serotonin Transporter
SIADH	Syndrome of Inappropriate Antidiuretic-hormone Secretion
SNRI	Selective Serotonin and Norepinephrine Uptake Inhibitor
SPECT	Single Photon Emission Tomography
SSRI	Selective Serotonin Reuptake Inhibitor
SUD	Subjective Units of Distress
TNF- $\alpha$	Tumor Necrosis Factor-alpha
VHD	Valvular Heart Disease
VMAT2	Vesicular Monoamine Transporter 2
WBC	White Blood Cell Count
8-OH-DPAT	8-Hydroxy-2-(di-n-propylamino)tetralin

## 1.0 Summary

The Multidisciplinary Association for Psychedelic Studies (MAPS) is a U.S.-based non-profit research and educational organization supporting research of the therapeutic potential of 3,4-methylenedioxyamphetamine (MDMA). MAPS is sponsoring clinical trials of MDMA-assisted psychotherapy for patients with chronic disorders such as Posttraumatic Stress Disorder (PTSD), social anxiety associated with autism, and anxiety related to terminal illnesses. MDMA-assisted psychotherapy is an experimental treatment that combines psychotherapeutic techniques with administration of MDMA, a pharmacological adjunct that enhances aspects of psychotherapy. Prior to placement on the Drug Enforcement Administration's (DEA) list of Schedule I substances, MDMA was administered to thousands of people in psychotherapeutic practice outside of clinical trials. According to the 2011 United Nations World Drug Report, 11 to 28 million people aged 15 to 64 used Ecstasy, material represented as containing MDMA, around the world in various non-medical settings [1-5, 631]. The information presented in this Investigator's Brochure (IB) is summarized from published research studies of MDMA conducted by groups outside of the sponsor, sponsor collected data and published studies of Ecstasy use. For the purposes of this document MDMA will be used to refer to drug of known purity used in a controlled setting and Ecstasy will be used to describe drug-related information gathered from epidemiological settings.

MDMA is a ring-substituted phenethylamine also known as methylenedioxyamphetamine. MDMA is structurally similar, but functionally distinct, from amphetamines. MDMA is a chiral molecule, the sponsor uses racemic MDMA in the form of white crystalline powder compounded with inert material into capsules. The hydrochloride salt of MDMA is readily water soluble and once ionized is lipophilic. A substantial amount of data, both clinical and nonclinical, has been collected for over half a century of research on the physiological and psychological effects of MDMA in humans and animals. Estimates from animal data suggest a median lethal dose (LD50) in humans between 10 to 20 mg/kg [632]. Due to a wide range of responses to identical milligram per kilogram (mg/kg) dosing [7], the sponsor's human trials use fixed doses equivalent to between 1 and 4 mg/kg (active doses in studies range from 75 mg to 225 mg). Onset of MDMA effects occurs 30 to 60 minutes after oral administration [7, 8, 9], peak effects appear 75 to 120 minutes post-drug [10, 11, 12], and duration of effects lasts from 3 to 6 hours [10, 12, 13], with most effects returning to baseline or near-baseline levels 6 hours after drug administration. The elimination half-life of active doses is 8 to 9 hours [14].

The pharmacokinetics of MDMA in humans has been characterized using oral doses of up to 150 mg MDMA. MDMA disposition in the body follows nonlinear pharmacokinetics. As described in [Figure 1](#) (see [Section 5.2.1 Pharmacokinetics](#)), metabolism of MDMA results in *N*-demethylation to 3,4-methylenedioxyamphetamine (MDA). The parent compound and MDA are further *O*-demethylated to 3,4-dihydroxymethamphetamine (HHMA) and 3,4-dihydroxyamphetamine (HHA), respectively. Both HHMA and HHA are subsequently *O*-methylated mainly to 4-hydroxy-3-methoxy-methamphetamine (HMMA) and 4-hydroxy-3-methoxy-amphetamine (HMA). These four metabolites, particularly HMMA and HMA, are known to be excreted in the urine as conjugated glucuronide or sulfate metabolites [14].

MDMA is a triple monoamine reuptake inhibitor, and similar drugs in this class have been found to exert potent anti-depressant activity with a favorable safety profile in clinical trials [15, 16]. MDMA concomitantly promotes release, inhibits reuptake, and extends duration of serotonin, norepinephrine, and dopamine in the synaptic cleft to increase serotonergic, noradrenergic, and dopaminergic neurotransmission. MDMA has self-limiting subjective and physiological effects due to inhibitory activity on tryptophan hydroxylase [17-19], which prevents additional serotonin from being produced and released. This inhibition is reversible [20]. MDMA produces anxiolytic



and prosocial effects through release of the monoaminergic neurotransmitters, with the greatest effect on serotonin, followed by norepinephrine and dopamine [21-25]. MDMA has been shown to acutely decrease activity in the left amygdala and increase blood flow to the prefrontal cortex (PFC) in the brain [26-28]. MDMA has also been found to increase serum levels of the neurohormones oxytocin and arginine vasopressin (AVP) in humans [19, 29-33]. Some studies in healthy volunteers suggest that MDMA increases trust and attenuates reactivity to threatening cues, which are at least partially associated with oxytocin release [29, 34, 35]. The combined neurobiological effects of MDMA can increase compassion for self and others, reduce defenses and fear of emotional injury, and make unpleasant memories less disturbing while enhancing communication and capacity for introspection [36-39]. These factors taken together can provide the opportunity for a corrective emotional experience in the context of psychotherapy. Many of the therapeutic effects of MDMA-assisted psychotherapy are evident within a short period of treatment, often after the initial session.

Increased feelings of interpersonal closeness, changes in social perception and reduced anxiety may make MDMA a suitable pharmacological adjunct to enhance psychotherapy for anxiety disorders, such as PTSD and social anxiety in autistic adults [40]. MDMA may provide a much-needed option in the treatment of PTSD and anxiety associated with other conditions. Published results from MAPS study (MP-1) showed clinically and statistically significant improvements in PTSD severity in 20 per protocol subjects [41]. Findings from the long-term follow-up of MP-1 suggest that therapeutic benefits were sustained for an average of 41 months post-treatment [42]. The sponsor's second Phase 2 pilot study conducted in Switzerland (MP-2) demonstrated clinically significant improvements in PTSD symptoms, with results in the 125 mg MDMA dose group numerically but not statistically superior to the 25 mg MDMA dose group [43]. Long-term follow-up data 12 months later suggest that therapeutic benefits continued to increase in this subject population. There were no drug-related Serious Adverse Events (SAEs) or safety concerns in either study.

Data from MAPS studies and published literature show that MDMA produces sympathomimetic effects that include significant transient, self-limiting increases in heart rate (HR) and blood pressure that are likely to be well tolerated by healthy individuals [7, 9, 10, 12, 26, 44-46]. Most people do not experience elevations that exceed those seen after moderate exercise. These results were reproduced in MAPS Phase 1 safety study [47]. Risks posed by elevated blood pressure are addressed by excluding candidates with a history of cardiovascular, cerebrovascular disease, or with pre-existing uncontrolled hypertension and by regularly monitoring blood pressure and pulse throughout experimental sessions. Common reactions reported in the literature and clinical trials from MDMA are transient and diminish as drug effects wane during the session and over the next one to 7 days. The effects include lack of appetite, insomnia, dizziness, tight jaw or bruxism, difficulty concentrating, headache, impaired gait or balance, muscle tension, ruminations, feeling cold, and thirst (see [Section 5.3.9 Adverse Events](#)). MDMA is also a mild immunosuppressant [48]. Due to their limited duration, these sub-acute reactions are not likely to have clinical significance.

As of 01 October 2015, with 1180 individuals exposed to MDMA in controlled research settings (which includes 122 in MAPS-sponsored studies), there have been no unexpected drug-related SAEs to date, and expected SAEs have been rare and non-life threatening. As of the data cut-off, a single expected related SAE (increased ventricular extrasystoles), and 10 unrelated SAEs after drug administration have been reported in MAPS-sponsored clinical trials.

There have been a number of SAEs reported in individuals who use Ecstasy (material represented as containing MDMA, as defined above) around the world in various non-medical settings [1-5]. These include fatalities reported after Ecstasy and poly-drug use in unsupervised and uncontrolled

settings. These events are relatively rare given the prevalence of Ecstasy use, estimated to be in the millions worldwide [49, 50]. The most common adverse effects in Ecstasy and poly-drug use include hyperthermia, psychiatric problems, hepatotoxicity, and hyponatremia [51-55] (see [Section 4.4 Toxicology in Animals and Epidemiological Settings](#) and [4.5 Serious Reports, Mortality, and Morbidity in Animals and Epidemiological Settings](#)).

## 2.0 Introduction

MDMA is not a novel compound. The history of its use in humans predates controlled studies in healthy volunteers and clinical trials. MDMA was first synthesized and patented by Merck in 1912 [56] and is currently not covered by a patent. MAPS holds the Drug Master File (DMF) and an Investigational New Drug (IND) for MDMA with the U.S. Food and Drug Administration (FDA). After MDMA was rediscovered by the chemist Alexander Shulgin in 1976 [57], he and his colleagues provided initial reports of its pharmacology, with 80 mg to 160 mg MDMA required to produce desired subjective effects in humans [58, 59]. MDMA was found to robustly influence human emotional status in a unique way [59] without adversely effecting physiological functions or perception, such as visual perception or cognition [8, 11, 13].

MDMA possesses a complex pharmacological profile that is dominated by its effects as a monoamine releaser and reuptake inhibitor, with additional effects on limiting neurotransmitter production and degradation. Its prominent effects on serotonin differentiate it from amphetamine and methamphetamine, which primarily act to increase catecholamines such as norepinephrine and dopamine [21, 60]. In the Merck Index, MDMA resides in the Entactogen class [61]. Entactogens contain a ring-substituted amphetamine core, belong to the phenethylamine class of psychoactive drugs, and are described as promoting acceptance and compassion for self and others, changing recognition and response to emotions, and increased interpersonal closeness [19, 37, 62, 63]. In comparison to anxiolytics, antidepressants and atypical antipsychotics, MDMA does not require steady state levels in the blood to function as a catalyst to psychotherapy. Two to six administrations of MDMA, spaced approximately 1 month apart at active doses of 75 mg to 125 mg, may be an alternative to other medications that require daily dosing. This infrequent dosing regimen mitigates adverse event (AE) frequency and improves the risk/benefit ratio of MDMA, which may provide a significant advantage over daily dose medications.

Shulgin and Nichols were the first to report the effects of MDMA in humans [59]. MDMA-assisted psychotherapy first occurred during the mid-to-late 1970s after Shulgin introduced MDMA to a psychotherapist, Leo Zeff. Reported effects of MDMA include enhanced feelings of closeness to others, wellbeing, and insightfulness [64-66]. Prior to placement in Schedule I, MDMA was used in psychotherapy for individuals, couples, and groups to treat diverse psychological disorders, including moderate depression and anxiety [65, 67-69]. It was also found to be useful in reducing physical pain secondary to certain kinds of cancer [68]. No formal controlled clinical trials of safety and efficacy were conducted at the time [65, 70].

During the early 1980s, increasing numbers of people began using MDMA, sold as "Ecstasy" outside of therapeutic contexts [1]. The first wave of non-medical use occurred not only in dance clubs, but also in groups of people who used the drug in a self-exploratory or spiritual context. Non-medical use continues today in the same contexts [4, 71]. In the U.S., an estimated 800,000 people reported initiating Ecstasy use in the past year [72], and approximately 2.1 million Europeans between the ages of 15 and 64, or approximately 0.6% of the population, reported using Ecstasy in 2013 [73].

MDMA was added to the list of Schedule I controlled substances in the U.S. in 1985, defining it as a drug with a high potential for abuse and no accepted medical use [74, 75]. Classification as a Schedule I controlled substance, combined with the early research in animals and recreational users, hampered clinical research into the medical uses of MDMA until the 1990s. Shortly after it was scheduled, animal studies described long-term decreases in markers of serotonergic functioning after high or repeated doses of MDMA administration [76], however these were not relevant to doses in clinical trials [77, 78]. A recently published meta-analysis took careful steps to overcome methodological limitations in previous work, and found only modest indicators of long-term impairment in cognitive function in humans [53]. A systematic review of brain imaging studies in moderate ecstasy users found no convincing evidence for structural or functional changes [79]. Reports of AEs, such as hyperthermia, following Ecstasy use [80-82] and studies in Ecstasy users reporting changes in serotonin transporter (SERT) density, impaired memory and executive function raised concerns regarding the safety of MDMA administration [83-87]. However uncontrolled studies of Ecstasy use and preclinical animal studies that use inappropriately high doses of MDMA produce findings that are open to several interpretations [78, 88]. The vast majority of publications of Ecstasy users are retrospective reports in polydrug-users [53, 89].

While the initial studies in the 1990s conducted in humans examined the physiological effects of MDMA strictly from a safety perspective, current investigations have examined the effects on attention, prosocial effects, memory and brain activity, and human drug discrimination. Findings from an initial sponsor-funded study indicated that MDMA-assisted psychotherapy could be conducted safely in people with chronic PTSD who had failed first line treatments [90, 527]. This was repeated in a chronic, treatment-resistant PTSD sample in a sponsor-supported study (MP-1) [42] which demonstrated durable improvement in PTSD severity, with no difference in cognitive function between placebo and MDMA groups after an active dose of MDMA was given on two occasions, 1 month apart. In addition, placebo-controlled Phase 1 clinical trials confirmed that MDMA produces an easily controlled intoxication characterized by euphoria, increased well-being, sociability, self-confidence, extroversion, transient increases in anxiety, and minor alterations in perception [8, 10-12, 29, 30, 35, 91, 92].

MAPS is completing Phase 2 investigations of MDMA-assisted psychotherapy. Significant durable improvement in PTSD symptoms lasted for at least 12 months after MDMA-assisted psychotherapy in two completed studies (MP-1, MP-2) [42, 43]. There are four Phase 2 studies for treatment of PTSD that have completed treatments and are in follow-up: two studies in the U.S. (MP-8, MP-12), one in Canada (MP-4), and one in Israel (MP-9). Data from Phase 2 studies will be submitted to FDA for an End-of-Phase 2 meeting to support an application for Phase 3 multi-site MDMA/PTSD research studies. Based on the current state of scientific knowledge and the risk/benefit profile of active doses of MDMA, it appears favorable to continue the research of MDMA as an adjunct to psychotherapy.

Based on clinical experience with PTSD, MAPS is exploring new indications for this treatment. Studies for additional indications include one Phase 2 study (MAA-1) of MDMA-assisted therapy for social anxiety in people on the autism spectrum and one study of MDMA-assisted psychotherapy to address anxiety associated with a life-threatening illness (MDA-1). In addition, there is one ongoing Phase 1 study of MDMA-assisted psychotherapy to assess psychological effects in healthy volunteers (MT-1). When completed, this will be the first Phase 1 investigation to assess acute effects in a therapeutic setting that is comparable to MDMA-assisted psychotherapy studies for PTSD.

This IB will present preclinical and clinical studies of MDMA, as well as epidemiological studies in Ecstasy users.

### 3.0 Physical, Chemical, and Pharmaceutical Properties and Formulation

MDMA is structurally similar, but functionally distinct, from amphetamines and mescaline. MDMA, also known as 3,4-methylenedioxy-N-methylamphetamine and N-methyl-3,4-methylenedioxyamphetamine, has the chemical formula of  $C_{11}H_{15}NO_2$ . It was first synthesized as a precursor of a haemostatic drug called methylhydrastinine as a phenylisopropylamine derivative of safrole, an aromatic oil found in sassafras, nutmeg, and other plants [56].

MDMA is a chiral molecule, possessing two enantiomers, S(+)-MDMA and R(-)-MDMA, with S(+)-MDMA being more potent than R(-)-MDMA [6, 58]. Research in humans to date and the majority of nonclinical studies have used racemic MDMA, or an admixture containing equal amounts of both enantiomers. Studies of drug discrimination in rodents [94, 95] and studies of self-administered and experimenter-administered MDMA enantiomers in primates [23, 96-99] suggest that MDMA enantiomers may produce different physiological and rewarding effects, and there may be some synergy between the two when administered as a racemate. It seems that R(-)-MDMA may have hallucinogen-like effects, compared to S(+)-MDMA, which exhibits psychomotor stimulant-like effects. Findings comparing the effects of the enantiomers of the related compound methylenedioxyethylamphetamine (MDE) suggest that these different effects of MDMA enantiomers may occur in humans [100]. According to an *in vivo* microdialysis study in rodents, S(+)-MDMA may be associated with greater dopamine release in specific brain areas [101]. A study conducted in 2014 in monkeys found that S(+)-MDMA, but not R(-)-MDMA, significantly increased extracellular dopamine levels in the dorsal striatum, whereas S(+)-MDMA significantly increased serotonin levels [23]. *In vitro* studies reported greater binding at a specific alpha nicotinic acetylcholine (ACh) receptor by R-MDMA compared with S-MDMA [102]. MDMA available for humans in clinical trials is racemic, containing roughly equal amounts of both enantiomers. Any differential effects of the enantiomers remain untested in humans. The sponsor will use racemic MDMA in all current and planned studies. Unless otherwise stated, MDMA is used throughout this document to refer to the racemic mixture.

For clinical trials, the sponsor used racemic hydrochloride salt of MDMA from two sources. Since this is the formulation used in all prior investigations in humans, the sponsor will continue to use the hydrochloride salt of MDMA. The hydrochloride salt of MDMA is readily water soluble with a  $pK_a$  of 9.9 [103], which influences whether it is ionized in plasma and slightly reduces its ability to cross into oral fluid. MDMA is also more lipophilic, which drives it into oral fluid, and may influence its ability to pass the blood brain barrier and influence signaling in the central nervous system (CNS) [104].

Sponsor-supported studies in the U.S. use MDMA manufactured in 1985 by David Nichols, Ph.D., at the Department of Medicinal Chemistry and Pharmacology, Purdue University, West Lafayette, IN. The MDMA was manufactured as a single lot for use in federally approved clinical research. A stability analysis conducted in 2006 indicates that the compound remains highly stable and pure after 21 years of storage [105]. Studies conducted outside of the U.S. use MDMA from a single batch manufactured in 1998 by Lipomed AG in Arlesheim, Switzerland. The most recent analysis of drug stability and purity conducted on February 2, 2010 confirmed that this MDMA is 99.9% pure with no detectable decomposition. For sponsor-supported studies, MDMA in the form of white crystalline powder is compounded with inert material into capsules. Capsules are administered orally with a glass of water.

The sponsor has contracted with Shasun, a manufacturer in the United Kingdom, to manufacture active pharmaceutical ingredients (APIs) to produce 1 kg of MDMA following current Good

Manufacturing Practices (cGMP). The material is planned for use in all Phase 3 studies. Details of manufacturing are available from the manufacturer upon request.

MDMA doses in sponsor-supported studies are fixed within a therapeutic dose range, rather than based on body weight, based on epidemiological information and lack of linear dose response with behavioral effects in Phase 1 and sponsor-supported studies [7]. A typical active dose is 125 mg, which is equivalent to 2 mg/kg for the initial dose. The optional supplemental dose of 62.5 mg is equivalent to 1 mg/kg, for a cumulative dose of 3 mg/kg. Various comparator and active doses of MDMA are also being tested in the clinical trials.

MDMA does not require special conditions for storage. The capsules are stored in sealable containers placed within a dark safe at ambient temperature. MDMA is a Schedule I compound and is stored and handled in compliance with relevant federal and state regulations. In accordance with the requirements of the U.S. DEA and international drug regulatory authorities, license holders will be responsible for storing and dispensing the MDMA, and ensuring it is stored under appropriate protections, often in a floor-mounted safe.

Lactose is used as inactive placebo and as an inactive filler intended to maintain blinding by creating capsules of equal weight. Lactose has been in use as an inactive material of similar appearance and was selected because it can be safely consumed by most people and is inactive. Whenever conducting blinded studies, the sponsor will continue to employ lactose or inactive materials that exist as white powders without significant odor that can be safely administered in humans. The purpose of this excipient is solely to permit placebo or active placebo administration under blinded conditions.

#### 4.0 Nonclinical Studies

Findings from nonclinical animal research, retrospective studies of Ecstasy use and case reports of Ecstasy use in humans are presented. Research into the pharmacological, physiological, or psychological effects of MDMA began in the 1950s, when the U.S. Army administered MDMA to guinea pigs, monkeys, mice, rats, and dogs as part of a military research program, possibly intending to develop chemical incapacitants or means of enhancing interrogation [106]. Investigations of the pharmacology, functional effects, and toxicity of MDMA in animals have generally included injections of large and often repeated doses of MDMA that are not human-equivalent doses. Studies of MDMA have been conducted in primates and rodents. Primate species studied include baboon, macaque, rhesus monkey, and squirrel monkey, and rodents include mice and rats. Studies of circadian rhythm have occurred in hamsters. Beginning in the mid-2000s onwards, reports re-examining these effects have questioned the applicability of interspecies scaling models for MDMA, and have supported nonlinear pharmacology [78, 107, 108]. In general, doses in the range of 1 to 5 mg/kg in animals are relevant to human research and are described in more detail in [Section 4.2.2 Pharmacodynamics in Animals](#). Findings in doses above this that show a toxic effect are described when relevant in [Section 4.4 Toxicology in Animals and Epidemiological Settings](#).

Evidence exists for intentional human use of MDMA, known as Ecstasy among other names, as early as the late 1960s [57], and there are records of a police seizure of MDMA in the early 1970s [109]. MDMA was administered to thousands of people prior to scheduling and many continue to use Ecstasy around the world in various non-medical settings [1-5]. In this IB, “Ecstasy” (or other common names) refers to material assumed to be MDMA used in naturalistic settings (see [epidemiology sections](#)), however when used in these uncontrolled settings the drug may not contain only or any MDMA. One of the problems in assessing the effects of Ecstasy in users is determining the purity and identity of the substance. It may contain other substances along with



or instead of MDMA, and when present, the amount of MDMA can vary widely [110-112]. The majority of studies rely on self-reported use and do not attempt to confirm that material used is MDMA. Synthesis of MDMA is relatively simple, and is often produced illegally in laboratories with no quality control, these synthesized tablets also may be cut or mixed with other psychoactive substances. Substances found mixed with MDMA have included amphetamine, methamphetamine, dextromethorphan, paramethoxymethamphetamine (PMMA), paramethoxyamphetamine (PMA), cathinones, ketamine, caffeine, and ephedrine. Retrospective studies in Ecstasy users are described in [Section 4.3 Physiological Effects in Epidemiological Settings](#) and case reports of morbidity and mortality in Ecstasy users are included in [Section 4.5 Serious Reports, Mortality, and Morbidity in Animals and Epidemiological Settings](#) to provide the context of potential safety information of a related compound to MDMA which has extensive use outside of a research setting.

## 4.1 Nonclinical Pharmacology

MDMA possesses a complex pharmacological profile that is dominated by its effects as a monoamine releaser and reuptake inhibitor. Its prominent effects on serotonin differentiate it from amphetamine and methamphetamine, which primarily act on dopamine and norepinephrine [21, 60]. In the following sections, the pharmacology of MDMA is presented based on nonclinical animal studies and epidemiological studies.

## 4.2 Pharmacology in Animals

### 4.2.1 Pharmacokinetics in Animals

MDMA is metabolized via two hepatic pathways. In the major pathway in rats, MDMA is *O*-demethylated by cytochrome P450 CYP2D1 and 3A2 to form HHMA, which is *O*-methylated to generate HMMA by catechol-*O*-methyltransferase (COMT). In the minor pathway in rats, MDMA is *N*-demethylated by CYP1A2 and 2D1 to form MDA, which is an active metabolite. MDA is *O*-demethylated by the same enzymes as MDMA, with subsequent metabolism by COMT. Metabolites of MDMA are excreted in urine as glucuronide and sulfate conjugates. MDMA and metabolites have shorter half-lives in rats than humans at comparable doses based on plasma  $C_{max}$  values. Rats tend to form more MDA and glucuronide-conjugated metabolites than humans [113]. As MDMA dose increases above 2.5 mg/kg s.c. or i.p. in rats, a larger percentage of the administered dose is shunted to the *N*-demethylation pathway, resulting in greatly enhanced formation of MDA [114]. Comparison of metabolic pathways between rats and mice given 10 mg/kg intraperitoneal (i.p.) MDMA indicate that 49.1% of MDMA is metabolized through the HMMA pathway in mice versus 72% in rats, and 18.3% of MDMA is metabolized through the MDA pathway in mice versus 28% in rats based on AUC ratios to MDMA. MDMA at 10 mg/kg was also found to be eliminated more rapidly in mice (0.4 hours, i.p.) than rats at (1.1 hours, subcutaneous (s.c.)) [78, 115].

To address questions of the applicability of interspecies scaling models and nonlinear pharmacology of MDMA, a study examining MDMA and metabolites in rats given 2.5, 5, and 10 mg/kg s.c. found that MDMA metabolism is nonlinear in rats, with 2.5 mg/kg producing plasma  $C_{max}$  levels approximating those seen in humans receiving between 75 and 100 mg [14, 114, 116]. Injections of 2 mg/kg s.c. or i.p. in rats were found to be similar to oral administration of 100 mg MDMA in humans based on plasma MDMA and metabolite concentrations [78]. Based on plasma values, a dose of 3 mg/kg i.p. MDMA administered in mice was comparable to a single oral dose of 100 mg in humans [94]. Studies in rats and mice provide compelling evidence of nonlinear pharmacokinetics, likely due to saturation of metabolic enzymes, determined by greater

than expected AUC values for MDMA and MDA after subsequent MDMA doses, while AUCs for HHMA and HMMA remained lower than expected [114, 115].

Single dose pharmacokinetics of oral 7.4 mg/kg MDMA in squirrel monkeys shows two to three-fold higher plasma MDMA concentrations than humans receiving an oral dose of 100 mg, although allometric interspecies scaling predicts an equivalent dose [107]. A study directly comparing MDMA pharmacokinetics in humans and monkeys found that the two species metabolized MDMA in a similar but not identical manner - MDMA half-life in monkeys was less than half the duration seen in humans (1.1 hours at a dose of 2.8 mg/kg in squirrel monkeys versus 8.4 hours after 1.5 mg/kg in humans). Both monkeys and humans exhibit nonlinear pharmacokinetics [14, 118, 119], and it appears they exhibit similar plasma MDMA levels after receiving the same dose of MDMA [119, 120]. These pharmacokinetic findings suggest that nearly all toxicological and behavioral preclinical studies of MDMA use overestimated doses that exceed human-equivalent doses by 2.7 to 10.7 times, depending on route of administration, due to both simple dose conversion and allometric scaling. As a consequence, it is difficult to interpret the relevance of findings in preclinical studies employing these dosing regimes.

**Table 1: Pharmacokinetic Constants for Plasma MDMA After Various Routes of Administration to Humans or Animals**

	$C_{\max}$ (ng/ml)	AUC (h•ng/ml)	$T_{\max}$ (h)	$t_{1/2}$ (h)	References
<b>Rat<sup>A</sup></b>					
2 mg/kg i.p.	210±108	163±56	0.14±0.08	0.80±0.16	[78]
2 mg/kg s.c.	196±50	304±65	0.75±0.29	0.79±0.14	[78]
2 mg/kg p.o.	46±15	61±42	0.56±0.31	0.77±0.11	[78]
2.5 mg/kg s.c.	164.1±47.1	272.1±71.6	0.6±0.2	1.1±0.9	[114]
5 mg/kg s.c.	370.8±41	879.1±133.2	0.9±0.6	0.9±0.1	[114]
10 mg/kg s.c.	893.9±90.7	2879.9±491.5	1.1±0.4	2±0.6	[114]
<b>Mouse<sup>B</sup></b>					
3 mg/kg i.p. <sup>C</sup>	369.8	---	0.17	0.6	[94]
10 mg/kg i.p.	1109±87	1233±53	≤0.3	0.4	[115]
20 mg/kg i.p.	2152±82	2611±86	≤0.3	0.6	[115]
<b>Squirrel Monkey</b>					
1.4 mg/kg p.o.	100.2±51.5	340.3±248.4	1±0.4	1.8±0.9	[121]
2.8 mg/kg p.o.	312.7±92.8	1314.2±581.5	1.1±0.4	2.1±0.8	[121]
5.7 mg/kg p.o.	723.6±228	3866.2±891	1.3±0.9	2.6±0.7	[121]
10 mg/kg p.o.	1594.5±295.6	12,839.2±2144.6	1.3±0.9	4.2±1.5	[121]
7.4 mg/kg s.c.	1227±167	5006±528	---	3.5±0.9	[107]
7.4 mg/kg p.o.	773±157	3408±821	---	3.1±0.5	[107]
<b>Human</b>					
1.0 mg/kg p.o.	147±10	1389±119	2.3±0.2	7.2±0.6	[122]
1.6 mg/kg p.o.	292±76	3485±760	2.4±0.6	8.1±2.1	[116]
1.6 mg/kg p.o.	254.7±60.4	3070.6±673.4	2.4±0.6	8.4±1.6	[119]
2.0 mg/kg p.o.	442-487	5133-5232	1.5-2.0	6.9-7.2	[14]

<sup>A</sup> Male Sprague-Dawley rats

<sup>B</sup> Male FVB mice

<sup>C</sup> Fantegrossi et al. reported mean pharmacokinetic parameters of R(-)-MDMA and S(+)-MDMA after administering racemic MDMA. In this table, plasma racemic  $C_{\max}$  values estimated by taking sum of R(-) and S(+), while  $T_{\max}$  and  $t_{1/2}$  presented as an average of the enantiomers' values.

#### 4.2.2 Pharmacodynamics in Animals

Most effects of MDMA likely arise directly from monoamine reuptake inhibition and release, and indirectly from activation of downstream monoamine receptors and subsequent secretion of

neuromodulators oxytocin and AVP. MDMA binds primarily to membrane-bound monoamine transporters, which remove monoaminergic neurotransmitters from the space between neurons, known as the synaptic cleft. MDMA appears to alter the conformation of the transporters, enabling monoamines to diffuse out of the neuron rather than being actively transported into the presynaptic neuron [60, 123, 124]. MDMA prevents the reuptake of serotonin, and to a lesser extent, norepinephrine and dopamine, and facilitates release of these neurotransmitters [60, 125-127]. The selectivity of MDMA for specific monoaminergic neurotransmitters is species-dependent, and cannot solely be attributed to differences in binding affinity for specific reuptake transporters observed *in vitro*, as described below. In *in vitro* studies, MDMA was also found to compete with monoamines for sites on the vesicular monoamine transporter-2 (VMAT2), suggesting MDMA also promotes active release of monoamines from vesicular stores, in addition to inhibiting reuptake [128-130].

MDMA can inhibit monoamine oxidase A (MAO-A) *in vitro* at high concentrations, which preferentially degrades serotonin, and leads to accumulation of extracellular serotonin in the synaptic cleft [131, 132]. Inhibition of MAO-A may have played a role in fatalities and medical emergencies seen after combining Ecstasy with MAO inhibitors in epidemiological settings [133, 134]. Spurred on by prior reports hypothesizing that apparent greater serotonergic toxicity of MDMA in primates, as compared to rodents, could be attributed to greater SERT affinity [135], researchers specifically examined affinity in cells transfected to express human monoamine transporters [127, 136]. These studies found that even though binding affinity of MDMA for the human norepinephrine transporter (NET) exceeded the affinity for SERT and dopamine transporters (DAT), serotonin was preferentially released over norepinephrine and dopamine [127], which may account for primarily serotonergic effects of MDMA. On the other hand, in rodents MDMA affinities for transporters are ordered as SERT>NET>DAT [137]. MDMA does not have as strong an affinity for the DAT as methamphetamine [21].

The ability of MDMA to stimulate release of pre-synaptic serotonin, norepinephrine, and dopamine in multiple brain regions and inhibit reuptake has been well documented [138]. *In vivo* microdialysis and voltammetry results show significant enhancement of serotonin, and to a lesser extent dopamine following MDMA administration, a response attenuated by various transporter inhibitors. MDMA-stimulated serotonin and dopamine release has been measured in the striatum, nucleus accumbens, PFC, and the hippocampus of freely moving rats [139-142]. In addition, enhancement of Ach release has been demonstrated in the PFC, striatum, and hippocampus by both a dopaminergic and serotonergic dependent mechanism [143, 144]. The subjective and physiological effects of MDMA are produced by the dynamic interaction of these transmitter systems on numerous brain networks that modulate learning and memory, emotion, reward, attention, sympathetic/parasympathetic activity, and neuroplasticity.

In addition to carrier-mediated monoamine release, MDMA has affinity *in vitro* for specific serotonin, norepinephrine, Ach, and histamine receptors, although the concentrations tested may not translate to standard human MDMA doses [24, 145-147]. An *in vitro* binding study comparing MDMA with a number of drugs that include cathinone derivatives suggests that contrary to an earlier report of low affinity for 5HT<sub>2A</sub> serotonin receptors, MDMA may have significant effects at the receptor [25]. MDMA likely modulates 5HT<sub>1A</sub> serotonin receptors indirectly through serotonin release, though it is possible that MDMA may also act as a partial 5HT<sub>1A</sub> agonist in some brain areas [148]. Findings from other studies suggest that MDMA shares qualities with 5HT<sub>1A</sub> agonists. Early studies in rats suggest that pharmacological activation of 5HT<sub>1A</sub> receptors reduce anxiety and aggression [149, 150], and some drug discrimination studies suggest that the 5HT<sub>1A</sub> agonist 8-hydroxy-2-(di-n-propylamino)tetralin (8-OH-DPAT) partially or fully substitutes for MDMA [151-153]. In addition to its primary effects, both enantiomers of MDMA enhance Ach release in the PFC [144, 154] and promote changes in GABA-ergic systems



that correlate with sociability [155]. At least some direct or indirect effects of MDMA on serotonin receptors may alter GABA uptake in the ventral tegmental area of rats [156]. An *in vitro* study found that S-MDMA showed signs of competitive interaction with the alpha-4 beta-2 nicotinic receptor which are implicated in learning [157], while R-MDMA did not produce this effect [102].

Infusion of serotonin in the rat brain stimulates secretion of oxytocin into peripheral blood via activation of 5HT<sub>1A</sub>, 5HT<sub>2C</sub>, and 5HT<sub>4</sub> receptor subtypes, as well as AVP secretion via activation of 5HT<sub>2C</sub>, 5HT<sub>4</sub>, and 5HT<sub>7</sub> receptor subtypes [158]. MDMA was shown to increase oxytocin and AVP secretion in rats [159] through a 5HT<sub>1A</sub> mechanism [160]. Administering a 5HT<sub>1A</sub> antagonist attenuates the prosocial behavior of rats, measured by preference to lie adjacent to each other, possibly because it prevents elevation in oxytocin [160, 161]. MDMA also promotes norepinephrine release through reuptake inhibition, which is an additional pathway that can contribute to oxytocin secretion and may control emotion regulation. Both oxytocin and AVP are implicated in the widespread regulation of behavioral aspects of mood and also act on different target organs to modulate physiological functions in the periphery [162]. Taken together, MDMA has been shown to have a diverse array of pharmacodynamic effects in animals, with findings of interest presented below by topic.

#### 4.2.2.1 Stable Effects on Gene Expression in Animals

Epigenetic modifications, including deoxyribonucleic acid (DNA) methylation, demethylation, and histone acetylation, are thought to be involved in dynamic regulation of memory reconsolidation in the adult nervous system and play a role in memory formation [163]. Early childhood adversity and trauma is associated with transcriptional silencing of the brain-derived neurotrophic factor (BDNF) gene through DNA methylation, which can either be a risk factor in development of PTSD or a result of having PTSD in adulthood [164]. In a 2015 report, MDMA showed DNA hypermethylation and hypomethylation activity in cardiac tissue by microarray analysis in mice [165], and this activity may extend the CNS. Epigenetic effects on BDNF and other gene expression is a hypothesized mechanism by which MDMA in combination with training in animal studies modeling anxiety disorders, or psychotherapy in humans, exerts its therapeutic effects.

A number of research teams have studied the effects of MDMA on gene expression in rodents [166-169]. However, many of these reports used 10 to 20 mg/kg MDMA, a dose range that is 5 to 10.7 times greater than the 1.5 to 2 mg/kg doses employed in human trials, making it less likely that these changes can be generalized to humans given lower doses. However, even at these doses toxicity was not observed, and a self-administration study at clinically relevant doses reproduced findings of elevation of genes such as serum/glucocorticoid kinase 1 and 3 (*Sgk1*, *Sgk3*), which regulate glutamatergic signaling and are associated with neuroplasticity and learning, as well as processes involved in memory consolidation in serotonergic neurons [170]. These studies also report an increase in expression of genes that regulate the GABA transporter [166], which is expressed in GABA-ergic neurons indirectly regulated by glutamatergic afferent neurons. Serotonin-transporter knockout mice did not display some of these changes in gene transcription, suggesting that serotonin release is required for this activity [166]. In the acute period 24 to 48 hours after MDMA exposure, a study in rats found 33 to 70% upregulation of BDNF messenger ribonucleic acid (mRNA) transcripts in the frontal cortex, with a time-dependent decrease, up to 73%, of BDNF transcripts in the hippocampus [171]. The frontal cortex and hippocampus are both regions known to play a causal role in memory retrieval and reconsolidation in animals and humans [172], mediated in part through GABA-ergic signaling [173], suggesting that these transcriptional changes may be functionally related.

Examining rat brains after repeated MDMA administration for 2 weeks detected a sharp drop in SERT expression [174], suggesting a compensatory downregulation in response to repeated high doses of MDMA. A study in rats found repeated administration of MDMA at 1 or 5 mg/kg weekly for 4 weeks increased transcripts for 5HT<sub>1B</sub> receptors in various brain regions and 5HT<sub>2C</sub> receptors in the cortex and hypothalamus, likely due to serotonin depletion and subsequent need to increase serotonin receptor availability [175]. Increased levels of gene transcripts regulating extracellular signaling in mice were also reported after MDMA [176]. Serotonin may play a more significant role than dopamine in transcription changes mediated by MDMA [175]. Mouse brains examined 8 hours after 8 days of self-administration or non-contingent administration detected increased transcription of genes related to inflammation and immune modulation in both groups and transcription of genes related to neuroadaptation in mice self-administering MDMA [170]. Transcripts in these studies were assessed 8 to 10 hours after the last of repeated MDMA administrations and it is unclear whether these changes reflect residual acute effects of the MDMA or changes related to repeated MDMA administration. In addition, changes in transcription do not always correlate with functional consequences in proteins levels. BDNF has been shown to have multiple functionally distinct splice variants which have tight temporal and spatial control in an activity-dependent, stimulus-specific manner [177]. However, MDMA produces a durable enhancement of fear extinction in mice, an effect mediated by an MDMA-associated increase in BDNF expression specifically in the context of fear extinction training, suggesting that gene expression changes after MDMA are functionally relevant [178].

#### 4.2.2.2 Immunological Effects in Animals

MDMA acts as a mild immunosuppressant in rodents. MDMA administration at 5 mg/kg in rats is associated with impaired macrophage activity as evidenced by inhibition of Tumor Necrosis Factor-alpha (TNF- $\alpha$ ) secretion for 12 hours post-drug [179]. In mice injected with 10 mg/kg MDMA for 5 days, increases in epithelial tissue of cytokines interleukin 1-alpha (IL-1 $\alpha$ ), granulocyte-colony stimulating factor (G-CSF), and interleukin 3 (IL-3) were found, while decreased serum levels of many cytokines were reported [180]. MDMA decreased neutrophil oxidative bursts and phagocytosis, and increased the number of circulating neutrophils while decreasing the number of lymphocytes. MDMA also increased hypothalamus-pituitary-adrenal (HPA) axis activity through a noradrenergic pathway in the hypothalamus [181]. MDMA also suppresses interferon- $\gamma$  secretion and signaling in mice [182]. Interestingly, MDMA was shown to reduce inflammation and airway reactivity in a mouse model of allergic asthma, suggesting that MDMA could have beneficial immunomodulatory effects in cases of heightened inflammation [183]. This constellation of findings was in the 10 mg/kg dose range, which calls to question the applicability to moderate dosing regimens. However, a microarray study found that mice self-administering MDMA at moderate doses had transcriptional changes in many genes related to immune and inflammatory responses as well as neuroplasticity and learning [170], suggesting that immunosuppressant effects of MDMA at clinically relevant doses could be beneficial in the treatment of psychoneuroimmunological disorders such as PTSD [184].

#### 4.2.2.3 Thermoregulatory Effects in Animals

Rodents have generally been used to study the hyperthermic effects of MDMA. Rodents have a much smaller body mass and do not perspire, but use their tail to regulate body temperature which has a large surface to volume ratio, and is perfused with many blood vessels for thermoregulation. Since thermoregulation is different in rodents and humans [185], findings may have limited applicability to humans. MDMA doses that are moderate to high elevate body temperature and disrupt thermoregulation in mice [124], and doses of MDMA in the 1 to 2 mg/kg range only cause a slight increase in body temperature [186]. Rats given doses of 10 mg/kg MDMA (s.c. and i.p.), but not 2 mg/kg, experienced increases in body temperature correlated

with levels of the active metabolite MDA [78, 114]. A study of rats receiving subcutaneous injections of 1 and 3 mg/kg MDMA demonstrated minimal effect on brain hyperthermia using thermal couplers installed in the nucleus accumbens, however ambient temperatures of 29°C and social interaction had a potentiating effect on body temperature and malignant hyperthermia at higher doses [187], described in [Section 4.4 Toxicology in Animals and Epidemiological Settings](#). MDMA effects on body temperature are susceptible to changes in ambient temperature in rodents, with high ambient temperature significantly increasing body temperature in mice and rats, and low ambient temperatures reducing it [188-190]. The MDMA-induced impairment in thermoregulation is caused, at least in part, by peripheral vasoconstriction in the tail, an effect mediated by brain neurotransmitter activity [191, 192].

High doses of MDMA also produce significant elevations in body temperature in primates [107, 193, 194]. At doses closer to those humans ingest [195], monkeys exhibit only slight to moderate elevation in body temperature [196, 197]. In contrast to findings in rodents, primates are not susceptible to changes in ambient temperature when given MDMA, exhibiting slight to moderate increases in body temperature regardless of the temperature of the environment [195-197], though at least one study found that the ambient temperature influenced the effects of 1.5 mg/kg i.v. MDMA on body temperature in monkeys, with lower body temperatures seen after MDMA administered in cool temperatures and higher body temperatures in another group given MDMA at warm temperatures [198]. Findings in rodents do not extrapolate well to primates in this area. Given that the thermoregulatory effects in rodents are highly dose-dependent, the majority of physiological effects seen after low to moderate MDMA administration suggest that a controlled environment and moderate doses would be sufficient to mediate physiological complications associated with hyperthermia, including cardiovascular, osmoregulatory, neurological, and immunological effects.

#### 4.2.2.4 Cardiovascular Effects in Animals

*In vivo* assessments of cardiovascular effects of MDMA in animals detected increased sympathomimetic activity, as seen in humans [124, 199]. An injection of 2 mg/kg MDMA elevated heart rate in rabbits [633]. Ten mg/kg MDMA produced a relatively larger increase in heart rate in rats than blood pressure, an effect possibly controlled by beta adrenergic receptors [199]. The researchers found that MDMA has both pressor and depressor effects, acting through adrenergic receptors [201-203]. Another study in rodents also suggests that norepinephrine may play a role in cardiovascular effects [204], findings that have been more intensively investigated in humans [205-208]. Given the affinity of MDMA for the NET, it is possible that the cardiovascular effects of MDMA could be attributed to norepinephrine signaling in the peripheral nervous system.

#### 4.2.2.5 Osmoregulatory Effects in Animals

AVP is a key regulator of water balance in the body, and has antidiuretic actions when acting at its V2 receptor subtype in the kidneys [209, 210]. MDMA can influence water regulation by activation of the AVP system, as shown in several animal studies. A study of isolated *in vitro* rat hypothalamus initially reported AVP and oxytocin release after MDMA and its metabolite HMMA [33]. *In vivo* drug-discrimination studies in rats suggest that AVP receptors are involved in producing interoceptive effects of MDMA [162]. When 10 mg/kg i.p. MDMA was administered at 30°C ambient temperature to male Wistar rats, MDMA induced expression of Fos, a marker of neural activation, in the supraoptic nucleus, a brain region important for osmoregulation and a key mediator of oxytocin and AVP release [211]. This finding suggests that MDMA can have osmoregulatory effects in rats at high doses administered at warm ambient temperatures.

#### 4.2.2.6 Neurobiological Effects in Animals

It appears that single doses of MDMA (2.5 mg/kg i.p. in monkeys, 7.5 mg/kg i.p. in rats), approximately four times a human equivalent dose, reduces brain serotonin production for 2 weeks or more [107] but does not increase validated markers of neurotoxicity associated with neurodegeneration [108]. Monkeys allowed to self-administer MDMA for an 18-month period had no reductions in brain dopamine, slight reductions in brain serotonin, and no chemical markers of neuronal injury [509]. One report detected a reduction in N-acetylaspartate to creatine ratio, which the authors considered a sign of neuronal injury, although no decreases in brain serotonin were detected after administration of two human-equivalent doses of MDMA to marmoset monkeys for 2 days [213]. A study examining the rat hippocampus reported indications of apoptosis after 5 or 10 mg/kg given daily for 1 week but not after 2.5 mg/kg [214]. Doses of 10 mg/kg administered s.c. and i.p., but not 2 mg/kg, produced signs of serotonin syndrome in rats, but neither dose reduced total serotonin levels in the brain 2 weeks after drug administration. Serotonin syndrome is defined as an excess of serotonin in the CNS causing a suite of specific signs and symptoms that can require intervention [215-217]. Serotonin syndrome severity correlates with MDMA plasma concentrations [78]. Taken together, MDMA doses up to 2.5 mg/kg appear to alter regulation of serotonergic signaling in the rat brain without producing damage to serotonin axons, based on transient reductions in brain serotonin and SERT levels, in the absence of indicators of neuronal injury or decreased expression of the SERT gene [88].

In rats, large doses of MDMA (10 or 20 mg/kg) elevated serum corticosterone (a rodent cortisol analog) and prolactin [218-220], with elevation lasting up to 4 hours after dosing, and with hormone levels attenuated by a 5HT<sub>2A</sub> serotonin receptor antagonist. Given the dosage used was five to 10.7 times larger than an active dose in humans, it is unclear if this response is analogous to elevated cortisol in humans or whether it reflects a different process. Administering 1 to 3 mg/kg doses found that R(-)-MDMA, but not S(+)-MDMA, significantly increased prolactin levels in rhesus monkey plasma, suggesting that at least the R(-) enantiomer of MDMA can influence endocrine signaling at doses relevant for studies in humans [23]. Fluoxetine attenuated prolactin release after administration of racemic MDMA, and fluoxetine and a 5HT<sub>2A</sub> antagonist attenuated prolactin release after R(-)-MDMA, indicating that prolactin release is associated with serotonin release and indirect action on 5HT<sub>2A</sub> receptors by R(-)-MDMA [99].

Serotonergic deficits are associated with disruption of sleep patterns and architecture. In drug-naïve rats, a single dose of 15 mg/kg MDMA i.p. contributed to marked increases in motor activity, deep slow wave sleep, and wakefulness for 5 to 6 hours. Circadian patterns of motor activity and sleep/vigilance parameters were altered for up to 5 days post-treatment, after which most parameters returned to normal. In a single exposure to MDMA 3 weeks prior to the same procedure, rats experienced the same acute effects, but with shorter duration, suggesting that MDMA has the ability to influence sleep architecture and patterns acutely after this dose in drug-naïve rats, but these effects are mediated by experience with MDMA and do not persist beyond 1 week [221].

#### 4.2.2.7 Neuropsychological Effects in Animals

In rodents, doses of MDMA equivalent to human doses produce few behavioral effects. However, several dose-dependent differences on behavioral tests in rats have been reported, including increased locomotor activity and anxiety-related behaviors thought to be associated with serotonin syndrome [161, 222], and decreased social anxiety at 5 mg/kg i.p. [161]. Rats given 7.5 mg/kg MDMA, equivalent to four times the dose tested in humans, exhibited increased anxiety in the elevated plus maze [223], while rats given 15 mg/kg MDMA, equivalent to eight times the

dose tested in humans, exhibited reduced anxiety on the maze. A study of the sub-acute effects of four different doses of MDMA given daily for 1 week, found reduced anxiety with 1.25 and 2.5 mg/kg and increased anxiety with 5 and 10 mg/kg [214]. Lower doses used in these studies are comparable to dose used in human research and nonmedical settings. However, sample sizes used in the study were small. Rats given higher doses also reduced aggressive behavior as well as social investigation. MDMA produces some repetitive behavior in rodents, but not to the same degree as psychostimulants. Rats on MDMA walk around a cage perimeter, interpreted as an indicator of thigmotaxis, which is a sign of anxiety [124]. However, it is notable that a 2007 publication failed to find thigmotaxis in rats given 5 mg/kg MDMA [224]. In contrast, rhesus monkeys do not exhibit increased locomotor activity after receiving up to 2.4 mg/kg MDMA [197]. Some researchers have proposed that behavioral tests of anxiety may instead be measuring risk-taking behavior, or impulsivity [225]. It is also notable that the majority of rat studies with deleterious behavioral findings were conducted specifically in inbred male Wistar rats, suggesting that individual and gender-based differences could influence these findings [226, 227]. Preclinical data in animals suggests that the profile of neurotransmitter release observed after MDMA administration may increase the risk of mania in some individuals [228], although mania has not been reported as a side effect of MDMA or Ecstasy in humans. Conflicting findings on anxiogenic and anxiolytic dose-dependent effects of MDMA are likely to have limited applicability to humans, with transient anxiety being a possible side effect.

To date, no empirical investigations have been conducted on the effects of MDMA on primate social interactions, which limits the generalizability of rodent studies to the more complex and relevant social behavior of primates and humans. Morley and colleagues observed rat behavior after receiving 5 mg/kg MDMA, noting that this dose correlated with prosocial behavior, such as lying next to each other [161]. Subsequent studies suggest that MDMA increases prosocial behavior in rats by elevating oxytocin in the paraventricular nucleus through 5HT<sub>1A</sub> receptor agonism, with the oxytocin increase arising from the indirect effects of MDMA on 5HT<sub>1A</sub> receptors via serotonin release [160, 229, 230]. There have been no human pharmacological challenge studies combining MDMA with 5HT<sub>1A</sub> agonists, while 5HT<sub>1A</sub> antagonists have negligible effects on subjective or physiological effects of MDMA in humans [92, 231-233]. As a result, it is unclear whether the rat behavior is analogous to human reports of increased feelings of empathy or interpersonal closeness while under the influence of MDMA [2, 13, 234, 235].

MDMA given before training persistently enhances fear extinction learning in mice through a BDNF-dependent mechanism [178], which could be a possible mechanism of action for MDMA in combination with psychotherapy as a treatment for anxiety disorders. The dose of 5.6 mg/kg was approximately two times a human equivalent dose based on plasma values, but the findings are the first biological evidence of a lasting effect of MDMA on disruption of anxiety-related behavior in mice.

#### **4.3 Physiological Effects in Epidemiological Settings**

The vast majority of non-clinical epidemiological studies are retrospective comparisons of people who have previously self-administered Ecstasy, a study design that is unable to eliminate the possibility that one or more predisposing factors may lead to repeated Ecstasy use and the variables compared [5, 89, 236]. Samples are often selected on the basis of moderate to heavy self-reported Ecstasy use, with very few studies conducted in samples reporting the levels of moderate exposure seen in clinical trials. Many investigations have compared people reporting use of Ecstasy with non-Ecstasy using controls, mostly as a means of detecting long-term effects of Ecstasy use. Many of the studies do not appropriately match samples for substance use behavior, there is often concurrent use other illicit substances and the Ecstasy used is of unknown purity, dosage, and composition.



The acute effects of MDMA have an initial onset of approximately 30 minutes after oral intake and are characterized by anxiety, tachycardia, and elevated blood pressures [237]. Typical effects include diaphoresis, bruxism, jaw clenching, paresthesias, dry mouth, increased psychomotor activity, and blurred vision. Within an hour, these sympathomimetic effects are replaced by feelings of relaxation, euphoria, increased empathy, and communication. Taking a smaller supplemental dose may prolong these effects and this is being tested in the context of clinical trials. However, when too much additional MDMA is consumed in an uncontrolled setting, individuals report unpleasant symptoms of autonomic hyperarousal associated with feelings of restlessness, paranoia, and anxiety. With increased dosage sympathomimetic effects predominate, placing the patient at risk for cardiovascular instability, arrhythmias, and hyperthermia (see [Section 4.4 Toxicology in Animals and Epidemiological Settings](#)).

Retrospective surveys of Ecstasy use offer similar accounts of subjective effects to those reported in controlled studies of MDMA. Study respondents report experiencing stimulant-like effects, such as greater energy or talkativeness, and hallucinogen-like effects, including perceptual changes, visual distortions, or poor concentration, as well as feelings of closeness, compassion, or empathy toward the self or others [2, 234, 235, 238, 239]. The disparity in detection of entactogenic effects in retrospective versus controlled studies is largely due to failure to measure these effects, but might also relate to aspects of setting in controlled studies that do not permit enough unstructured interpersonal contact to produce or facilitate feelings of interpersonal closeness. Starting in the 2010s, more researchers are seeking to assess the prosocial effects of MDMA [35, 37, 38, 240].

The findings discussed in this section are of effects in low to moderate users of Ecstasy. Serious and life threatening events and effects in heavy users are discussed in [Section 4.4 Toxicology in Animals and in Epidemiological Settings](#). Because of these many confounds and issues, findings discussed from retrospective comparisons and case reports of Ecstasy using samples and controls are considered cautiously with respect to their degree of relevance for safety in clinical trials.

### 4.3.1 Immunological Effects

As supported by mild immunosuppressant effects found in rodents, a longitudinal study of regular Ecstasy and cannabis users found a sustained reduction in IL-2, increased levels of Transforming Growth Factor-Beta (TGF- $\beta$ ), and reduced CD4 cells, and regular Ecstasy and cannabis users reported experiencing a greater number of mild infections than occasional Ecstasy and cannabis users on a structured questionnaire [241]. Immunological effects of MDMA in humans are likely to involve serotonergic pathways and are discussed in more detail in [Section 5.3.2 Immunological Effects](#).

### 4.3.2 Thermoregulatory Effects

Thermoregulatory effects of Ecstasy taken in epidemiological settings are highly dependent on dose [242] and permissive factors, including high ambient temperature [243, 244], crowded conditions involving overwhelming social interaction, physical exertion, reduced fluid intake [243], and thyroid dysregulation [245, 246]. In the absence of these permissive factors from use in epidemiological settings, hyperthermia is rarely reported. For a detailed discussion on thermoregulatory effects when Ecstasy is combined with permissive factors, see [Section 4.4.6 Hyperthermia](#).

### 4.3.3 Cardiovascular Effects

Studies in Ecstasy users indicate that only people reporting average lifetime exposure of 900 tablets had cardiac abnormalities [247]. No abnormalities were found in people reporting lifetime exposure of approximately 200 tablets in the same study. Previous to this, echocardiographic readings in eight Ecstasy users also failed to find any cardiac abnormalities [45]. Valvular heart disease (VHD) only occurred after extremely heavy Ecstasy use, it is unlikely to be a risk within the research or therapeutic context where subjects are screened for relevant pre-existing conditions. For more information on toxicological effects, see [Section 4.4.7 Cardiovascular Toxicity](#).

### 4.3.4 Osmoregulatory Effects

Ecstasy use has been associated in the literature with acute symptomatic hyponatremia with the syndrome of inappropriate antidiuretic-hormone secretion (SIADH) involving raised antidiuretic-hormone, also known as AVP [248]. SIADH refers to disorders related to water and sodium balance characterized by the impairment of urinary dilution and hypotonic hyponatremia in the absence of renal disease or other identifiable non-osmotic stimuli known to activate the release of AVP [249]. MDMA is known to cause central release of both oxytocin and vasopressin through indirect effects of serotonergic signaling as previously described, and this activity indicates that it is not accurate to attribute the osmoregulatory effects of Ecstasy to SIADH, but rather this should be characterized as a pharmacological effect on AVP secretion.

AVP plays a key role in osmoregulation, and is released upon a change in plasma osmolality [250]. AVP is also involved in the response and adaptation to stress, through its effects on the HPA axis [250]. The rise in AVP does not seem to be part of a generalized stress response, but results from a pharmacological effect compounded by excessive fluid ingestion [251]. In Ecstasy users with confirmed urinary MDMA, a significant association was found between plasma osmolality, plasma sodium, and CYP2D6 extensive metabolizer/ intermediate metabolizer genotypes and COMT low-activity genotypes [252]. Effects of Ecstasy, combined with increased consumption of water and permissive factors, such as strenuous exercise in warm ambient temperatures, can be further exacerbated in the context of poor metabolism. Gauging appropriate water intake may be difficult for users to estimate because MDMA reduces perception of thirst and impairs judgment [253]. For more information on the risk of hyponatremia, see [Section 4.4.8 Hyponatremia](#).

### 4.3.5 Neurobiological Effects

Spurred on by animal studies that found repeated or high doses of MDMA damaged the axons of serotonin neurons, researchers began studying the effects of repeated non-medical or recreational use of Ecstasy in humans [83-85, 254]. Early investigations had a number of methodological flaws, including retrospective design and poor matching of Ecstasy users with appropriate controls [89, 255]. Later studies sought to remedy some of these problems by using carefully matched polydrug user or cannabis user controls, or by relying on a sample with relatively low exposure to psychoactive substances, including alcohol [256-259]. Some of these investigators also conducted longitudinal studies, comparing Ecstasy users, sometimes alongside controls, at two separate time points [260-262].

Researchers using slightly different methods have reported differing results. These include finding no differences between Ecstasy user and polydrug user controls in SERT binding sites [263], modest reductions in estimated SERT sites in Ecstasy users versus non-drug using or cannabis-using controls [264], and an association between decreased SERT sites and lifetime

Ecstasy use [265]. This study also reported finding slightly fewer 5HT<sub>2A</sub> receptor binding sites in both “Ecstasy preferring” and “hallucinogen preferring” groups. Studies in low to moderate Ecstasy users did not report an increase in this marker [266], and only one of three studies in heavy users detected a change in 5HT<sub>2A</sub> receptor density. [267-269]. A prospective study in moderate Ecstasy users also failed to find any chemical markers of neuronal injury, and only found decreased cerebral blood volume in the dorsolateral frontal cortex [266, 270]. A re-examination of brain imaging using the less specific SERT marker Beta-CIT indicate an inverse relationship between age of first use of Ecstasy and mid-number of midbrain serotonin sites without detecting any relationship between age of first use and frontal SERT sites [271]. A retrospective imaging study using a radioligand that maps serotonin synthesis found lower ligand presence (“trapping”) in prefrontal, orbitofrontal and parietal areas and higher presence in brainstem, frontal and temporal areas in Ecstasy users versus polydrug user controls, with a greater difference seen in men [272]. The researchers reported relationships between differences in trapping and cumulative use, duration and temporal proximity of use. The samples were not well-matched for drug use.

Studies comparing brain activity in Ecstasy users and non-Ecstasy using controls reported some but not many differences in brain activity. These included greater brain activation in the occipital cortex, with concomitant methamphetamine use contributing to increased activation to a visual stimulus [273]. The same group of researchers detected less within-region coherence in the thalamus in Ecstasy users who averaged 29 episodes of use when compared with non-Ecstasy-using controls [274]. In a retrospective study, Ecstasy users exhibited lower brain activity in bilateral dorsolateral prefrontal cortex compared with controls reporting no illicit drug use, with neither group exhibiting impaired task performance [275]. Ecstasy users exhibited a single difference in brain activity compared to polydrug using controls. A prospective study comparing brain activity before and after use of Ecstasy failed to detect differences in working memory, attention or brain activity [276], suggesting a relationship between repeated, regular use of Ecstasy and other drugs and changes in brain activation. Investigations of the interaction between genotype and regular Ecstasy use have supported differential effects upon reward-based attention or visual or verbal memory [277-279], with some findings supporting differences due to genotype and some failing to do so. A systematic examination of imaging studies comparing ecstasy users reporting consumption of 100 or fewer tablets with controls reported finding no evidence for an association between moderate Ecstasy and signs of structural or functional changes in the brain [79]. Given the small samples and uneven numbers with different genotypes, any conclusions await further support.

Sleep disturbances are thought to be associated with deficiencies in serotonergic signaling [280]. Examining sleep architecture in Ecstasy users, investigators found less total sleep time and less stage 3 and 4 sleep on the adaptation night, but no overall differences in sleep architecture [281]. Another study comparing heavy Ecstasy users with non-drug using controls found no differences in baseline sleep using electroencephalography (EEG) [282]. Early studies in mostly heavy Ecstasy users reported significant decreases in total sleep as well as stage 2 sleep [283], while studies conducted in the 2000s found Ecstasy users were able to fall asleep more easily upon depletion of catecholamine neurotransmitters suggesting an underlying difference in serotonergic control of sleep architecture [284, 285]. Findings of sleep disruption in Ecstasy users are not likely to be applicable to the exposures seen in research or therapeutic settings.

A study of breathing during sleep in 71 Ecstasy users and 62 polydrug users did not find overall differences in disrupted breathing, assessed via nasal cannula, but found that all moderate and severe breathing disruptions occurred in the Ecstasy using sample [286]. McCann and colleagues reported a relationship between cumulative (lifetime) Ecstasy exposures and instances of disrupted breathing during non-REM sleep and suggested Ecstasy users could be vulnerable to



potentially fatal sleep apnea. In contrast, other researchers failed to find greater night-time awakenings indicative of sleep apnea in Ecstasy users [281, 282], and the high rate of disrupted breathing McCann and colleagues detected even in the controls suggest that this measure may not provide clinically significant assessments. Taken together, it appears that MDMA acutely produces lighter sleep with fewer REM periods.

#### 4.3.6 Neuropsychological Effects

Previous reports have found an association between Ecstasy use and symptoms of depression or anxiety [287, 288]. A meta-analysis of self-reported depressive symptoms detected an association between Ecstasy use and scores on the Beck Depression Inventory (BDI), a popular self-report measure of depression symptoms [289]. However, the association was strongest in studies with small samples, and drug use variables were often incompletely reported and not verified through any methods save self-report in the studies analyzed. Many studies found that increases in self-reported anxiety or depression were more strongly related to polydrug use rather than to use of any one substance [290-293]. Two studies found an equal or stronger association between regular use of cannabis, and not Ecstasy, with anxiety, depression or other psychological problems [294, 295]. Anxiety regarding loss of control under the influence of Ecstasy could develop to a degree where it could lead to panic attacks. Case reports have been published describing panic attacks in individuals under the acute influence of Ecstasy [296]. Enduring panic attacks have been reported in individuals after repeated Ecstasy use [297, 298] and in one case, even after a single dose [299].

Neuroendocrine response to oral citalopram did not differ between Ecstasy users, cannabis users and controls [300]. People reporting regular drug use and Ecstasy use had higher levels of salivary cortisol in the evening, and higher salivary cortisol on the day of a multitasking activity [301], and higher salivary cortisol on waking that was unrelated to prefrontal SERT binding or self-reported depression symptoms [302]. A 4-year longitudinal study reported that factors other than Ecstasy use, including female gender and presence of financial and relationship difficulties, were more closely related to self-reported symptoms of depression [303]. Comparison of self-reported psychological symptoms in samples of people grouped by self-reported drug use found current Ecstasy users had lower global symptom severity scores than polydrug users [304]. In conclusion, it appears that the relationship between Ecstasy use on self-reported mood or psychiatric problems is not strong, with equal or stronger involvement of other factors.

In a prospective study comparing cognitive function in people before and up to 18 months after reported initiation of Ecstasy use, Schilt and colleagues found an association between Ecstasy use and performance on measures of verbal memory, but not attention or working memory [305]. All scores were within normal range; people who did not use Ecstasy showed greater improvement in performance at the second time of assessment than people reporting some use. A second prospective study examined working memory in people reporting Ecstasy use similar to subjects in Schilt's study with controls, and failed to find any significant differences in working memory and selective attention [276]. An analysis of findings from largely retrospective studies of Ecstasy users reported a small deficit in verbal or working memory [53]. Retrospective studies of polydrug users who use Ecstasy and controls reported impaired global motion processing without changes to local processing [306].

Not all studies report that Ecstasy users fare worse on measures of cognitive function than controls. A number of reports detected little or no significant differences between Ecstasy users and polydrug user controls in performance on tasks of cognitive function [236, 275, 276, 307-311], though other studies continue to find consistent differences, particularly in verbal memory [285, 312-315]. Regular use of many substances, including alcohol, may affect cognitive

function, with Ecstasy being only one of those substances [316]. Several reports have found relationships between cognitive function and use of other drugs as well as or instead of Ecstasy [278, 307, 309, 312, 317, 318].

The only study attempting to address effects of Ecstasy use on cognitive function in middle aged versus younger users did not find a greater degree of impairment. Schilt and colleagues reported impaired verbal memory in people who began using Ecstasy in their 30s compared with age-matched drug-naïve and polydrug using controls reporting some lifetime Ecstasy use, but did not find a greater effect size for Ecstasy use in this sample than in samples of younger Ecstasy users, leading them to conclude that Ecstasy use does not have a greater impact on cognitive function in older users [319].

The relationship between Ecstasy use and impulsivity has also been extensively examined, with some researchers reporting greater impulsivity in Ecstasy users and others failing to find any differences [84, 320]. Recent studies using both behavioral and self-report measures of impulsivity reached contradictory conclusions [311, 321, 322]. Two recent studies using the same measure of behavioral impulsivity in samples of heavy Ecstasy users obtained different findings [311, 321]. It is notable that Quednow and colleagues compared Ecstasy users with abstinent cannabis users and drug-naïve controls while Roiser and colleagues compared Ecstasy users with former Ecstasy users, polydrug users and drug-naïve controls, raising the possibility that results might have differed in part due to control group selection. It is possible that people who self-administer Ecstasy may already possess above-average levels of sensation-seeking and impulsiveness. To date, all such studies have used retrospective study designs and cannot rule out this possibility, and studies published in the last 2 years suggest that polydrug use may be equally or more strongly related to impulsivity in Ecstasy users [323-325]. The relationship between drug use, including Ecstasy use, and impulsivity, is complex.

#### **4.4 Toxicology in Animals and Epidemiological Settings**

In the sections below, nonclinical toxicological findings are presented for animals and epidemiological studies or case reports of morbidity and mortality in Ecstasy users. Data from epidemiological studies are provided, subject to the limitations in interpretation that result from unknown purity, dose, and quantity of MDMA existing in Ecstasy use in naturalistic settings.

##### **4.4.1 Single Dose Studies in Animals**

Single doses between 5 and 60 mg/kg have been administered in rodents. Since rodents are similar to primates in mg/kg dosing, the doses of 5 mg/kg and above, administered by any route of administration in rodents, are inappropriately high for comparison to human studies utilizing doses less than or equal to 125 mg, so findings are only useful as models of toxicology or abusive use in humans. A study of the long-term effects of a single dose of 5.7 mg/kg MDMA on estimated SERT sites in the brains of squirrel monkeys reported reduced sites in some frontal, temporal and parietal areas [326]. The plasma  $C_{max}$  of 725 µg/L in squirrel monkeys was three times greater than what is observed in humans after a single dose of 100 mg MDMA ( $C_{max}$  of 202.92 to 222.5 µg/L) [113, 327, 328], even after administration of a supplemental dose twice that of the initial dose 2 hours later, which increased  $C_{max}$  to 311.16 µg/L [328]. A handful of studies in rats have examined the effects of single toxic doses in comparison to low doses and determined that single doses have transient effects on serotonin depletion [78, 114, 108], likely due to reversible inhibition of tryptophan hydroxylase [17, 18, 20], which prevents additional serotonin from being produced and released.

#### 4.4.2 Repeated Dose Studies in Animals

The majority of toxicological studies employed multiple dosing regimens to account for the shorter drug half-life in animals compared to humans, with doses ranging from 5 mg/kg to 20 mg/kg, via s.c., i.p., oral, or gavage administration. Frequently, doses are administered at regular intervals of two to four times per day. Other regimens employ these doses once daily for 5 or 7 days. Nearly all preclinical toxicology data is derived from repeated dose studies. Preclinical research selected doses through use of simple dose conversions or allometric scaling, a method of modeling human equivalent doses in other species [329]. Comparison of pharmacokinetic data ( $C_{max}$ , AUC,  $T_{max}$ ) for plasma MDMA concentrations between humans and rodents, in light of the impact of route of administration, it is difficult to translate the relevance of high dose multi-day dosing findings in preclinical toxicity studies to intermittent dosing regimens in humans.

In order to establish the DMF and IND for MDMA, the sponsor supported randomized 28-day general toxicity studies in both genders of the rat (0, 10, 50, 100 mg/kg oral) and the dog (0, 3, 9, 15 mg/kg oral)[330]. Both sexes of dogs on 9 and 15 mg/kg MDMA and rats on 50 and 100 mg/kg MDMA gained less weight than those on control and 3 mg/kg, with significant differences in food consumption observed as early as the first week which were no longer significantly different by the fourth week. Gross observations at necropsy in the dog possibly related to MDMA included reduced testicular size on 9 and 15 mg/kg in the dog and prostatic enlargement in two dogs on 15 mg/kg. No gross lesions were seen in the rats at necropsy. Blood chemistry and urinalysis values were unremarkable in the dog. Clinical pathology findings showing a trend to decrease with dose in the rat were urinary pH, blood urea nitrogen, glucose, creatinine (females), lactate dehydrogenase (females), and chloride, in contrast total white blood cell count (WBC) and phosphorus showed a trend to increase with dose. No MDMA-related lesions were seen in the brains of either species.

#### 4.4.3 Genotoxicity

An Ames test of Ecstasy tablets with 0 to 57.5% MDMA, quantified by GC-MS, found no evidence of genotoxicity [331]. Micronuclear and chromosomal aberrance tests were performed in Chinese hamster ovary cells with MDMA purified from seized Ecstasy tablets and with N-nitroso-MDMA (N-MDMA), a putative metabolite of MDMA [332]. MDMA did not produce increases in either *in vitro* genotoxicity test.

#### 4.4.4 Carcinogenicity

There are no preclinical findings directly addressing the carcinogenicity of MDMA. No tumors were reported after 28 days of daily MDMA administration in rats (0, 10, 50, 100 mg/kg) and dogs (0, 3, 9, 15 mg/kg) in a sponsor-supported preclinical study [330]. In the absence of positive results in genotoxicity tests, carcinogenic potential from intermittent dosing of limited number of exposures to MDMA in controlled settings is not of concern.

#### 4.4.5 Reproductive and Developmental Toxicity

MDMA (15 mg/kg, s.c.) administered to pregnant rats was detected in amniotic fluid [333] indicating the potential for neonatal exposure. Preliminary teratological studies in rats (N=12 per dose) given 0, 2.5, or 10 mg/kg MDMA by gavage on alternate gestational days (GD) 6 to 18 found no abnormalities in gestational duration, litter size, neonatal birth weights, or birth defects (N=10 litters per dose), despite statistically significant reduction in maternal weight gain at 10 mg/kg [334]. These results are in contrast to physiological abnormalities resulting from prenatal methamphetamine and d-amphetamine exposure in mice and rabbits [335].

In a single-generation fertility and developmental toxicity study, C57BL/6 mice (N=25 per dose per gender) received a daily dose of 0, 1.25, 5, or 20 mg/kg MDMA via gavage [336]. Dosing for females spanned 2 weeks before mating through GD15 of pregnancy. Dosing for males spanned 4 weeks through the first day of pregnancy. There were no cases of MDMA-related mortality in females at all treatment levels. Gross necropsy of organs of MDMA-treated groups of male and female mice were unremarkable. No changes in copulation or fertility indices arose in MDMA-treated animals, but fewer pregnancies arose in all three MDMA-treated groups. When the fetuses were examined, no external, visceral, or skeletal malformations were detected in control or 1.25 mg/kg groups, but at 5 mg/kg (2 of 129) and 20 mg/kg (5 of 138) fetuses exhibited a cleft palate, anophthalmia, or skeletal malformations (short tail). Taken together, these studies suggest that MDMA has weak reproductive or developmental toxicity at high doses when MDMA exposure starts 2 weeks prior to mating and continues through GD15, which temporally covers ovulation through organogenesis and closure of the hard palate, in the females and spermatogenesis in the males.

In a separate perinatal/postnatal toxicity study done by the same researchers, C57BL/6 female mice (N=25) received a daily dose of 0, 1.25, 5, or 20 mg/kg MDMA via gavage daily from GD6 slightly after implantation through postnatal day (PND) 21 end of lactation [336]. Pup viability was assessed daily and gross external examination of pups occurred on PND 0, 4, 7, 14, 21, and 28. Behavioral and physical indices of development were observed in the F1 animals, such as pinnae detachment and righting reflex. Testes descent in males occurred on PND20 and vaginal opening occurred in selected females on PND30. Delivery and post-partum (nesting) behavior did not differ across treatment groups, and no MDMA-related differences in pup viability were detected, including pup survival rate and sex ratios per litter. No significant abnormalities were observed at necropsy of mice either found dead at lactation nor killed at PND20. In contrast to the first study described above where MDMA was given 2 weeks before mating through GD15, when MDMA was given to only the females from GD6 to the end of lactation (both studies covered the period of organogenesis and closure of hard palate), there were no signs of impaired development and no significant differences in sexual development or reproductive capacity of F1 and F2 mice. This suggests that either dual exposure of male and female breeding pairs exacerbated reproductive toxicity, or possible evidence of a critical period for MDMA reproductive toxicity prior to organogenesis.

Male fertility after prenatal exposure was studied in male pups born to female Sprague-Dawley rats (N=6 per group) that received 0, 0.5, 5, or 10 mg/kg s.c. daily for three consecutive days per week for 10 weeks, including gestation and 3 weeks of lactation [337]. These females were mated with untreated males. The 5 mg/kg s.c. dose is two-fold greater than a human-equivalent dose based on plasma levels in other studies [78, 114, 119] and s.c dosing leads to higher plasma levels than dosing by gavage which was used in the studies above. There were no signs of toxicity in the 0.5 and 5 mg/kg groups, but dams in the 10 mg/kg group showed signs of sickness the week before delivery, and four of the six receiving 10 mg/kg and one of the five receiving 5 mg/kg were found dead at or prior to GD16. Mortality at 10 mg/kg s.c. indicates that this dose is too high for use in reproductive toxicity studies; the authors subsequently discontinued the 10 mg/kg dose after week 10. Vestibular and motor function were assessed on PND21, with no differences between groups. Balano-preputial separation happened later than controls after 5 mg/kg in male pups on PND37-54. There were no differences in mating or fertility rate in F1 males. Hormone levels were similar across groups at PD81 and sperm morphology was unaffected. However, MDMA administration resulted in a significant higher incidence of DNA damage in Comet Test of sperm DNA at 5 mg/kg in relation to the control group. Minor dose-dependent alterations were seen in testicles, spleen and kidneys. There were no pathologies of the epididymis. Testicles showed a slight decrease in numbers of germ cells in 5 mg/kg treated rats.

A second study investigated male fertility after 0.5, 5 and 10 mg/kg administered s.c. once daily three times per week in rats (N=20 per group) for 12 weeks, covering puberty to onset of sexual maturity [338]. Ten rats per dose were mated with untreated females, with mating behavior alone serving as measure of reproductive function without reporting signs of conception. The other 10 rats per group were examined for testicular and sperm parameters, including sperm count and motility and morphology. There was a dose-dependent increase in tubular degeneration in testes in MDMA-treated rats, but sperm motility and morphology was unaffected. In a sponsor-supported preclinical study, microscopic evidence of possible testicular atrophy and prostatic enlargement was also found in one of three dogs after 28 days of 9 mg/kg oral MDMA and in two of three dogs after 15 mg/kg oral MDMA [330]. Taken together, these studies suggest minimal male fertility toxicity at human-equivalent doses, with signs of increased toxicity at higher doses.

In an initial developmental toxicity study, pregnant rats were administered twice-daily injections of high doses of MDMA (15 mg/kg) or saline from embryonic days (E) 14 to 20. Rat pups that had received MDMA showed reductions in the dopamine metabolite homovanillic acid, along with reductions in the serotonin metabolite 5-HIAA. Prenatally exposed MDMA animals also had reduced dopamine and serotonin turnover in the nucleus accumbens [339]. The same team reported postnatal exposure to MDMA correlated with reductions in serotonin and its metabolite, as well as significant increases in dopamine turnover and the prevalence of a dopamine metabolite in multiple forebrain structures and the brainstem. BDNF was significantly increased (19% to 38%) in all forebrain structures and in the brainstem in MDMA-exposed neonates [340]. The researchers proposed that the increase in BDNF was compensating to minimize MDMA effects. However, later studies found that neonatal MDMA exposure did not affect hippocampal concentrations of serotonin or dopamine [341] and that enhanced BDNF detected in the occipital lobe did not mediate the abnormal serotonergic signaling observed following neonatal MDMA exposure [342]. PND 11 and 20 were proposed to be equivalent to the third trimester of gestation in humans [340], so it is possible that exposure to high doses of MDMA *in utero* could have developmental effects, but these do not appear to be related to BDNF levels. The doses used in the rat studies are approximately eight to 10 times greater than a human equivalent dose.

Prenatal MDMA exposure at high doses significantly increased locomotor activity of rat pups in a 20-minute novel cage environment test [339]. Rodents treated with MDMA during development were not significantly different than rodents who received MDMA as adults. The results of several behavioral tests indicate that developmental MDMA exposure combined with adult exposure may interfere with some aspects of learning, including visual-spatial memory and time spent with a novel object [341]. Neonatal MDMA administration did not alter working memory in the object-recognition test in young adulthood (PD 68 to 73) and there were no differences in binding of the radiolabeled selective serotonin reuptake inhibitors (SSRI) citalopram to the SERT at this age. However, the pretreated animals showed increased thermal dysregulation and serotonin syndrome responses following MDMA challenge, especially with respect to head-weaving stereotypy [343]. Another team also found that neonatal rat MDMA exposure exacerbated hyperthermic response to a subsequent dose of MDMA [344]. A study in neonatal rats suggests two distinct critical periods wherein repeated doses affected learning versus acoustic startle [345]. Serotonergic factors may be involved in the developmental effects of MDMA, with the SSRI citalopram producing similar learning impairments in neonatally exposed rats [346]. Given differences between human and rodent development and thermoregulation, it is not clear whether such findings can be generalized to humans (see [Section 4.2.2.3 Thermoregulatory Effects in Animals](#)).

Previous research supported a possible link between Ecstasy use and birth defects [347], while an epidemiological study of a large cohort of pregnant women in England conducted in 2004 failed



to support this link, at least in respect to a specific cardiac defect [348]. However, the authors also stated that exposure to MDMA in their sample was too low to establish risk. An earlier survey of a drug-using population suggests that most women cease using Ecstasy when they learn they are pregnant [349]. A 2012 survey of 96 women in the UK interviewed about their drug use during pregnancy found a link between self-reported extent of prenatal MDMA exposure and delays in infant development at 12 months, with heavily exposed infants delayed in mental and motor development, but not language or emotional development [350]. These results were repeated in a 2016 survey of 96 mothers who reported heavier MDMA use ( $1.3 \pm 1.4$  tablets per week) during pregnancy. Infants had motor delays from 4 months to 2 years of age that were not attributable to other drug or lifestyle factors [351]. Since there may be a critical period during which exposure to MDMA could alter development, and as a result of the relative lack of information concerning its developmental toxicity, women who are pregnant or who are not using an effective means of birth control should not receive MDMA in clinical trials. None of the sponsor's studies enroll pregnant or lactating subjects.

#### 4.4.6 Hyperthermia

At least one case series of individuals seen on the same night and near or in the same nightclub suggest a relationship between Ecstasy dose and likelihood of hyperthermia [352]. A case report and some findings in rodents suggest that hyperthyroidism or thyroid dysregulation may play a role in MDMA-related hyperthermia in humans [245, 246]. When assessing acute effects of Ecstasy, hyperthermia is one of the more frequently reported acute harms of Ecstasy [53, 242].

A study of rats receiving subcutaneous injections of 9 mg/kg MDMA, just under half the LD50 of 20 mg/kg in rats housed together, reliably produced malignant hyperthermia in the context of warm ambient temperatures of 29°C and during social interaction [187]. At this dose, MDMA monotonically increased intracerebral heat production and muscle temperature while causing strong and sustained peripheral vasoconstriction, which inhibits heat dissipation. Social interaction on its own also induced metabolic brain activation and transient vasoconstriction in rats, which compounds the hyperthermic effects of MDMA observed at toxic doses and warm ambient temperatures. These effects are likely to be mediated through dopaminergic pathways [353, 354], which have been shown to play a minor role in producing the effects of MDMA in humans [34].

#### 4.4.7 Cardiovascular Toxicity

Injections of 20 mg/kg MDMA in conscious rats assessed by radiotelemetry (10.7 times the equivalent dose in humans), found that MDMA caused a prolonged increase in blood pressure [202]. In the same study, MDMA was found to produce mild isotonic contractions of aorta and vas deferens vascular tissue in anesthetized rats, but could also inhibit prejunctional contractions evoked by stimulation [202].

The elevation of blood pressure and increased heart rate produced by MDMA, similar to that produced by other sympathomimetic drugs, can lead to additional risks and complications [355-357], such as stroke, cardiac events, or other cerebrovascular events, including cerebral venous sinus thrombosis [358] and cerebral or subarachnoid hemorrhage [80, 359-363]. In two such cases, a previously existing underlying arteriovenous malformation appeared to play a role in the event [359, 361]. Intra-cardiac pressures, intra-arterial pressures, angiotensin II, pain, and adrenergic ( $\alpha_2$ ) central nervous stimuli can also influence AVP secretion [364]. Increased AVP concentration is described in several studies as a strong predictor of mortality in patients with chronic heart failure and acute heart failure, and contributes to increases in blood pressure [365]. As with any amphetamine, increased heart rate (tachycardia) and elevated blood pressure can also

lead to cardiac events, such as arrhythmias or myocardial infarction [366, 367]. Fatal dysrhythmias have been reported following heavy MDMA use, resulting in ventricular fibrillation and asystole. Individuals with underlying cardiac and/or pulmonary disease and preexisting conditions such as Wolff-Parkinson-White syndrome are especially at risk for heart failure and fatal arrhythmias. Although the presence of MDMA was rarely confirmed in reported cases, these types of events are all well-established complications of hypertension and can occur after use of amphetamines. There have been no such events to date in any clinical trial of MDMA.

Some researchers have expressed concern that MDMA activity at 5HT<sub>2B</sub> receptors might be indicative of increasing risk of VHD with repeated use [24]. Studies in Ecstasy users indicated that only people reporting average lifetime exposure of 900 tablets had cardiac abnormalities indicative of potential VHD [247], and a case of VHD has occurred in a man reporting approximately 16 years of heavy Ecstasy use, from age 17 to 33 years old. [368]. No abnormalities were found in people reporting lifetime exposure of approximately 200 tablets in the same study. Echocardiographic readings in eight Ecstasy users also failed to find any cardiac abnormalities [45]. Since VHD-associated changes and VHD only occurred after extremely heavy Ecstasy use, they are unlikely to be a risk within the research or therapeutic context.

#### 4.4.8 Hyponatremia

A number of case reports describe hyponatremia after uncontrolled, non-medical Ecstasy use [54, 369-371]. A recent meta-analysis showed that a moderate reduction of serum sodium concentration is associated with an increased risk of death in different pathologic conditions [372]. Relationships have been found between reduced plasma sodium, a measure of hyponatremia, and variations in COMT and CYP2D6 genotypes, possibly related to increased AVP and oxytocin release associated with MDMA [252]. Active doses of MDMA likely inhibit CYP2D6 in most individuals, as described in Section 5.2.1 Pharmacokinetics. Behavioral factors, including vigorous exercise and excessive consumption of water without an attempt to replace electrolytes, and an increase in the anti-diuretic hormones AVP and oxytocin, likely all contribute to this very rare but SAEs in Ecstasy users [32]. Women are generally more likely to exhibit hyponatremia than men [373, 374], including Ecstasy or MDMA related hyponatremia [54]. Heart failure is commonly associated with hyponatremia, and is also characterized by increased concentrations of AVP [375-377]. Hyponatremia has not occurred during a controlled clinical trial with MDMA.

#### 4.4.9 Hepatotoxicity

*In vitro* studies and studies employing high, repeated doses of MDMA, estimated as being at least five times greater than expected in a clinical trial [378], report damage to liver cells [379-381]. Though many of these studies employed MDMA concentrations much higher than would occur after human ingestion, there are reports of liver disease in Ecstasy users. Studies in rats suggest a role of body temperature in promoting liver toxicity. A review of the literature highlights a number of potential factors, including body temperature and metabolism in preclinical studies and polydrug use, including alcohol, and environmental factors in humans [382]. Due to disparities in dosing and method, it is hard to establish whether these findings are relevant for liver toxicity in human Ecstasy users.

Hepatotoxicity (liver disease or damage) was reported in approximately 16% of 199 case reports from Ecstasy users in non-medical, uncontrolled settings, collected from the mid-1990s to 2001, making it the third most common serious adverse report in the literature. There appears to be more than one pattern of Ecstasy-related hepatotoxicity, and a number of factors, including polydrug use and setting of use may be involved [382]. Acute liver failure or hepatitis has

occurred after reported ingestion of a single Ecstasy tablet [383-386]. In other cases, hepatotoxicity has occurred after months of regular Ecstasy use [387]. Standard toxicity studies failed to find liver damage after MDMA in rats or dogs after 28 days of exposure [330], nor have any cases of liver disease arisen during controlled studies. Examinations of case reports and a number of *in vitro* studies suggest an association between hyperthermia and hepatotoxicity. However, liver disease also occurred in some individuals without the occurrence of hyperthermia, appearing after continued use and resolving after abstinence. These reports suggest a potential immunological mechanism. Since hepatotoxicity has been noted in Ecstasy users, *in vitro* and *in vivo* studies have examined the hepatotoxicity of MDMA. These studies show that high repeated doses of MDMA can impair liver cell viability *in vivo* [379], and can increase profibrogenic activity in cultured stellate cells [381] while reducing cell viability without producing lipid peroxidation *in vitro* [379, 388]. At higher ambient temperatures, a toxic dosing regimen was capable of increasing lipid peroxidation and activating apoptosis due to oxidative stress [389]. A single intraperitoneal dose of 20 mg/kg in rats was still capable of disrupting glutathione homeostasis, decreasing antioxidant enzyme activity, and lipoperoxidation activating apoptosis in one study [390]. However, peak liver exposure to MDMA in sponsor-supported studies should be approximately one-eleventh the concentration shown to impair cell viability in these studies. No cases of liver disease or hepatotoxicity have occurred in controlled clinical trials with MDMA. See [Section 5.3.6 Hepatic Effects](#) for discussion of liver panel results in sponsor-supported clinical trials.

#### 4.4.10 Neurotoxicity

Repeated very high doses of MDMA in animals reduce total serotonin levels in the brain, impair transport of serotonin, and cause psychobehavioral changes such as increased anxiety [124, 226, 391-393]. In combination with other drugs or in high dose binge administration studies, MDMA may provoke serotonin syndrome. For example, rodents respond to high doses of MDMA by exhibiting flat body posture, forepaw treading and an erect tail (“Straub tail”) [393]. These behaviors are considered indicators of serotonin syndrome. Doses used in most preclinical studies of neurotoxicity are at least five times the amount used in clinical trials or nonmedical settings, and can be as high as 20 times that amount. Studies in rodents and primates suggest that repeated high doses of MDMA could reduce regional serotonin, damage serotonin axons and cause neurotoxicity [124, 135, 394-397] and promote apoptosis in the hippocampus after 5 or 10 mg/kg MDMA given daily for 1 week [214]. However, the majority of these studies employed large doses of MDMA that overestimated human-equivalent doses, with findings now clearly indicating that doses used in nearly all rat and most primate studies are inappropriately high for comparison to use in clinical settings and are more pertinent toxicological effects of MDMA [78, 114, 119].

Most studies suggested that heavy but not moderate Ecstasy users had impaired verbal memory and lower numbers of estimated SERT sites, assessed via imaging with radioactively labeled ligands in positron emission tomography (PET) or single photon emission tomography (SPECT), with heavy use often defined as 50 or more times or tablets. Taken together, findings from these studies suggest there is some risk of long-term effects in heavy Ecstasy users with respect to number of estimated SERT sites in specific brain areas and performance on measures of memory. However, interpreting findings of changes in serotonin receptors or cognitive function after repeated Ecstasy use are complicated by the possible impact of polydrug use and other potential pre-existing factors in retrospective reports, and the findings are not readily transferrable to use of MDMA in a therapeutic or research context.

Many investigations have examined cognitive function in Ecstasy users with the goal of demonstrating long-term effects of purported neurotoxicity of Ecstasy. Rogers and colleagues



performed a meta-analysis on a large number of retrospective studies of Ecstasy users and various cognitive functions. Given methodological flaws in this type of analysis, the investigators cautiously concluded that there might be a significant effect of Ecstasy use on verbal memory, and a lesser effect on visual memory [53]. Retrospective designs and inappropriately matched samples continue to appear in the literature [398-400], even when using multiple control groups. Two meta-analyses of memory in Ecstasy users arrived at somewhat contradictory conclusions [401, 402]. Both detected an association between Ecstasy use and impaired performance on at least some measures of memory. However, one reported that this association had a medium to large effect size with no effect of Ecstasy dose [401], while the other reported that the association had a small to medium effect size with an Ecstasy dose effect, and that polydrug use itself contributed to impaired cognitive function [402]. A meta-analysis comparing current Ecstasy users and drug-using controls on visuospatial skills reported that current users performed less well on measures of visual recall, recognition and item production than controls [403], but found no significant relationship between lifetime Ecstasy use and visuospatial task performance. A longitudinal study comparing people who continued to use Ecstasy with those who did not do so detected lower performance on immediate and delayed visual memory [404]. In a second follow-up in the same sample reported lower scores in visual memory, at marginal significance and no further impairment [405]. An examination of the relationship between elements of Ecstasy use history and verbal memory reported that use in the past year, especially in men, was associated with impaired verbal memory [406]. The authors suggest that gender differences in polydrug use may be involved. A study comparing performance on a test of verbal memory in 65 ecstasy users enrolled in clinical trials of MDMA and an equal number of age and gender matched non-drug using controls from other trials failed to detect significant differences between the two groups [407]. This study employed a pre-determined measure of clinical significance, 1.5 times the average standard deviation of the healthy controls, and used a Bayesian statistical test suited for assessing a null hypothesis. It is notable that none of the subjects were enrolled in studies designed to compare cognitive function in ecstasy users, which may have reduced anxiety and potential risk of “stereotype threat” that may be faced by substance users completing assessments of cognitive function, which was done to reduce expectancy in the study [408].

The nature and strength of the association between regular Ecstasy use and any impairments in executive function remains inconclusive, with studies reporting conflicting results [5, 258, 259, 409, 410]. Findings from a study published in 2014 did not find differences in multitasking [301]. A meta-analysis comparing executive function in Ecstasy users and non-Ecstasy using controls found a significant effect of Ecstasy use on one component of executive function (updating), no effect on another (shifting) and mixed results when looking at other components (response inhibition and access to long-term memory) [411]. Polydrug use likely contributes to findings of impaired executive function seen in Ecstasy users [292, 412]. Current research has not settled the question.

Psychiatric problems after uncontrolled, non-medical Ecstasy use were reported in 22.1% of 199 case reports from the early 1990s to 2001, and are the most common reason for appearance at an emergency department [52, 55]. Psychiatric symptoms included affective responses, such as dysphoria, anxiety, panic, and psychotic response, as well as cases with mixed psychotic and affective features. The most common problem reported included panic, restlessness and psychotic response, as seen a systematic review and several epidemiological case series [53, 413]. The mechanisms behind Ecstasy-associated psychiatric problems remain unclear, but are likely the result of an interaction between pharmacology and individual susceptibility. The difficulty of assessing the frequency of these events is increased given that pre-existing psychiatric problems occur in people who choose to use Ecstasy [414] and findings of an association between use of Ecstasy and other drugs and self-reported symptoms of anxiety and depression. As described earlier, most cases of psychological distress after Ecstasy use resolved after supportive care [52, 55].

Anxiety responses associated with MDMA administration reported in controlled trials have resolved over time, usually either during the period of acute drug effect or with the waning of drug effects.

#### **4.5 Serious Reports, Mortality, and Morbidity in Animals and Epidemiological Settings**

Intravenous MDMA doses that cause lethality in 50% of the cases, known as the LD50, are 97 mg/kg in mice, 49 mg/kg in rats, 14 to 18 mg/kg in dogs, and 22 mg/kg in monkeys [106]. LD50 may vary across strains, sexes, and housing conditions [415-417]. For example, LD50 in mice housed together is 20 mg/kg, which is considerably lower than in isolated animals [189, 418]. Reducing ambient temperature and administering the 5HT<sub>2A</sub> antagonist ketanserin reduced lethality, suggesting that amplified elevation in body temperature and activity at serotonin receptors may promote lethality in group-housed mice given MDMA [189]. Considerable variation across studies in environmental factors, that are often underspecified in published reports, contribute to challenges in extrapolating findings in animal studies that may be relevant in epidemiological settings.

A number of SAEs, including fatalities, have been reported in humans after Ecstasy use in unsupervised and uncontrolled settings. These events are relatively rare given the prevalence of Ecstasy use [49, 50]. These include hyperthermia (potentially arising from “serotonin syndrome”), psychiatric problems, hepatotoxicity (secondary to hyperthermia), cardiac disorders and hyponatremia [49, 52-54, 419]. Set and setting likely play a role in the development of some Ecstasy-related AEs, such as vigorous exercise, lack of attention to somatic cues, and too little or too much hydration combined with pharmacological action on AVP resulting in hyperthermia or hyponatremia [51, 371]. Even if ambient temperature does less to moderate the effects of MDMA on body temperature than originally believed based on animal studies, other environmental and behavioral factors, as those related to vigorous exercise, may be involved. It is important to note that not all reports of AEs in Ecstasy users provide information on whether MDMA was detected in plasma or other fluids, with some relying on self-report or the reports of friends as to identity of substances consumed. Reports indicating detectable MDMA will thus be the best indicators of an actual association. Unexpected drug-related SAEs have not occurred in any of the human MDMA research studies thus far.

While case reports do not provide an appropriate basis for estimating the relative frequency of these events, they can provide information on the possibility of an event occurring. Most Ecstasy-related emergency department admissions are the result of people experiencing anxiety or panic reactions after use and involve supportive care only [52, 55, 420]. An extensive systematic review reached similar conclusions concerning the frequency and nature of emergency department admissions, though also noting that owing to complexities of nonmedical and recreational use, the researchers found it hard to establish a lethal dose [53]. However, a pair of case series drawn from two different events suggests a general relationship between estimated dose and number of emergency department admissions after exhibiting seizures, unresponsiveness or hyperthermia, with both series reporting high doses of MDMA (230 and 270 mg) in sample tablets or capsules [421, 422]. As is the case with fatalities associated with reports of Ecstasy use, medical emergencies after Ecstasy use are more likely to occur in men [52]. Individuals consuming Ecstasy with pre-existing conditions are at increased risk when consuming drugs of unknown purity, identity, and dose in uncontrolled settings.

**Table 2: Summary of Published Morbidity and Mortality Reports**

Body System	Reports	Morbidity Reports	Mortality Reports	Total Reports
Thermoregulatory Disorders	Hyperthermia, Hyperprexia, Rhabdomyolysis, Hypoglycemia	135 [80, 421, 423-438]	43 [80, 217, 421, 423, 435, 439, 440]	178
Cardiac Disorders	Cardiac valve disease, Ventricular fibrillation, Cardiac arrest, Arrhythmia, Myocardial infarction, Generalized tonic-clonic seizure, Acute coronary syndrome, Myocardial necrosis, Cardio-respiratory arrest, Cardiomyopathy	15 [367, 368, 441-447]	12 [366, 423, 448-452]	27
Osmoregulatory Disorders	Cerebral oedema, SIADH, Urinary retention, Hyponatremia, Acute renal failure	18 [453-465]	6 [367, 466-470]	24
Hepatobiliary Disorders	Acute fulminant hepatitis, Liver disease, Disseminated intravascular coagulation	4 [386, 447, 471, 472]	5 [473-477]	9
Blood and Lymphatic System Disorders	Aplastic anemia	3 [478, 479]	1 [480]	4
Injuries, Poisonings, and Procedural Complications	Anaphylactic shock, Facial rash eruption	1 [481]	1 [482]	2
Nervous System Disorders	Hemorrhage, Infarct, Hippocampal sclerosis (suspected), Encephalopathy, Amnestic syndrome	13 [355, 356, 483-490]	0	13
Dental Disorders	Xerostoma, Bruxism, Dental erosion	15 [491-493]	0	15

<b>Body System</b>	<b>Reports</b>	<b>Morbidity Reports</b>	<b>Mortality Reports</b>	<b>Total Reports</b>
Psychiatric Disorders	Psychotic episode, Depressive episode, Obsessive-compulsive disorder, Autoenucleation	4 [494-496]	0	4
Respiratory, Thoracic, and Mediastinal Disorders	Subcutaneous Pneumomediastinum, Epidural pneumatosis, Diffuse alveolar hemorrhage, Asthma	9 [366, 497-504]	0	9
Ophthalmic Disorders	Lagophthalmos, Keratopathy, Bilateral sixth nerve palsy	4 [505, 506]	0	4
Injuries, Poisonings, and Procedural Complications	Unknown cause of death	0	204 [366, 507]	204

Four hundred ninety-three case reports, with 272 of these resulting in death, associated with Ecstasy use from 1986 through 2016 are summarized in Table 2. Of these 272, 32 were described in a cumulative 2002 literature review with incomplete citations of sources, and are reported in addition to individual case reports of morbidities in the literature [423]. Detectable levels of MDMA in blood or urine are reported in less than half of these case reports, and range from 50 ng/mL (reported as less than 0.05 mg/L) in the case of anaphylactic shock [482] to 1500 ng/mL (reported as 1.5 mg/L) in a fatal case of hyperthermia and rhabdomyolysis [440]. It is more difficult to associate events with MDMA when the compound is not detected or when detection is for amphetamines in general. Some events, such as VHD, acute hepatitis with gallbladder inflammation, liver disease, or urinary retention occurred in individuals who self-reported daily use for months to years prior to the event. In the majority of the 202 poisoning cases with unknown cause of death, Ecstasy was used in combination with opiates by drug addicts who died in the UK and Wales between 1996 and 2002 [507], and polysubstance use is common in the majority of serious reports presented.

Thermoregulatory disorders play a part in the development of a constellation of disorders across body systems described below. Primary symptoms are hyperthermia resulting rhabdomyolysis described in 135 reports of morbidity and 43 reports of mortality, constituting the most common acute adverse effect associated with Ecstasy. Sympathomimetic effects of MDMA, at unknown doses and purity, in combination with permissive factors in uncontrolled settings, can lead to serious reports of acute and persisting adverse effects on multiple organs. In research settings, the risk of hyperthermia is limited by controlling ambient temperature, conducting treatment sessions in relaxed, private environments, and generally limiting permissive factors.

Cardiac disorders associated with Ecstasy in the context of hyperthermia resulted in 15 reports of morbidity and 12 reports of mortality. Several fatal cases of cardiac arrest were reported. In addition, a non-fatal cardiac arrest occurred in the context of a genetic arrhythmia disorder, catecholaminergic polymorphic ventricular tachycardia [442]. Apparent use of Ecstasy, with concurrent use of other amphetamines during pregnancy, was associated with seizures and myocardial infarction [445, 446]. As evidenced by these reports, individuals consuming Ecstasy

with pre-existing conditions that can influence cardiovascular and cardiac function are at increased risk when consuming drugs of unknown purity, identity, and dose in uncontrolled settings.

Osmoregulatory disorders associated with Ecstasy in the context of hyperthermia resulted in 18 reports of morbidity and six reports of mortality, with acute renal failure (ARF) as the most common cause of death. As described in [Section 4.4.8 Hyponatremia](#), increased AVP secretion caused by MDMA in combination with permissive factors in uncontrolled settings can lead to serious reports of acute and persisting adverse effects on multiple organs, including the liver. Individuals consuming Ecstasy with pre-existing conditions that can influence renal function are at increased risk. In response to this risk, many users tend to overcompensate with excessive consumption of water, leading to dilutional hyponatremia. Prevention of hyponatremia with limited consumption of electrolyte containing fluids and controlled ambient temperatures are required to preserve the body's homeostatic maintenance of fluid balance.

Hepatobiliary disorders associated with Ecstasy use resulted in four reports of morbidity and four reports of mortality. One of the mortality reports happened 1 week after Ecstasy use and was consistent with acute fulminant hepatitis in the absence of viral infection. This patient died despite liver transplantation efforts [\[473\]](#). Typically, mortality results from disseminated intravascular coagulation (DIC) caused by platelet dysfunction associated with liver failure. Non-fatal morbidity reports range from acute hepatitis associated with daily usage of five to eight tablets of Ecstasy for 3 months in combination with alcohol [\[471\]](#) to liver damage in combination with congestive cardiomyopathy [\[447\]](#). Given that polysubstance use and prior insult to liver function cannot be ruled out, the frequency of isolated serious hepatotoxicity cases in the absence of hyperthermia are rare among serious reports associated with Ecstasy use. Hepatotoxicity is more common among serious reports in combination with hyperthermia and acute renal failure.

Blood and lymphatic system disorders associated with Ecstasy use resulted in three morbidity reports and one mortality report of aplastic anemia. The death after aplastic anemia occurred from complications of immunosuppressant therapy followed by an allogeneic stem cell transplant, 17 months after the first admission [\[480\]](#). The patient had initially presented with progressive weakness and epistaxis, resulting from daily Ecstasy use for 7 months, combined with heavy alcohol intake. Further examination revealed the replacement of bone marrow tissue with fatty deposits, likely due to alcohol consumption and exacerbated by chronic Ecstasy use. Three reports of morbidity ranged in prior Ecstasy use levels from once to four times in the prior year, with two cases spontaneously resolving within 2 months and the treated case failing immunosuppressive therapy and recovering 4 months after subsequent bone marrow transplant [\[480\]](#).

The report of possible anaphylactic shock and subsequent death occurred in a 13-year old girl who had at least one previous exposure to Ecstasy [\[482\]](#). Her friends reported that she experienced swelling lips after her first exposure. After approximately 1.5 tablets, the girl experienced nausea and vomited, and later had difficulty breathing. On admission she was hypothermic and hypotensive. A low level of MDMA (<0.5 mg/dL) was detected in blood. None of the other individuals consuming tablets from the same batch underwent similar experiences. Autopsy found a massive brain edema as well as laryngeal oedema and lung congestion. Chemical analyses ruled out hyponatremia. The reaction may have been to MDMA or to an adulterant in the tablet. The authors of the report do not report whether tablets were assessed for contents.

Memory difficulties arising immediately after Ecstasy use have been reported in a sporadic user [\[487\]](#). The memory difficulties arose in a man reporting use of Ecstasy five or six times, with

confusion and cognitive impairment reportedly occurring after taking a single tablet at a party. Cognitive function was assessed 7 years later. Imaging showed signs of hippocampal sclerosis. It is not clear from the report whether the individual used Ecstasy prior to or after this event. The individual had hypertension, raising questions concerning possibility of a cerebrovascular event. In a neurological serious report with 0.83 ng/mL MDMA detected in the hair of a girl who developed encephalopathy [486] during chronic low or moderate Ecstasy use, cognitive function and memory problems associated with neurological damage was reported. Upon cessation of use 16 months later, extensive hippocampal remodeling was reported assessed through PET scans. This finding is consistent with hippocampal dendritic spine remodeling observed in rats receiving 20 mg/kg MDMA for four days intended to simulate chronic usage in humans [508], however the clinical presentation was also similar to CNS herpes infection, so it is difficult to attribute this isolated case report to only Ecstasy use. Two reports have identified bilateral lesions in the globus pallidus of ecstasy users during magnetic resonance imaging (MRI) or autopsy, with a third report finding hippocampal changes in imaging associated with amnesic syndrome [488-490]. Due to the retrospective and infrequent nature of these reports, it is difficult to determine causality.

Overall, the risks of serious reports appear to be minimal in controlled settings with adequate screening with eligibility criteria defined in study protocols. None of these events have occurred within the context of human clinical studies with MDMA.

#### 4.6 Abuse Potential in Nonclinical Studies

Studies in Ecstasy users and animals suggest MDMA possesses some abuse potential, but not nearly that of amphetamine. Mice, rats, and monkeys self-administer MDMA, indicating that MDMA has rewarding properties in animals [509-511]; however, the rate and response-acquisition of self-administration is much lower than other drugs of abuse, such as cocaine or heroin. In rodents, acquisition of MDMA self-administration requires a lengthy training period with consecutive sessions [510, 512, 513]. Physical dependence and drug withdrawal was investigated by treating mice with 10 mg/kg i.p. MDMA twice daily for 5 days. Results showed that mice did not exhibit aversive/dysphoric or anxiogenic behaviors after treatment, indicating that high doses of MDMA do not induce classical symptoms of physical dependence [514]. Monkeys choose to self-administer MDMA in doses equivalent to or only slightly higher than doses used by humans [509], but typically reduce their MDMA intake over time. While monkeys work hard to obtain MDMA, they work harder to obtain other psychostimulants, such as cocaine or methamphetamine [512, 513]. Taken together, results in animals suggest that the abuse liability of MDMA is moderate.

Drug discrimination studies investigating the discriminative stimulus effects of MDMA as either hallucinogenic or stimulant have reported inconsistent findings. Some drug discrimination studies have shown MDMA to completely substitute for S-(+)-amphetamine in rats [634], monkeys [635], and pigeons [636]; where as other reports did not [637]. In a two-lever procedure, MDMA did not substitute for the hallucinogens (+)-lysergic acid diethylamide (LSD) or (+)-2,5-dimethoxy-4-methylamphetamine (DOM) [66, 638-639]. A three-lever procedure found that LSD produced dose-dependent increased substitution for MDMA while neither cocaine nor 2,5-dimethoxy-4-bromoamphetamine (DOB) substituted for it [640]. Serotonin and dopamine may be involved in producing stimulus characteristics in rats [641]. On the other hand, MDMA has been shown to substitute for mescaline [638]. Given MDMA's unique pharmacological profile and its ability to produce stimulant-like, mild hallucinogen-like, and empathogenic effects, in 1986 Nichols coined a novel pharmacological class, the 'entactogens' [66].

Research of Ecstasy dependence comes from a combination of published case studies and assessment of symptoms based on the Composite International Diagnostic Interview,



the Diagnostic and Statistical Manual of Mental Disorders Version IV (DSM-IV), and/or the Severity of Dependence Scale [521]. Of the small number of individuals assessed in a representative sample of Munich residents aged 14 to 24, only 1% were diagnosed with Ecstasy abuse and 0.6% with dependence [520], though other reports of large (N=173) but non-representative samples, including subjects recruited from substance abuse programs, reported 30% (N=52) had used Ecstasy and of these, 43% met DSM-IV criteria for dependence [519]. In a large Australian sample (N=329), approximately 25% of polydrug users wanted to reduce their Ecstasy use and 20% had received treatment for an Ecstasy-related problem, although this sample likely had an over-representation of chaotic intravenous polydrug users [522]. In a study of self-reported cravings in Ecstasy users, exposure to Ecstasy-related cues induced greater subject ratings of craving. Although over 50% of subjects agreed on some level with two or more statements regarding Ecstasy-related craving, the average score for craving was negative [523]. It also appears that MDMA has fewer or less intensely rewarding effects than stimulants, and even heavy Ecstasy users fail to report the intensive patterns of use seen with other stimulants [2, 4, 515]. Based on two structural analyses, Ecstasy dependence is bifactorial [517]. Although Ecstasy dependence does have a compulsive use factor as well as an escalating use factor, withdrawal symptoms do not include significant physical symptoms such as alcohol, cocaine, methamphetamine, opioids, and tobacco [516, 518]. In a prospective longitudinal study (N=2446), German polydrug users reported low prevalence of initial Ecstasy abuse or dependence, as well as substantial decline in use factors at 12-month follow-up, suggesting that Ecstasy use is a self-limiting transient phenomenon in many cases [520]. Features of Ecstasy abuse and dependence in humans are consistent with preclinical findings in self-administration studies of moderate abuse liability that is greater than that for serotonergic hallucinogens, but less than that for stimulants [510, 524].

## 5.0 Effects in Humans in Clinical Settings

### 5.1 History of Use in Clinical Settings

Shulgin and Nichols were the first to report on the effects of MDMA in humans [59]. In the 1970s, psychotherapists used MDMA-assisted psychotherapy to treat psychological disorders, including anxiety [65]. Legal therapeutic use continued until its placement on the U.S. list of Schedule I drugs in 1985 [64, 68, 525]. An estimated 500,000 doses of MDMA were administered during psychotherapy sessions in North America prior to its scheduling [57, 525]. A few uncontrolled human studies of MDMA occurred in the 1980s [44, 62], including Greer and Tolbert's study of MDMA in a psychotherapeutic context.

Controlled human studies of MDMA commenced in the mid-1990s with a MAPS funded investigator-initiated Phase 1 dose-response safety study [47, 526]. MAPS also funded a Phase 2 investigator-initiated dose-response safety and efficacy pilot study in Spain that was terminated early due to political concerns. This study enrolled six subjects, with four receiving a single session of MDMA-assisted psychotherapy without any safety concerns and experiencing some PTSD symptom reduction [527].

Based on past reports of MDMA use, preclinical studies and the results from these investigator-initiated trials with MDMA, the sponsor launched a Phase 2 Clinical Development Program in 2001 to develop MDMA-assisted psychotherapy for the treatment of chronic PTSD under U.S. IND. Eight sponsor-supported Phase 2 studies of MDMA-assisted psychotherapy for PTSD have been conducted. Two have been published, one main study with an extension in three subjects who relapsed in the U.S. (MP-1, MP1-E2) [41, 42], and one in Switzerland (MP-2)[43]. Four additional studies have completed treatments (MP-4, MP-8, MP-12) and are in follow-up, one

study in Israel was terminated early (MP-3) and re-initiated with a new study team (MP-9) and has completed enrollment.

MP-1, the first Phase 2 proof of principle study, explored the effect of MDMA-assisted psychotherapy for PTSD with a 125 mg initial dose and 62.5 mg supplemental dose of MDMA, as compared with inactive placebo in a chronic PTSD population (N=23). MP-1 enrolled eighteen women and five men, all European-American, average age  $41.3 \pm 7.1$  years. Subjects had no history of major medical conditions, psychotic disorders, dissociative identity disorder, or borderline personality disorder. Safety data obtained included: cognitive function before and after study participation, vital signs, liver panels, psychological distress during experimental sessions, concomitant medications, and AEs. Two subjects experienced unrelated SAEs, including a fractured clavicle from a motor vehicle accident and vasovagal syncope nearly 2 months after the second and final MDMA administration. Three MP-1 subjects relapsed after treatment, two of the them during the 3.8-year follow-up period and one after the follow-up. These three subjects were enrolled in an extension study, MP1-E2, to understand if a single MDMA-assisted psychotherapy session would improve PTSD symptoms after a relapse. The study has been completed. One subject experienced an unrelated SAE, a major depressive episode with suicidal ideation. MP-1 and MP1-E2 are now complete.

MP-2, the second Phase 2 proof of principle study, was conducted in Switzerland (N=14). This study explored reproducibility of MDMA-assisted psychotherapy for PTSD with a 125 mg initial dose and 62.5 mg supplemental dose of MDMA, as compared with 25 mg active placebo initial dose and 12.5 mg supplemental dose of MDMA (N=14). MP-2 enrolled 11 women and three men, average age  $41.8 \pm 10.9$  years. Most were of European ethnicity, one woman was South African and one man was Middle Eastern. Subjects enrolled had no psychotic disorders, dissociative identity disorder, or borderline personality disorder. One subject had a previous history of breast cancer, but had been in remission for over 10 years and was not symptomatic at screening. Safety data obtained from this study included: vital signs and psychological distress during experimental sessions, liver panels before and after treatment, concomitant medications, and AEs. One subject was diagnosed with a metastatic brain tumor during follow-up that resulted in death, which was an unrelated SAE. A second subject was hospitalized prior to dosing for psychiatric crisis, also reported as an unrelated SAE. MP-2 is now complete.

MP-3, the third Phase 2 study, was conducted in Israel with two Israeli therapist teams. This study was designed to explore reproducibility of MDMA-assisted psychotherapy for endemic PTSD with a 125 mg initial dose and 62.5 mg supplemental dose of MDMA, as compared with 25 mg active placebo initial dose and 12.5 mg supplemental dose of MDMA (N=5). MP-3 enrolled five male subjects, average age  $39.4 \pm 15.9$  years, with PTSD symptoms that failed to respond to at least one course of psychotherapy or at least one course of pharmacotherapy. Two subjects were Middle Eastern and three were European. This study was terminated early due to personnel turnover at the clinical site and difficulty of ensuring consistent training of site staff. These subjects are included in demographics data, and excluded from all other data due to inconsistencies in data collection. No SAEs or severe AEs were reported in this study.

There are three Phase 2 studies currently in follow-up (MP-8, MP-12, MP-4) and one that is completing treatments (MP-9). These studies explore the reproducibility of treatment outcomes of MDMA-assisted psychotherapy in people with chronic PTSD that failed to respond to at least one course of psychotherapy or at least one course of pharmacotherapy. Two of the randomized, blinded studies are taking place in the U.S. MP-8 (N=26) compares 30 mg versus 75 mg versus 125 mg initial dose of MDMA, with an optional supplemental dose equivalent to half the initial dose, in military veterans, firefighters and police officers ("first responders") with service-related PTSD, with an average age of  $37.2 \pm 10.3$  years. MP-12 (N=28) compares 40 mg versus 100 mg



versus 125 mg initial dose of MDMA, with an optional supplemental dose equivalent to half the initial dose, in subjects with PTSD from any cause, with an average age of  $42.0 \pm 12.9$  years. The Canadian study MP-4 (N=6) compares placebo to 125 mg initial dose of MDMA, with an optional supplemental dose equivalent to half the initial dose, in subjects with an average age of  $47.7 \pm 6.0$  years, and MP-9 (N=10) in Israel compares an initial dose of 25 mg to 125 mg MDMA, with an optional supplemental dose equivalent to half the initial dose, in subjects with an average age of  $36.7 \pm 8.0$  years.

The sponsor is also supporting two additional Phase 2 studies of MDMA-assisted therapies in parallel indications: one for treatment of social anxiety in autistic adults (MAA-1, N=12), and another for anxiety associated with a life-threatening illness (MDA-1, N=18). Subjective effects, mood, and reactions are also being assessed in the ongoing Phase 1 placebo-controlled study of MDMA-assisted psychotherapy, in healthy volunteers who have completed training in manualized MDMA-assisted psychotherapy (MT-1).

In sponsor-supported studies, MDMA or placebo/comparator is administered after preparatory psychotherapy during two or three 8-hour experimental sessions scheduled 2 to 5 weeks apart, each followed by at least three sessions of integrative psychotherapy. This treatment model is based on historical experience with MDMA use as an adjunct to psychotherapy.

Most data reported is from the Phase 2 studies of MDMA-assisted psychotherapy for PTSD. The studies have employed a range of comparator and active doses, from an initial dose of 25 mg to 150 mg MDMA. The highest dose (150 mg) was offered to a limited number of subjects in MP-2 as part of "Stage 3," an open-label arm for non-responders in Stage 1 and/or Stage 2. All studies have employed 125 mg usually followed 1.5 to 2 hours later by a supplemental dose of 62.5 mg MDMA as the primary active treatment.

The effects in humans presented in the sections below will include findings from both sponsor-supported clinical trials in patient populations as well as studies conducted in controlled laboratory settings in healthy volunteers without sponsor support. Findings from extensive human research being conducted on the pharmacology and mechanism of action will be presented in addition to the information required by FDA in order to support the safety profile of MDMA.

## 5.2 Pharmacology in Humans

As of 2015, the sponsor has not conducted studies on the pharmacodynamics or pharmacokinetics of MDMA, but relies on published literature. Beginning in the early to mid-1990s, several research teams conducted studies of the pharmacodynamics and pharmacokinetics of MDMA [10, 14, 22, 29, 116, 327, 528-530] without receiving sponsor support. Findings from these teams are described below, with specifics of metabolism detailed in [Section 5.2.1 Pharmacokinetics](#).


### 5.2.1 Pharmacokinetics

Onset of MDMA effects occurs 30 to 60 minutes after administration [8, 9], peak effects appear 75 to 120 minutes post-drug [7, 10-12], and duration of effects lasts from 3 to 6 hours [10, 12, 13], with most effects returning to baseline or near-baseline levels 6 hours after final drug administration. Self-reported duration of effects may increase as the dose of MDMA increases [7]. Administering a second dose of MDMA 2 hours after the initial dose, twice that of the initial dose, does not significantly extend the duration of measureable physiological or subjective effects [328]. Orally administered MDMA has a half-life of 7 to 8 hours in humans, with one report listing a half-life of 11 hours [531], and half-life is marginally extended if an additional dose is administered 2 hours after an initial dose [328]. Metabolites of MDMA are summarized in [Figure](#)

1 [532-537]. Metabolites are primarily excreted as glucuronide and sulfate conjugates [534]. Studies examining metabolism of 100 mg MDMA reported excretion values similar to those reported by de la Torre and associates [118, 531, 538-540]. MDMA and its metabolite MDA appear in oral fluid samples at much higher concentrations than plasma, for 24 to 48 hours for the former and 12 to 47 hours for the latter after oral administration of 1 to 1.6 mg/kg MDMA [541]. Urinary excretion of the MDMA metabolite HHMA after 100 mg MDMA in four men was  $91.8 \pm 23.8$  mol and 17.7% recovery [540]. By contrast, urinary recovery of the major metabolite HMMA after 100 mg was 40% [542]. As was the case for maximal plasma values, urinary recoveries for MDMA and MDA were higher when a second dose of 100 mg MDMA was administered 24 hours after an initial dose of 100 mg MDMA when compared with a single dose [118]. In one study, urinary excretion of the metabolite HMMA exceeded that of MDMA by 33 hours after a dose of 1.6 mg/kg MDMA [543], suggesting that secondary metabolism of MDMA continues during this period. Findings support the enantioselective nonlinear metabolism of MDMA and its metabolites measured in blood and urine [544, 545].

A study comparing the effects of a single 100 mg dose with an initial administration of 50 mg followed 2 hours later by 100 mg reported higher peak plasma MDMA than might be expected, and lower levels of the MDMA metabolites HMMA and HMA [328], findings further supported by examining plasma MDMA after two doses of 100 mg given 4 hours apart [546], likely due to metabolic autoinhibition. Comparison of pharmacokinetic-pharmacodynamic relationships for MDMA reveals acute pharmacodynamic tolerance. Despite 8 hours of plasma half-life of MDMA, and persistent high drug levels in the blood, most pharmacodynamic effects of the initial dose rapidly return to baseline within 4 to 6 hours [530]. These findings suggest that intensity of most subjective and physiological effects of MDMA would not be significantly impacted by the supplemental doses in sponsor-supported studies due to acute tolerance to its prototypical effects [546]. This acute tolerance could be caused by functional depletion of stores of serotonin so that no more can be released despite MDMA still being present [530], or suggests that MDMA transport into intracellular spaces is saturable due to limited transport capacity [127]. Additionally, reversible inhibition of tryptophan hydroxylase as observed in rodents [20], or internalization of serotonin reuptake transporters from the plasma membrane leading to less serotonin release [78], would support self-limiting effects of MDMA. On the other hand, although SERT can be internalized, evidence suggests that accumulation of extracellular serotonin stimulated by MDMA affects SERT trafficking by perpetuating cell-surface SERT expression, but in contrast promotes internalization of DAT and NET [127, 547].

**Figure 1: Metabolism of MDMA in Humans**



Metabolism of MDMA in humans (in red) compared to metabolism in rats (in blue). Reproduced with permission of R. de la Torre [113].

MDMA is metabolized in the liver by several cytochrome P450 CYP enzymes, including CYP1A2, CYP3A4, and CYP2D6. It is likely that active doses of MDMA inhibit CYP2D6 function, as measured by examining the effects of MDMA on dextromethorphan metabolism. Inhibition of CYP2D6 by MDMA was demonstrated first in a physiological model derived from data collected after oral administration in humans [548]. O'Mathuna and colleagues present evidence that CYP2D6 activity may not fully recover until 10 days after MDMA [549, 550]. After reviewing their data and the literature on MDMA pharmacokinetics, de la Torre and colleagues concluded variation in CYP2D6 genotype is not clinically significant, due in part to the fact that the enzyme is inhibited in most people after administration of an active dose [327]. In contrast, MDMA may produce increased activity of the enzyme CYP1A2, as evidenced by comparing caffeine metabolism before and after MDMA [551]. The enzyme COMT and monoamine oxidase may also be involved in the metabolism of MDMA [542]. At least one variation in COMT genotype may affect MDMA elimination rate ( $K_e$ ) and systolic blood pressure (SBP) after MDMA [552]. As a monoamine reuptake inhibitor that leads to monoamine release and inhibits monoamine oxidase-A [132] combining MDMA with a monoamine oxidase inhibitor (MAOI) medication presents a risk for provoking serotonin syndrome and increases in sympathetic activity. Fatalities have occurred apparently as a result of combining MAOI medications with MDMA [133, 134]. For this reason, MAOI medications are tapered for at least five half lives of the medication and active metabolites, plus 1 week for symptom stabilization in sponsor-supported studies.

Researchers have attempted to compare MDMA pharmacokinetics in humans and other species, including other primates, as discussed in Section 4.2.1 Pharmacokinetics in Animals and Section 5.2.1 Pharmacokinetics. These investigations sought to establish human-equivalent doses given nonlinear pharmacokinetics. Doses that researchers assumed to be human-equivalent produced greater plasma concentrations. However, duration of exposure expressed in half-life was often shorter. For example, a dose of 1.6 mg/kg MDMA produced a half-life of 8.4 hours in a small sample of humans while a dose of 2.8 mg/kg had a half-life of 2.1 hours [119]. A dose of 7.4

mg/kg in squirrel monkeys, four times a human-equivalent dose and never administered in a human trial, had a half-life of 3.4 hours [107]. Researchers have detected nonlinear pharmacokinetics of MDMA in all species studied to date, leading Mueller and colleagues to conclude that a preclinical study cannot accurately and simultaneously model human-equivalent plasma levels and equivalent duration of exposure [119].

### 5.2.2 Pharmacodynamics

Estimates from animal data suggest the LD50 in humans is probably between 10 to 20 mg/kg [6]. Typically, human trials have used doses between 1 and 2 mg/kg, with therapeutic studies using fixed dosing rather than adjusting dosing on a mg/kg basis, in order to achieve a more consistent subjective response between subjects. The pharmacokinetics of MDMA in humans have been characterized in blood and urine samples using oral doses of up to 150 mg MDMA [14]. MDMA is a triple monoamine reuptake inhibitor, and similar drugs in this class have been found to exert potent anti-depressant activity with a potentially favorable safety profile [15, 16]. MDMA concomitantly promotes release, inhibits reuptake, and extends duration of serotonin, norepinephrine, and dopamine in the synaptic cleft to increase serotonergic, noradrenergic, and dopaminergic neurotransmission. MDMA has self-limiting subjective and physiological effects as previously described.

Many researchers categorize MDMA as belonging to a unique class of drugs referred to as the Entactogens [13, 66], defined as substances that produce changes in mood and social interaction, as well as feelings of interpersonal closeness and changes in perception. MDMA shares some of the pharmacological effects of stimulants and serotonergic hallucinogens [8, 10, 11, 553], as well as a small number of pharmacologically related compounds, such as MDE [553]. Initially, narrative reports and surveys supported the social cognitive effects of MDMA or Ecstasy [2, 234, 235, 554]. Controlled trials detected self-reported empathy or closeness to others in healthy volunteers [7, 12, 91], and starting in the late 2000s to 2010s, controlled studies measured effects of MDMA on social cognition or emotion [29, 30, 35]. Although researchers have offered several models and explanations for the effects of Entactogens, it appears that serotonin and norepinephrine release play a significant role in producing at least some of these effects. Indirect action on 5HT<sub>1A</sub> or 5HT<sub>2A</sub> receptors and neuroendocrine responses such as increases in the hormones oxytocin, AVP, prolactin, and cortisol may also play a role in producing the unique effects of MDMA.

In addition to neuroendocrine and norepinephrine-mediated effects, MDMA may target similar binding sites on the SERT, as do already approved PTSD medications Paxil and Zoloft, which are both SSRIs. Similar to the SSRI Prozac, MDMA also inhibits MAO-A to extend presence of serotonin in the synaptic cleft [132]. Pre-treatment or co-administration studies of SSRIs with MDMA appear to attenuate or eliminate most subjective, physiological and immunological effects of MDMA due to competition for binding sites on the SERT which may prevent transporter-mediated serotonin release [91, 555-558]. Pre-treatment or co-administration with SSRIs attenuates serotonergic effects of MDMA on mood and perception, without influencing specific effects, such as nervousness or excitability [555]. Some researchers report that SSRIs attenuate MDMA-induced increases in heart rate and blood pressure [91, 556], while others report that SSRIs only attenuate elevated heart rate [558]. Additional effects of each SSRI beyond reuptake inhibition on production, release, and degradation of serotonin are likely responsible for variations between SSRI co-administration findings. All three studies of SSRI pre-treatment suggest that co-administration of SSRIs with MDMA is safe, but the combination prevents or significantly reduces the subjective effects of MDMA. The role of serotonin release on the potentially therapeutic effects of MDMA-assisted psychotherapy has yet to be investigated, however reduced feelings of sociability and closeness to others after paroxetine pre-

administration suggests that serotonin release is at least partially involved in prosocial effects that are thought to be therapeutically relevant [91]. These subjective effects are predominately mediated by direct or indirect action on 5HT<sub>2A</sub> receptors [92, 233, 559], with at least one study concluding that the effects of MDMA upon positive mood are at least due in part to 5HT<sub>2A</sub> receptor activation [92]. In contrast, the 5HT<sub>1A</sub> receptor appears to be partially involved in producing the subjective effects of MDMA [92, 231-233]. Co-administration of the beta-blocker and 5HT<sub>1A</sub> antagonist, pindolol, along with 1.6 mg/kg MDMA to 15 men attenuated self-reported “dreaminess” and pleasantly experienced derealization after MDMA without attenuating MDMA-related reduction in performance on a task requiring visual attention, and co-administration of pindolol failed to alter the acute effects of 75 mg MDMA on self-reported mood [92, 231].

Human MDMA studies suggest that norepinephrine release also contributes to the pharmacodynamic, physiological and psychological effects of MDMA [205, 208, 560, 561]. Tricyclic antidepressants, as well as many of the current antidepressant medications, are known to promote norepinephrine signaling, as does MDMA. Studies with the norepinephrine uptake inhibitor reboxetine, and the  $\alpha_1$ -adrenergic receptor antagonist doxazosin, suggest that norepinephrine plays a role in the effects of MDMA on blood pressure and subjective effects of positive mood and excitement [206, 560], but not in “entactogenic” or “empathogenic” effects. Most of the psychostimulant-like and psychological effects of MDMA are blocked after administration of the dual selective Serotonin and norepinephrine uptake inhibitor (SNRI) duloxetine [208, 561]. There is evidence that norepinephrine and serotonin may play a role in the elevation in the neuroendocrine hormone copeptin, the C-terminal precursor of pre-pro-AVP, detected in women acutely after MDMA administration [561]. Some *in vitro* findings with human monoamine transporters expressed in cells indicate that MDMA displays a higher affinity for the NET than the serotonin or dopamine transporter, while still producing greater detectable release of serotonin versus norepinephrine, suggesting a role for both transmitter systems [127]. As the NET unexpectedly has a greater affinity than the DAT for dopamine, it preferentially clears dopamine in brain areas where there is a greater concentration of NET, such as the frontal cortex [562]. The relative affinities of MDMA for various monoamine reuptake transporters, and the affinity of the respective transporters for each neurotransmitter, can thus influence the selectivity of signaling pathways MDMA activates in a region-specific manner depending on transporter density and availability.

Some MDMA effects on human mood and anxiety may be attributed to dopamine release based on the finding that pretreatment with haloperidol, a dopamine receptor antagonist with partial selectivity for the D<sub>2</sub> receptor subtype, diminished MDMA-induced positive mood and increased anxiety [563]. However, the control group receiving haloperidol alone also experienced dysphoric mood, suggesting that this finding may overestimate the dopaminergic effects of MDMA. Studies comparing MDMA with the dopaminergic and adrenergic drug methylphenidate (Ritalin) suggest that dopamine release and inhibition of uptake play a minor role, if any, in producing the effects of MDMA [34]. Co-administration of MDMA with the potent dopamine reuptake inhibitor methylphenidate neither enhanced nor attenuated the effects of MDMA [530]. MDMA, but not methylphenidate, increased trust, openness, and closeness to others. Co-administration of MDMA with the dopamine reuptake inhibitor bupropion prolonged, but did not reduce subjective effects of MDMA, supporting that dopamine does not have a part in MDMA effects on mood [564].

MDMA produces a robust increase in the neurohormone oxytocin [29], a finding first seen in a naturalistic study that reported elevated levels of oxytocin in clubgoers with detectable blood MDMA levels when compared to clubgoers without detectable levels of MDMA [32], as described in [Section 4.3.5 Neurobiological Effects](#). It is likely that all neuroendocrine changes are part of a signaling cascade downstream of monoamine release. Exogenous oxytocin increases trust and improves accuracy of emotion perception, and increased cortisol, in some



circumstances, may serve as a signal to seek affiliation or to increase positive mood [565-568]. However, studies comparing increases in empathy or prosocial effects of MDMA with intranasal oxytocin have failed to find indications that the two substances produce similar effects, with MDMA producing greater feelings of sociability and emotional empathy than oxytocin [63, 569]. Peripheral oxytocin has been suggested to be a reliable indicator of central oxytocin, but peripheral effects of oxytocin need to be ruled out when assessing central effects [570]. The potential significance of elevated oxytocin in producing changes in social cognition are discussed in [Section 5.3.8.3 Social Effects](#), and include potentially therapeutic effects, such as increased feelings of closeness to others or greater ability to detect expressions of positive mood in others.

MDMA acutely increases cortisol, prolactin, and adrenocorticotrophic hormone concentrations in a dose dependent manner [9, 12, 19, 30, 47, 118, 526, 572-574], whereas growth hormone levels are unchanged by up to 125 mg MDMA [9]. Increases in cortisol and prolactin peak at about 2 hours after MDMA administration [9, 47]. A second dose of 100 mg MDMA, given 4 hours after an initial 100 mg, produces a second increase in cortisol during an interval when cortisol levels are declining [575], and a dose of 100 mg MDMA, given 24 hours after an initial dose, stimulates a greater release of cortisol but not prolactin [118]. In a study of the effects of 0.5 and 1.5 mg/kg MDMA in eight people, there was a trend for increased levels of the hormone dehydroepiandrosterone (DHEA) after 0.5 mg/kg MDMA, and a significant increase after 1.5 mg/kg MDMA, with peak levels appearing 2 to 3 hours post-drug [12]. A crossover study comparing the effects of MDMA and methylphenidate found that MDMA increased serum cortisol while methylphenidate did not, and that neither drug altered testosterone levels [574]. These findings suggest a relationship between serotonin release and increased serum cortisol. Pre-treatment with the cortisol synthesis inhibitor metyrapone blocked MDMA-induced increase in cortisol levels in blood without preventing impaired performance on verbal memory tasks or altering the effects of MDMA on mood [572]. A study investigating the emotional effects of MDMA found no correlation between those changes and the MDMA-induced increases in oxytocin, cortisol, and prolactin [573].

The pharmacological basis for reported acute shifts in memory, including impaired visual recall and improved recall for life events, after MDMA administration remains undetermined. Initial findings suggest a relationship between MDMA and activation of temporal areas in the brain and response to positive memories, as well as increases in medial PFC and response to negative memories [36]. It is possible that elevation in cortisol could be tied to specific acute effects on mood or memory. Another study found MDMA-associated changes in inferior parietal lobule and acute impairment in working memory [576]. Animal studies have postulated a role of Ach release triggered by upstream serotonin and dopamine neurons in MDMA-induced shifts in memory described above. A human study revealed no difference in MDMA-induced memory changes following pretreatment with the cortisol synthesis inhibitor metyrapone or the  $\alpha_7/nAChR$  receptor antagonist memantine, suggesting cortisol is not involved in these effects [572, 577]. It is unclear what contributions, if any, elevated neuroendocrine levels make to the subjective and memory effects of MDMA.

### 5.3 Safety of MDMA in Humans

Safety data from studies in controlled research settings show that MDMA produces sympathomimetic effects that include statistically significant, self-limiting increases in body temperature, heart rate, and blood pressure that are likely to be transient and well tolerated by healthy individuals [7, 9, 10, 12, 26, 41-47, 526, 527]. Risks posed by elevated blood pressure are addressed in clinical trials by excluding candidates with a history of cardiovascular or cerebrovascular disease or with pre-existing uncontrolled hypertension and by monitoring blood pressure and pulse during MDMA-assisted experimental sessions. Common reactions from



MDMA research studies are transient and diminish as drug effects wane during treatment sessions and over the next 24 hours. In studies conducted with and without sponsor support in controlled clinical settings, with 1180 individuals exposed to MDMA, there have been no published or reported unexpected drug-related SAEs to date, and expected SAEs have been rare and non-life threatening. One subject to date experienced an expected related SAE (increased premature ventricular extrasystoles in MP-8), and 10 unrelated SAEs after drug administration have been reported in MAPS-sponsored clinical trials.

All sponsor-supported data presented in this IB was collected through 01 October 2015. There are three completed (MP-1, MP-2, MP1-E2) and four ongoing Phase 2 studies of MDMA-assisted psychotherapy in people with PTSD that have completed enrollment (MP-8, MP-12, MP-4, MP-9). A Phase 2 study of MDMA-assisted therapy treating social anxiety in autistic adults (MAA-1) and another Phase 2 study of MDMA-assisted psychotherapy treating anxiety associated with life-threatening illness (MDA-1) are ongoing. Safety is addressed and closely monitored through several measures in these studies. Vital signs, concomitant medications, unexpected and expected AEs are collected in all studies. Suicidal ideation and behavior are formally measured with the Columbia Suicide Severity Rating Scale (C-SSRS) in all but MP-1 and MP-2. One completed (MP-1) and two ongoing studies (MP-12, MP-4) measure cognitive function before and after treatment. Psychological distress during psychotherapy sessions is assessed in all studies with the single-item Subjective Units of Distress (SUD) scale.

Partial safety data from the Phase 1 study MT-1 in healthy volunteers is not presented in the current report since data remains blinded. There have been no severe or serious AEs during the study, and there were no clinically significant changes in vital signs. No medical intervention has been required during this study to date.

Physiological effects of MDMA-assisted psychotherapy in sponsored studies are similar to those reported in studies conducted outside of sponsor support, including elevated blood pressure, body temperature, and heart rate. The following common reactions are found in published literature and are collected in the sponsor's Phase 2 clinical trials: anxiety, depressed mood, insomnia, obsessive rumination, restlessness, irritability, headache, disturbance in attention, dizziness, paresthesia, judgment impaired, hypersomnia, nausea, diarrhea, fatigue, asthenia, feeling cold, muscle tightness, decreased appetite, hyperhidrosis, disturbed gait, dry mouth, thirst, sensation of heaviness, somnolence, and nystagmus. These common reactions are transient and diminish as the drug is metabolized during treatment sessions and excreted over the next 24 hours, with the majority of reactions resolving within several days and up to 1 week after dosing. Among spontaneous reports of reactions to MDMA, muscle tightness (jaw), anxiety, decreased appetite, headache, and fatigue were most commonly reported acutely during MDMA-assisted psychotherapy. During the week following treatment, the most frequently reported reactions were anxiety, fatigue, insomnia, depressed mood, and hypersomnia. The half-life of MDMA doses used in these studies is 8 to 9 hours and the majority of AEs have been transient, resolving within 2 to 3 days after MDMA has been metabolized and excreted. Severe anxiety, insomnia, fatigue, nausea, muscle tightness, and depressed mood are commonly reported in PTSD studies supported by the sponsor. These reactions also overlap with symptoms of pre-existing conditions in medical history associated with PTSD (depression, somatic symptoms, insomnia, anxiety), which may influence the reaction frequency observed during clinical trials of MDMA-assisted psychotherapy.

### 5.3.1 Reproductive and Developmental

All research studies with MDMA, with and without sponsor support, require measures to limit pregnancy risk prior to receiving each dose of MDMA. Women of childbearing potential must

use an effective method of birth control to be enrolled in sponsor-supported studies, and pregnancy tests must be negative prior to each experimental session. There is no information on reproductive and developmental risks reported as there have been no pregnancies in these studies. See [Section 4.4.5 Reproductive and Developmental Toxicity](#) for information gathered on reproductive and developmental risks in Ecstasy users.

### 5.3.2 Immunological Effects

Various groups have studied immunological effects of MDMA in laboratory settings, with none found to be clinically significant from a safety standpoint. Studies in men conducted by researchers in Spain have found 100 mg MDMA to have immunosuppressive and anti-inflammatory effects [117, 557, 575, 578, 579]. Findings included a decline in CD4 cells, smaller CD4/CD8 ratio, attenuated lymphocyte proliferation in response to mitogen, and an increase in natural killer (NK) cells, with effects diminishing but still detectable 24 hours after drug administration. These researchers also found that MDMA decreased production of pro-inflammatory cytokines, including IL-6, IL-1 $\beta$ , TNF- $\alpha$ , and INF- $\gamma$ , and increased production of anti-inflammatory cytokines, including IL-10 and TGF- $\beta$ . Generally, MDMA appeared to decrease the concentration of Th1 cytokines, including IL-2, and increase the amount of Th2 cytokines, including IL-4, measured in blood. Changes of similar magnitude and duration have been previously noted after ingestion of other psychoactive agents, such as alcohol or cocaine [117, 579]. Due to their limited duration, these changes are not likely to have clinical significance beyond several days of possible increased risk of viral upper respiratory infection or similar illness. Interestingly, meta-analysis and meta-regression of 20 studies investigating inflammatory markers in PTSD found an association with increased IL-6, IL-1 $\beta$ , TNF- $\alpha$ , and INF- $\gamma$ , consistent with chronic low-grade inflammation [184], and any effects of MDMA on these immune markers remains to be tested.

Immunological changes seen after an initial dose of MDMA are enhanced by a second dose of identical size given 4 hours after the first dose [575, 580]. A second dose of identical size given 24 hours after the first dose produced the same immunological effects over the same time course, but with greater intensity than after the first dose [575]. Given this data, it is possible that administering a smaller supplemental dose 1.5 to 2.5 hours after the first dose will slightly enhance the immunological effects set in motion by the initial dose of MDMA. Previous Phase 1 studies mentioned above have not reported any indication of increased risk of illness occurring after MDMA administration.

### 5.3.3 Thermoregulatory Effects

In the first Phase 1 safety study funded by the sponsor, MDMA was found to cause a significant increase in body temperature in some healthy volunteers [47]. However, these increases were found to be transient and tolerable in a controlled clinical setting. Doses between 1.5 and 2 mg/kg produced only a slight elevation in body temperature that was not clinically significant [10, 556, 559] and this elevation was unaffected by ambient temperature [195]. Studies in MDMA-experienced volunteers given 2 mg/kg MDMA produced slight but statistically significant increases in core body temperature, at mean elevation of 0.6°C [195]. The same study found that ambient temperatures did not affect elevation in core temperature after administration of MDMA, which increased metabolic rate. A supplemental dose twice as large as the initial dose of MDMA elevates body temperature, but not beyond what would be expected after the cumulative dose [328]. While MDMA did not increase or decrease perspiration overall in this study, it was associated with a higher core temperature when perspiration began. Ambient temperature neither attenuated nor amplified the subjective effects of MDMA, with people reporting similar drug effects in warm and cool environments. As expected, people felt warm when the room was warm

and cold when the ambient temperature was cool, and MDMA did not distort perceptions of warmth or cold in either case. Unlike rodents given MDMA at higher mg/kg doses, humans do not exhibit reduced temperature when MDMA is given in a cold environment, and they do not exhibit significant hyperthermia in a warm environment. When compared with placebo, findings from 74 subjects given MDMA found that men exhibited a greater elevation in body temperature than women when given the dose of MDMA in mg/kg [10]. Subsequent studies have not confirmed this gender difference [26], and a report in a sample of 17 men and women reported higher oral temperatures in women [552]. A review of clinical placebo-controlled laboratory studies conducted without sponsor support found that route of measurement has an effect on variability in body temperature findings, with oral and tympanic, but not axillary, temperatures frequently rising above 38°C into moderate hyperthermia ranges at 125 mg MDMA [581]. Thermogenic effects of MDMA are distinct from malignant hyperthermia and are mediated by noradrenergic signaling, which contributes to peripheral effects of MDMA by affecting cutaneous vasoconstriction of blood flow and stimulation of heat production, and are attenuated by norepinephrine blocking drugs [582]. It is notable that subjects in studies in a clinical setting have not engaged in vigorous exercise and have remained either sitting or lying down throughout duration of drug effects. It may be the case that heat dissipation impaired by a hot environment, heat generation increased by exertion, interactions of serotonergic drugs, and potential disturbance of central heat regulation mechanisms contribute to the occurrence of hyperprexia (body temperatures >41°C) in people ingesting Ecstasy in uncontrolled settings. However, one of four naturalistic studies reported that Ecstasy users had a statistically significant increase in body temperature [583], while three others failed to find significant differences in Ecstasy-user body temperature at a club [584-586].

In all sponsor-supported studies to date, oral body temperature readings were taken at baseline, then every 60 to 90 minutes, with some differences in collection methods across studies. Peak values during each experimental session are ascertainable for all studies. Across studies, the final value was either at a relatively set time (MP-8, MP-12, MP1-E2) or as the final reading with time point varying (MP-1). MP-1 and MP-2 reported two pre-drug values (15 minutes and 5 minutes before dosing) and these were averaged. Average post-drug values serve as the final value for MP-2. If body temperature rose 1°C above the pre-drug reading, each duration above the pre-determined cut-off was collected in MP-2, MP-8, MP-12, MP-9, MP-4, MP1-E2, MAA-1, and MDA-1. Clinical signs and symptoms were monitored and more frequent readings were collected in cases where readings were above cut-off. Data presented below is final for completed studies and preliminary for ongoing studies.

**Table 3: Pre-Drug, Peak, and Final Body Temperature During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations**

Dose	Subjects (Observation)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with BT Above Cut-off (Observations)
0 mg	14 (27)	35.1/37.2 36.4 (0.5)	36.4/37.6 36.9 (0.3)	35.9/37.5 36.6 (0.3)	2 (2)
25 mg	8 (18)	35.8/37.1 36.5 (0.3)	36.0/38.5 37.2 (0.8)	36.0/38.0 36.9 (0.7)	4 (6)
30 mg	7 (15)	35.3/36.9 36.3 (0.5)	36.4/37.9 37.0 (0.4)	35.7/37.2 36.5 (0.4)	4 (6)
40 mg	7 (12)	35.6/37.2 36.4 (0.5)	36.6/37.6 37.1 (0.3)	36.5/37.6 37.0 (0.4)	3 (3)
75 mg	13 (20)	35.9/37.8 36.6 (0.4)	36.3/37.8 37.2 (0.5)	36.1/37.6 36.8 (0.4)	2 (2)
100 mg	25 (42)	33.9/37.5 36.1 (0.8)	35.5/37.9 37.0 (0.47)	34.8/38.0 36.7 (0.7)	8 (12)
125 mg	95 (232) <sup>A</sup>	34.3/37.7 36.5 (0.5) <sup>B</sup>	36.0/38.7 37.3 (0.5)	35.2/38.4 36.9 (0.5)	50 (83)
150 mg	3 (4)	36.6/36.7 36.7 (0.1)	37.3/38.2 37.7 (0.4)	36.8/37.7 37.3 (0.4)	1 (2)

<sup>A</sup> One endpoint temperature was excluded pending queries, and two listings are unavailable for endpoint temperature.

<sup>B</sup> One subject given 125 mg did not have pre-dose values for any vital sign, but post-drug values were collected.

Based on the literature, MDMA is expected to produce elevations in body temperature with possible influence of ambient temperature. Body temperature above 1°C above baseline was detected in 33% (114 of 343) of experimental sessions where MDMA was administered at any dose, and in 46% (72 of 157) of subjects in sponsor-supported trials. Maximum body temperature observed to date was 38.7°C in one MP-2 subject lasting 3 hours, where 125 mg MDMA was administered as the initial dose. This subject had no risk factors reported in medical history and temperature elevation was not clinically significant. Maximum duration above 1°C elevation was 9.2 hours in one MP-9 subject where 125 mg MDMA was administered as the initial dose. This subject experienced a maximum of 38.0°C temperature, which dropped to 37.6°C at final reading. By contrast, elevation of body temperature above 1°C was observed in 7% (2 of 27) of experimental sessions and in 14% (2 of 14) of subjects receiving inactive placebo. Perspiration was reported in 21% to 25% of experimental sessions with active dose MDMA, and was generally mild. Adjustments were made to the ambient temperature and to air circulation in the room, but no subjects required medical intervention to decrease body temperature, and values returned to baseline as drug effects waned. In conclusion, controlled setting for treatments with MDMA-assisted psychotherapy are optimized with the capacity to control ambient temperature for subject comfort, though there is no evidence that this will significantly influence or is needed for control of core body temperature.

### 5.3.4 Cardiovascular Effects

MDMA produces sympathomimetic effects that include elevation in blood pressure and heart rate, first recorded by Downing [44] and replicated by other research teams in the U.S. and Europe [9, 10, 45]. Subsequent trials confirmed that MDMA produced significant increases in heart rate and blood pressure that were likely to be well tolerated by healthy individuals [7, 12, 26, 46]. Most people do not experience elevations that are greater than those seen after moderate

exercise. MDMA has also been found to decrease respiratory sinus arrhythmia, the natural variation in heart rate over the course of each respiratory cycle [587]. Cardiovascular effects of MDMA first appear 30 to 45 minutes after administration [44] and peak between 1 and 2 hours post-drug [11, 45], with effects waning 3 to 5 hours after drug administration. Men given the same mg/kg dose of MDMA as women exhibited a significantly greater elevation in blood pressure and heart rate in a study summarizing and pooling data from a series of human MDMA studies [10]. These studies did not report any discomfort or increased distress accompanying cardiovascular effects.

Elevation in blood pressure above 140/90 occurred in approximately 5% of research subjects receiving a single dose of at least 100 mg of MDMA in Phase 1 research studies [9, 13]. Peiro and colleagues observed elevation in blood pressure above 150/90 as well in all 10 subjects given 50 mg followed 2 hours later by 100 mg MDMA [328]. When compared with 100 mg MDMA and placebo given 4 hours apart, two doses of 100 mg 4 hours apart significantly elevated SBP, while other physiological were not significantly elevated beyond values seen after a single dose. These studies used different dosing regimens than the one used in sponsor-supported studies, which employ an optional supplemental half dose. None of these individuals needed clinical intervention and blood pressure returned to normal as drug effects waned [9, 13, 328].

Greater elevations in blood pressure are seen in individuals with a specific COMT genotype (Val158/Met genotype), and greater elevations in blood pressure and heart rate are seen in individuals with a specific SERT (1/\* 5-HTTLPR) genotype [552]. However, the observed increases are not so severe as to suggest contraindication for these genotypes. The  $\alpha_1$ - and beta-adrenergic receptor antagonist carvedilol is capable of reducing MDMA-induced elevations in blood pressure, heart rate, and body temperature when administered 1 hour before MDMA without affecting the subjective effects of MDMA, indicating the norepinephrine release is primarily responsible for cardiovascular effects of MDMA [207]. Other concomitant antihypertensive medications either alter some of the effects of MDMA [588] or do not significantly reduce MDMA-induced blood pressure elevation [205].

Norepinephrine release induced by MDMA leads to indirect activation of the AVP system, stimulating secretion of copeptin (CTproAVP), a 39-aminoacid glycopeptide that is a C-terminal part of the precursor pre-proAVP. CTproAVP is secreted into circulation from the posterior pituitary gland in equimolar amounts with AVP. CTproAVP directly reflects AVP concentration and can be used as a surrogate biomarker of AVP secretion. In many studies CTproAVP behavior represents changes in plasma osmolality, stress and various disease states (diabetes, SIADH, heart failure, renal disorders), and is an indicator of osmoregulatory function in the body [365]. Heart failure is commonly associated with hyponatremia, and is also characterized by increased concentrations of basal AVP and CTproAVP [375]. Intra-cardiac pressures, intra-arterial pressures, angiotensin II, pain, and adrenergic ( $\alpha_2$ ) central nervous stimuli can also influence AVP secretion [364]. Increased CTproAVP concentration is described in several studies as a strong predictor of mortality in patients with chronic heart failure and acute heart failure. [365]. Taken together, the AVP system appears to be the main connection between MDMA and cardiovascular risk as well as hyponatremia.

In all sponsor-supported studies to date, blood pressure readings were taken at baseline, with study-specific differences in data collection times post-drug. Peak values during each experimental session are ascertainable for all studies. The final or endpoint was recorded as the final value, either at a relatively set time (MP-8, MP-12, MP1-E2) or as the final value available, or with timepoint varying (MP-1). MP-1 and MP-2 reported two pre-drug values (15 minutes and 5 minutes before dosing) and these were averaged, whereas all other studies reported single time point pre-drug. Average post-drug values serve as the final value for MP-2. If SBP rose above

160 mmHg or if diastolic blood pressure (DBP) rose above 110 mmHg, each duration above this pre-determined cut-off for more frequent measurement was collected in MP-8, MP-12, MP-9, MP-4, and MP1-E2. In MAA-1, if SBP rose above 180 mmHg or if DBP rose above 110 mmHg, each duration above the pre-determined cut-off was collected. If SBP rose above 180 mmHg and if DBP rose above 120 mmHg, each duration above the pre-determined cut-off is collected in MDA-1. MP-2 criteria for cut-off was exceeding both 160/110 mmHg. Clinical signs and symptoms were monitored and more frequent readings were collected in cases where readings were above cut-off. Data presented below is final for completed studies and preliminary for ongoing studies.

**Table 4: Pre-drug, Peak, and Final Systolic Blood Pressure During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations**

Dose	Subjects (Observations)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with SBP Above Cut-off (Observations)
0 mg	14 (27)	90/139 118.8 (13.0)	102/159 134.5 (16.3)	83/138 115.2 (13.5)	0
25 mg	8 (18)	110/130 119.9 (5.2)	117/147 133.6 (8.1)	107 /146 119.8 (11.3)	0
30 mg	7 (15)	94/134 114.2 (12.1)	110/155 132.3 (14.0)	98/140 118.5 (11.6)	0
40 mg	7 (12)	100/154 125.9 (14.1)	112/168 137.1 (17.7)	107/148 124.3 (12.1)	2 (2)
75 mg	13 (20)	101/145 124.2 (11.3)	116/179 144.7 (17.5)	107/156 127.8 (12.8)	3 (4)
100 mg	25 (42)	92/155 118.0 (13.4)	100/193 138.2 (22.8)	86/148 119.1 (14.6)	6 (8)
125 mg	94 (232) <sup>A</sup>	95/177 125.3 (14.9)	114/200 152.8 (17.4)	77/170 126.2 (15.7)	43 (78)
150 mg	3 (4)	102/146 128.0 (21.0)	128/185 156.5 (23.3)	117/161 141.0 (19.0)	1 (1)

<sup>A</sup> One subject given 125 mg did not have pre-dose values for any vital sign, but post-drug values were collected.

As described above, MDMA is expected to produce statistically significant but transient, self-limited increases in blood pressure. The supplemental half dose, when administered 1.5 to 2.5 hours after the initial dose, may cause further SBP increases above those resulting from the initial dose of MDMA. In one study (MP-1), 9 of 23 subjects received the supplemental dose, with four in the 125 mg MDMA group, in all subsequent studies, most of the subjects received the optional supplemental dose. A comparison of subjects receiving the supplemental dose to those who only received the initial dose in MP-1 indicate that the supplemental dose did not cause further elevation in blood pressure and heart rate beyond the initial dose, although the sample was underpowered to detect a small effect. Maximum SBP observed to date was 200 mmHg in a single MP-2 subject, lasting 5 hours, where 125 mg MDMA was administered as the initial dose. This subject had a medical history of controlled hypertension, and the traumatic event that caused PTSD was medical malpractice, with a secondary diagnosis of white coat hypertension. This subject was only enrolled after 24-hour monitoring of blood pressure at baseline to confirm this diagnosis. SBP above cut-off was detected in 27% (93 of 343) of experimental sessions where MDMA was administered, and in 35% (55 of 157) of subjects receiving MDMA in sponsor-supported trials. Maximum duration above SBP cut-off was 6 hours in two separate subjects with respective peak values of 172 and 174, where 125 mg MDMA was administered as the initial dose. Doses of 40 mg MDMA and greater were associated with elevations above cut-off. SBP



was elevated in 46% (43 of 94) of subjects and 34% (78 of 232) of experimental sessions where the 125 mg dose was administered. This was not observed in any of the sessions where inactive placebo or 25 mg to 30 mg MDMA was administered, supporting a dose dependent effect of MDMA on blood pressure. Despite elevations in SBP, no clinical signs or symptoms of hypertension were observed. In all cases, final values returned to pre-drug levels with no clinical intervention required. No clinically significant AEs were reported based on elevations in blood pressure.

**Table 5: Pre-drug, Peak, and Final Systolic Blood Pressure During Experimental Sessions in Controlled Hypertension Subjects in MAPS-Sponsored PTSD Study MP-8**

Dose	Subjects (Observations)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with SBP Above Cut-off (Observations)
30 mg	1 (1)	125/125 125	131/131 131	124/124 124	0
75 mg	1 (2) <sup>A</sup>	133/145 139.0 (8.5)	170/179 174.5 (6.4)	147/147 147 (0)	1 (2)
100 mg	1 (3) <sup>A</sup>	122/140 132.0 (9.2)	179/193 185.0 (7.2)	133/147 140.7 (7.1)	1 (3)
125 mg	2 (6)	124/171 137.2 (18.8)	144/177 160.0 (14.0)	126/158 134.3 (13.9)	2 (3)

<sup>A</sup> The same subject received these doses of MDMA in different stages of the study.

Candidates with hypertension are excluded from participation in all but one of sponsor-supported studies to limit cardiovascular risk during treatments. In MP-8, four subjects with hypertension controlled by medications were permitted to enroll after completion of carotid ultrasound and nuclear exercise test (per protocol) in addition to usual medical screening for the study. Results are depicted above. One subject dropped out after receiving a single experimental session with 30 mg MDMA and did not experience SBP above cut-off. SBP above cut-off was detected in 75% (3 of 4) of subjects and 67% (8 of 12) of experimental sessions where MDMA was administered to this sub-group. The prevalence of these elevations appears higher in this sub-group than the overall sample, although the prevalence could decrease in a larger group. Pre-drug SBP was typically higher in this sub-group, and peak SBP of these subjects was typically at the upper end of the range of the overall sample. Final SBP readings remained 11 to 14 mmHg higher on average than pre-drug SBP readings in the subject who received 75 mg of MDMA in two blinded experimental sessions and 100 mg in three open-label crossover experimental sessions. However, two subjects receiving 125 mg MDMA had final readings that returned to pre-drug values, suggesting this could be an individual case with a medical history of both hypertension and hyperlipidemia. None of the subjects with controlled hypertension experienced AEs of the cardiovascular system.

**Table 6: Pre-drug, Peak, and Final Diastolic Blood Pressure During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations**

Dose	Subjects (Observations)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with DBP Above Cut-off (Observations)
0 mg	14 (27)	56.5/94 74.6 (9.1)	65/103 84.5 (10.2)	48/100 70.85 (10.9)	0
25 mg	8 (18)	59/84 73.9 (6.2)	76/92 83.2 (5.0)	63/81 72.33 (5.3)	0
30 mg	7 (15)	60/87 74.3 (8.4)	75/99 85.5 (7.5)	68/91 76.7 (6.3)	0
40 mg	7 (12)	69/95 82.7 (8.3)	72/135 90.2 (16.7)	68/96 80.3 (9.4)	1 (1)
75 mg	13 (20)	56/95 75.1 (10.1)	73/118 88.6 (11.2)	59/100 76.1 (10.2)	2 (3)
100 mg	25 (42)	52/93 74.1 (10.6)	62/125 86.9 (15.2)	58/99 74.6 (10.0)	2 (4)
125 mg	95 (232)	54/120 79.1 (9.9)	69/126 92.5 (9.6)	53/104 78.2 (9.8)	6 (8)
150 mg	3 (4)	60/90 78.8 (14.0)	78/108 95.3 (12.6)	67/96 82.0 (12.2)	0

DBP exceeded cut-off in only 5% (16 of 343) of experimental sessions and in 7% (11 of 157) of subjects at any MDMA dose. Maximum duration above DBP cut-off was 5 hours in MP-2 subject 112, with a peak of 114, where 125 mg MDMA was administered as the initial dose. This subject had a high pre-drug DBP reading of 96, and also experienced the highest SBP in sponsor-supported studies to date, as described above. In contrast, 14 subjects participating in 27 experimental sessions with placebo did not experience any elevations in blood pressure above cut-off. In experimental sessions with 25 mg to 30 mg MDMA, elevations in blood pressure above cut-off were not observed either, supporting a dose-dependent effect of MDMA on blood pressure. In all cases, final values returned to pre-drug levels with no clinical intervention required. No clinically significant AEs were reported based on elevations in blood pressure.

**Table 7: Pre-drug, Peak, and Final Diastolic Blood Pressure During Experimental Sessions in Controlled Hypertension Subjects in MAPS-Sponsored PTSD Study MP-8**

Dose	Subjects (Observations)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with DBP Above Cut-off (Observations)
30 mg	1 (1)	85/85 85	86/86 86	77/77 77	0
75 mg	1 (2) <sup>A</sup>	89/95 92 (4.2)	113/118 115.5 (1.8)	91/100 95.5 (6.4)	1 (2)
100 mg	1 (3) <sup>A</sup>	77/91 83.7 (7.0)	121/125 123.0 (2.0)	82/99 90.7 (8.5)	1 (3)
125 mg	2 (6)	82/101 87.8 (8.4)	91/110 98.5 (7.5)	84/93 86.0 (7.5)	0 (0)

<sup>A</sup> The same subject received these doses of MDMA in different stages of the study.

DBP above cut-off was detected in one of four subjects (25%) and five of 12 (41%) of experimental sessions where MDMA was administered at any dose to subjects with controlled

hypertension. All five cases were in the same subject, who received both 75 mg and 100 mg MDMA and is described above. Of all observations of DBP above cut-off across studies and populations, 31% (5 of 16) of experimental sessions were attributed to this subject, suggesting that pre-existing risk factors are associated with elevations in blood pressure. However, this subject did not experience any AEs of the cardiovascular system and DBP resolved back to baseline at final reading in all cases.

In all sponsor-supported studies to date, heart rate readings were taken at baseline, with study-specific differences in data collection times post-drug. Peak values during each experimental session are ascertainable for all studies. The final or endpoint value was recorded as the final value, either at a relatively set time (MP-8, MP-12, MP1-E2) or as the final value available, with time point varying (MP-1). MP-1 and MP-2 reported two pre-drug values (15 minutes and 5 minutes before dosing) and these were averaged, whereas all other studies reported single time point pre-drug. Average post-drug values serve as the final value for MP-2. If heart rate rose above 110 bpm, each duration above the pre-determined cut-off was collected in MP-8, MP-12, MP-9, MP-4, and MP1-E2. Duration of pulse above cut-off was not collected in MP-2. Clinical signs and symptoms were monitored and more frequent readings were collected in cases where readings were above cut-off.

**Table 8: Pre-drug, Peak, and Final Heart Rate During Experimental Sessions with Placebo or any MDMA Dose in MAPS-Sponsored Studies Across Populations**

Dose	Subjects (Observations)	Pre-drug Min/Max Mean (SD)	Peak Min/Max Mean (SD)	Final Min/Max Mean (SD)	Subjects with HR Above Cut-off (Observations)
0 mg	14 (27)	45/111 69.9 (16.3)	54/108 81.2 (14.0)	45/92 70.7 (11.8)	0
25 mg	8 (18)	45/94 69.9 (13.7)	50/124 84.1 (19.8)	51/90 71.7 (12.3)	0
30 mg	7 (15)	45/91 67.1 (14.6)	54/102 81.1 (16.0)	50/89 72.7 (13.0)	0
40 mg	7 (12)	66/110 80.8 (14.3)	69/126 90.7 (15.6)	56/120 83.4 (18.7)	1(1)
75 mg	13 (20)	54/85 72.2 (8.8)	58/123 93.2 (16.9)	57/102 80.9 (13.2)	2 (4)
100 mg	25 (42)	42/114 68.5 (13.5)	63/139 96.6 (17.5)	55/103 78.6 (11.7)	6 (10)
125 mg <sup>A</sup>	95 (232)	36/122 74.9 (13.9)	63/160 104.7 (18.07)	47/135 85.0 (15.1)	51 (90)
150 mg	3 (4)	69/96 79.3 (11.7)	83/125 105.8 (17.3)	74/112 94.5 (15.8)	1 (1)

<sup>A</sup> A single value was not recorded for final readings in subjects receiving 125 mg.

Heart rate elevation above the pre-determined cut-off was detected in 31% (106 of 343) experimental sessions at any MDMA dose, and in 39% (61 of 157) of subjects receiving MDMA. Maximum peak pulse was 160 bpm reported in a subject who received 125 mg MDMA, with pulse remaining above cut-off for 60 minutes. At final reading 3.75 hours later, pulse had returned to below cut-off levels of 93 bpm. The maximum duration above cut-off was 9.5 hours in MP-1 subject 218, where 125 mg MDMA was administered as the initial dose. This subject experienced a peak pulse of 121, which dropped at final reading to 119. Subject 218 had no cardiovascular risk factors in medical history. In cases where blood pressure or heart rate was above cut-off, vitals were monitored more frequently. No subjects receiving MDMA in sponsor-

supported clinical trials have required any clinical interventions for elevated blood pressure or pulse, as all values returned to normal as the effects of MDMA diminished.

The values presented above suggest a dose-dependent action on SBP and heart rate, which is supported in the literature in healthy controls [7, 9, 12, 589]. Peak body temperature and values above cut-off do not appear to be strongly related to MDMA dose, with values above cut-off occurring at every dose, including inactive placebo. While peak DBP is higher after doses of 100 mg or greater, very few reports of DBP elevated above cut-off occurred during MDMA administration, suggesting that this is a less common response than elevated SBP or pulse.

On average, cardiovascular vital signs returned to baseline or near-baseline values by final reading, which is the case across all doses of MDMA. Blood pressure and pulse readings were used to assess AEs described in Section 5.3.9, but they were not the source of the event. There are far fewer observations of elevated DBP than SBP. None of the subjects have required medical intervention after elevations above cut-off, and the elevations were self-limiting and none were clinically significant.

Vital signs for subjects in the study of social anxiety in people on the autism spectrum appear to be similar to those made in people with PTSD receiving equivalent doses of MDMA. Only one measurement rose above pre-determined cut-off values in this sample (pulse above 110, for approximately 1 hour). Comparatively small sample size and use of somewhat lower doses may explain this difference. Differences in age may be involved, with the average age of MAA-1 subjects examined in the IB being 30.65, while mean age in PTSD studies is in the early to mid-40s [43, 590]. No subjects in this study have required any medical interventions.

### 5.3.5 Osmoregulatory Effects

The neuroendocrine hormone copeptin, described in Section 5.3.4 Cardiovascular Effects as correlating with AVP in blood, was detected in women acutely after 125 mg MDMA administration [561], and this finding was reproduced in another study reporting that 47.5 mg MDMA caused an acute rise in AVP and a small decrease in plasma sodium, at a time of day when it would not be expected to change, in an all-male sample. [251]. The sponsor-supported study MAA-1 includes AVP assessments in peripheral plasma samples before, during, and after MDMA-assisted psychotherapy. This study is ongoing and results are pending analysis.

### 5.3.6 Hepatic Effects

The first two sponsor-supported Phase 2 studies (MP-1, MP-2) assessed liver function after completion of two or three blinded experimental sessions. Values that differ from established, age-appropriate norms were evaluated for clinical significance. Laboratory assessments of liver function were not conducted after experimental sessions in subsequent sponsor-supported studies and no AEs related to liver function have been reported in these studies.

**Table 9: List of All Clinically Significant Changes in Laboratory Values in Two Subjects from MP-2**

Laboratory Value	Abnormal Test Value	Value at Baseline	Normal Value/Range	Condition
Bilirubin	2.8	2.2	<2.5 mg/dL	125 mg
ESR	32	2.4	<10 mm	125 mg

Two subjects in the MP-2 study reported two clinically significant abnormalities. One was an elevation in bilirubin in a subject with a family history of elevated bilirubin (probably Gilbert’s

syndrome), with the elevation occurring after open-label treatment with 125 mg to 150 mg initial dose of MDMA. Bilirubin levels can be indicative of decreased liver function, but the liver enzymes were normal at that time, supporting the interpretation that the bilirubin levels were slightly elevated compared to baseline due to hereditary factors. The other abnormal laboratory value, an elevation in erythrocyte sedimentation rate (ESR), a marker of inflammation, occurred in a subject with a medical history of breast cancer. This value was recorded 3 months after the last administration of MDMA as an AE unrelated to the study drug.

**Table 10: Average ALT Values at Baseline and 2-Month Follow-up After Two Experimental Sessions in Subjects from MP-1**

Timepoint	Placebo	125 mg
Baseline	25.6 (13.4) N=8	22.75 (12.89) N=12 <sup>^</sup>
Primary Endpoint After Two Experimental Sessions	26.4 (13.5) N=8	19.7 (12.7) N=13

<sup>^</sup> ALT value for one subject not recorded at baseline.

No clinically significant changes in liver function occurred in MP-1. Values for laboratory tests were within the normal range in MP-1. An independent t-test of differences between baseline and 2-month follow-up alanine aminotransferase (ALT) in placebo and MDMA subjects in MP-1 detected a trend toward a change that implied improved liver function that failed to reach statistical significance. Phase 1 studies conducted outside of sponsor support involving administration of MDMA to healthy volunteers have not published any results of liver function after MDMA administration. There have been no reported adverse effects on the liver from these studies.

### 5.3.7 Neurobiological Effects

Early investigations in healthy volunteers used PET to detect changes of brain activity after MDMA and found decreased left amygdalar activity and increased frontal activity [28]. PET brain scans 75 minutes after administration of 1.7 mg/kg MDMA found increased regional cerebral blood flow (rCBF) in ventromedial prefrontal, inferior temporal, and cerebellar areas and decreased rCBF in the left amygdala [28]. In a different study, arterial spin labeling has also found decreased cerebral blood flow (CBF) in the right amygdala and hippocampus after MDMA administration [27]. The decreased CBF correlated with drug intensity ratings after 100 mg MDMA. Blood oxygen level dependent (BOLD) MRI scans of resting-state functional connectivity in the same sample detected complementary decreases in medial PFC-hippocampal coupling and increases in right amygdala-hippocampal coupling, although the relationship did not achieve statistical significance [27]. Decreased activity in the amygdala may be indicative of reduced reactions to potential threats [591]. MDMA (100 mg) increased subjective ratings of positive mood in response to positive memories and decreased negative response to negative memories. Attenuated activity in the left anterior temporal area was detected after MDMA during worst memory recall. [36].

During a task that required keeping a visual target cue in mind, visual attention, and response inhibition, brain imaging detected changes in parietal activity after 75 mg MDMA compared with placebo [576]. MDMA increased activity in frontal areas and decreased activity in occipital sites as measured via functional MRI (fMRI) [592]. Reduced resting-state cerebral blood flow in right amygdala and hippocampus after MDMA was associated with greater intensity of self-reported subjective effects [27]. Subjects given MDMA exhibited similar brain activity when reading or encoding a word list, suggesting that they were investing similar effort into both tasks. Ten Ecstasy user subjects receiving a minimum of two doses of 1 to 1.25 mg/kg or 2.25 to 2.5 mg/kg

MDMA exhibited signal decreases in bilateral visual cortex, caudate, superior parietal, and dorsolateral frontal regions 10 to 21 days later, with increased rCBF measured in two subjects at a later time point. However, a comparison between heavy Ecstasy users and non-user controls failed to find differences in baseline rCBF [593], and a report assessing changes before and after initial Ecstasy use found increased rCBF in only one area of the prefrontal cortex [266], suggesting that the changes seen by Chang and colleagues are a transient effect. EEG recorded 2 hours after MDMA administration showed the following changes in EEG activity: overall increase in beta activity, reduction in alpha activity, localized decreases in alpha and delta in frontal areas, and increased frontotemporal beta signal [594]. The authors reported the EEG patterns after MDMA were similar to those seen with serotonergic and noradrenergic drugs, as well as, but to a lesser extent, dopaminergic drugs.

The sponsor is undertaking a small BOLD fMRI pilot study investigating brain activity in people with PTSD before and after MDMA-assisted psychotherapy, as a substudy of a sample of people enrolled in MP-8. Brain activity is recorded while the subject is listening to a neutral and a personalized trauma-related scripts. Preliminary findings are pending analysis.

Monoamine neurotransmitters are known to modulate sleep architecture and alertness. In a trial with 2 mg/kg MDMA given 6 hours prior to preparing for sleep, MDMA was found to increase Stage 1 sleep and produce fewer periods of REM sleep without increasing daytime sleepiness [281]. Sample size of seven in this study suggests that findings should be accepted with caution. PTSD patients suffer from poor sleep quality. Disturbed REM or non-REM sleep is a contributing factor to maladaptive stress and trauma responses and chronic sleep disruption associated with nightmares caused by PTSD may be an indicator of efficacy of PTSD treatments. The sponsor is collecting secondary outcomes in PTSD studies with the Pittsburgh Sleep Quality Index. Results are pending analysis from ongoing studies.

### 5.3.8 Neuropsychological Effects

MDMA alters mood, perception, and cognition in healthy volunteers, with effects on emotion and social behavior. At doses of at least 1 mg/kg (approximately 70 mg) and higher, active doses of MDMA alter mood and cognition, and produce slight alterations in perception [10, 529]. Acute subjective effects peak 90 to 120 minutes after oral administration and return to pre-drug levels 3 to 6 hours later [13, 595, 596]. Sub-acute effects assessed in controlled and naturalistic studies may occur 1 to 3 days after drug administration, but are no longer apparent seven to 14 days later [12, 324, 597]. Most of the therapeutic effects of MDMA are thought to result from changes in affect, cognition, and social interaction.

At least four research teams published relevant findings in studies of healthy volunteers during 2013 and 2014, examining the effects of MDMA on social cognition with several experimental paradigms assessing brain activity during episodic memory recall and assessing contributions of oxytocin and cortisol to the acute effects of MDMA. Findings include reduced reactivity to simulated social exclusion, reduced negative emotional response to self-selected “worst” memories, increased use of language related to interpersonal closeness, increased emotional empathy and increases in perceived partner empathy. One study reported greater social language after MDMA than with the psychostimulant methamphetamine [37], and another reported greater emotional empathy after MDMA and another psychostimulant, methylphenidate [34]. Taken together, this research lends greater support to the view that MDMA possesses unique psychological effects, distinct from psychostimulants that can be beneficial when combined with psychotherapy. As an entactogen, MDMA can promote increased trust, greater ability to face and cope with emotionally distressing memories, thoughts or feelings and greater emotional empathy toward the self as well as others.



When combined with psychotherapy, MDMA permits people to confront and consider emotionally intense memories, thoughts, or feelings, and perhaps through changes in mood and perception, increase empathy and compassion for others and oneself [41, 62, 527]. In a sub-study of MP-8, the Self Compassion Scale [598] was administered before and 2 months after MDMA-assisted psychotherapy. Preliminary results in this small sub-study (N=7) are trending upward; subjects were low in self-compassion with mean total score of  $2.4 \pm 0.63$  prior to the study and experienced an increase to moderate self-compassion with mean total score of  $2.8 \pm 0.84$ . In this assessment, self-kindness and a sense of common humanity increased, while self-judgment and feelings of isolation decreased on average within-subjects.

A Phase 1 study of the effects of MDMA-assisted psychotherapy on mood and social cognition in healthy volunteers who completed training in performing manualized MDMA-assisted psychotherapy is underway. Findings will include effect on mood and interpersonal closeness. The ongoing MAA-1 study in autistic adults is measuring symptoms of social anxiety, with secondary measures of emotion identification in the self and others, emotion regulation, alexithymia, and empathy. In this study, biomarkers associated with social behavior, including oxytocin, AVP, and cortisol, will also be assessed before, during, and after MDMA-assisted therapy. Taken together, findings from ongoing studies will assist the sponsor in evaluating how neuropsychological effects contribute to clinical development of MDMA-assisted psychotherapy.

### 5.3.8.1 Cognitive Function

MDMA does not affect responses on tasks requiring attention and response to visual stimuli or visually presented words [13, 28], but has been shown to interfere with performance on digit-symbol substitution, a measure of attention, psychomotor speed and visual memory [8]. A dose of 75 mg improved visual tracking speed, but impaired estimating the position of a blocked (occluded) object in a study of acute effects on skills used for driving cars [595]. A series of studies conducted in the Netherlands examined the effects of MDMA on skills needed for automobile driving reported transient and selective changes in verbal and visual attention, and memory after 75 or 100 mg MDMA [599-602]. MDMA caused difficulty learning or remembering lists of words and difficulty recalling object position within an array of objects. MDMA did not cause impairment in spotting scene changes and reduced weaving in a driving simulation. MDMA was associated with an excessively cautious response to the actions of another car in an assessment of actual driving [603]. While these studies have added to the literature of MDMA's cognitive effects, people in sponsor-supported studies are advised to never operate a vehicle while under the influence of MDMA or any other psychoactive substance.

MDMA acutely improved performance on one measure of impulsivity while failing to affect performance on other impulsivity measures [600]. The causes of these changes are unclear but may relate to changes in attention, salience of visual objects, and altered time perception. Changes in visuospatial recall and driving skills are likely associated with serotonin release or indirect action on serotonin receptors, as the noradrenergic and dopaminergic drug methylphenidate (Ritalin) did not produce similar changes [599, 602, 603]. A study on performance monitoring compared the effects of ethanol, MDMA, and both substances combined, found that MDMA had no effect on performance monitoring and no interaction when ethanol and MDMA are administered concurrently [604]. Administration of a 5HT<sub>2A</sub> receptor antagonist, but not a 5HT<sub>1A</sub> antagonist, reduced impaired performance on a word learning and recall task after MDMA, suggesting that interference is due in part to direct or indirect activation of these receptors [233]. Changes in cognitive function and psychomotor skills occurred during peak drug effects, but were not detectable 24 hours later.

Acute effects on cognitive function are not assessed in sponsor-supported studies. In three MAPS-sponsored studies, MP-1, MP-4, and MP-12, long-term effects on cognitive function was assessed by administering the Repeatable Battery for Assessment of Neuropsychological Status (RBANS), a relatively brief measure that assesses memory, attention and processing speed, visual-spatial and constructional abilities, and expressive language [605]; and the Paced Auditory Serial Addition Task (PASAT), a measure of auditory processing speed and mental flexibility [606, 607]. These instruments were given prior to and 1 to 2 months after psychotherapy assisted with either MDMA or comparator or placebo.

In MP-1, no significant differences in cognitive function were detected at the 2-month follow-up between subjects who received two sessions with 125 mg of MDMA compared to subjects who received placebo, as measured by RBANS and PASAT [41]. These findings suggest that MDMA did not impair cognitive function in this sample or that the effect was too small to attain statistical significance in this small pilot study. Two ongoing studies (MP-12 and MP-4) include these measures to assess reproducibility of this finding. Since both MP-4 and MP-12 were ongoing as of the data cut-off, available data pooled across studies are presented below by dose.

**Table 11: Neurocognitive Function - RBANS Mean Total Scores at Baseline, Primary Endpoint, End of Stage 1, and End of Stage 2 for MP-1, MP-4, and MP-12 as of 01 October 2015**

Dose	Baseline Mean (SD) N	Primary Endpoint Mean (SD) N	End of Stage 1 Mean (SD) N	End of Stage 2 Mean (SD) N
0 mg	100.9 (15.38) N=10	106.9 (15.15) N=10	---	119.0 N=1
40 mg	94.7 (5.20) N=6	102.0 (10.58) N=3	---	101.3 (5.51) N=3
100 mg	95.0 (17.87) N=6	104.9 (15.75) N=7	101.5 (18.97) N=6	---
125 mg	102.9 (15.88) N=27	103.1 (12.70) N=22	99.5 (9.33) N=6	---

On average, RBANS scores trend towards improvement after treatment with placebo and 40 mg to 100 mg initial dose of MDMA, whereas scores stay the same after treatment with 125 mg initial dose of MDMA. The trend towards improvement could be a practice effect from repeated assessments, although stimuli were varied across these, or could possibly be correlated with PTSD symptom reduction. One to three additional treatments with open-label active dose MDMA do not appear to worsen cognitive function based on preliminary End of Stage 1 and End of Stage 2 results. The significance of these pooled findings is yet to be determined.

**Table 12: Neurocognitive Function - PASAT Trial 1 and Trial 2 Mean Raw Total Scores at Baseline, Primary Endpoint, End of Stage 1, and End of Stage 2 for MP-1, MP-4, and MP-12 as of 01 October 2015**

<b>PASAT Trial 1</b>				
<b>Dose</b>	<b>Baseline Mean (SD)</b>	<b>Primary Endpoint Mean (SD)</b>	<b>End of Stage 1 Mean (SD)</b>	<b>End of Stage 2 Mean (SD)</b>
0 mg	42.1 (12.59) N=10	43.7 (12.03) N=10	---	37.0 N=1
40 mg	43.6 (10.36) N=5	53.3 (4.16) N=3	---	52.7 (5.03) N=3
100 mg	44.3 (12.44) N=6	46.7 (9.74) N=7	49.3 (9.09) N=6	---
125 mg	44.1 (11.12) N=27	49.1 (8.48) N=22	53.0 (6.36) N=6	---
<b>PASAT Trial 2</b>				
0 mg	34.2 (11.21) N=10	38.6 (11.66) N=10	---	45.0 N=1
40 mg	34.0 (13.36) N=5	43.0 (10.39) N=3	---	45.7 (8.15) N=3
100 mg	31.2 (12.67) N=6	29.0 (13.37) N=7	38.0 (10.33) N=6	---
125 mg	32.6 (9.62) N=27	35.4 (8.42) N=21	42.0 (11.8) N=6	---

On average, PASAT scores stay about the same after treatment with placebo and 100 mg initial dose of MDMA and trend towards improvement after treatment with 40 and 125 mg initial dose of MDMA. The trend towards improvement could be a practice effect from repeated assessments or could be correlated with PTSD symptom reduction. One to three additional treatments with open-label active dose MDMA do not appear to worsen cognitive function and continued to trend towards improvement on average based on preliminary End of Stage 1 and End of Stage 2 results. Cognitive function tests such as the PASAT are also known to be subject to individual variability, as they require basic proficiency with mathematical skills that are influenced by education level. The significance of these pooled findings is yet to be determined, but it does not appear that MDMA-assisted psychotherapy is negatively impacting cognitive function.

### 5.3.8.2 Perceptual Effects

MDMA causes slight changes in visual or auditory perception, including changes in the brightness or colors, sounds seeming closer or farther away, and simple visual distortions [7, 8, 10, 12]. Subjects also experienced altered time perception, and changes in meaning or significance of perceptions after MDMA [13]. On average, subjects maintained insight of their experience, with little indication that MDMA produces any strong alterations to the sense of self or control over the experience [11, 12]. Three healthy volunteers reported developing minimal to mild unusual beliefs or delusions under the influence of 1.5 mg/kg MDMA. Findings from a study with a small sample (five per group), perceptual alteration may be more pronounced after 2 mg versus 1 mg [596]. These beliefs resolved within a few hours, or by the next day at the latest. These subjects were aware that these beliefs were unusual [12]. Women reported experiencing all subjective effects of MDMA more intensely compared to men, but especially those related to perceptual changes [10]. The perceptual effects of MDMA appear to be the result of direct or indirect action on 5HT<sub>2A</sub> receptors, as co-administration of the 5HT<sub>2A</sub> antagonist ketanserin reduced reported perceptual alterations, as well as eliminated slight elevations in body temperature after 1.5 mg/kg MDMA [559], while co-administration with the 5HT<sub>1A</sub> antagonist

pindolol did not affect perceptual alteration [231]. The effects of MDMA upon perception have not been studied within sponsor-supported studies.

### 5.3.8.3 Social Effects

In controlled laboratory settings, an established measure of accurate facial expression reading found that MDMA improved detection of expressions of positive mood and reduced accuracy in detecting expressions of negative mood [30]. Despite initial findings in naturalistic studies suggesting that Ecstasy increased accuracy of assessing some emotional expressions, particularly fearful ones [608], an fMRI study found that 0.75 and 1.5 mg/kg MDMA reduced signaling in the amygdala in response to angry faces when compared with placebo without changing the response to faces showing fear [26]. These researchers also detected increased activity in the ventral striatum in response to happy faces. Taken together, these findings suggest that MDMA changes the way emotional facial expressions are processed or the response to them. Complementing these findings are results demonstrating that MDMA enhanced the accuracy of recognizing facial expressions of positive mood and impaired mind reading for facial expressions of negative mood, but had no effect on mind reading for neutral faces [30]. Enhanced mind reading of positive emotions may facilitate therapeutic relationships in MDMA-assisted psychotherapeutic settings. In addition, and contrary to the finding in the early naturalistic study described above, there is some evidence showing that MDMA produces selective difficulty in recognizing faces expressing fear [588]. Further investigation corroborates this finding, showing that MDMA reduced recognition accuracy of fear significantly more in women than in men, and reduced recognition accuracy of sadness in women, but not in men. The same study found MDMA-induced increases in both implicit and explicit emotional empathy in men, but not in women [19].

Findings in placebo-controlled trials suggest that MDMA enhances positive response to positive social stimuli. Wardle and colleagues observe this effect simultaneously with a decrease in positive response to positive stimuli with no social content, which suggests that the contrast in valuation of social and non-social emotional stimuli contributes to MDMA's prosocial effects [38]. MDMA also reduces the impact of rejection on mood and self-esteem [609], which manifests more strikingly at lower doses of MDMA than reduction in perceived social rejection, suggesting complex social and behavioral effects from MDMA. Moreover, results from Kirkpatrick and colleagues show a behavioral preference for social activities over non-social ones, with subjects reporting increased desire for only the social activity after 1.5 mg/kg MDMA [610].

In a study by Bedi and colleagues, MDMA induced changes in semantic speech content with natural language learning software. Through natural language processing (NLP), researchers found speech patterns after MDMA were distinct from those produced after methamphetamine and placebo [37]. Proximity of speech to the concepts of *friend*, *support*, *intimacy*, *rapport*, and *empathy* was increased in the MDMA drug condition, which may bear some significance for the use of MDMA in therapy. MDMA did not affect the overall structure of subjects' speech. These findings were confirmed in an additional sample through a standardized dictionary method and machine learning, indicating that MDMA increased the use of social words, as well as words connoting positive and negative emotions [240]. There is some evidence that the increases in affiliative and prosocial feelings are separable from romantic or sexual feelings. Men and women did not seek to prolong viewing of images with explicit sexual content after MDMA, and they did not impute increased romantic feelings to images of heterosexual couples [611].

While the hormone oxytocin is implicated in social interactions and bonding, evidence indicates that oxytocin alone does not explain MDMA's prosocial effects. One investigation found a positive correlation in subjective effects ratings between intranasal oxytocin and oral MDMA, but

only at the lower of the two oxytocin doses tested [63]. Using pindolol to block 5-HT<sub>1A</sub> receptor mediation of oxytocin's effects, Kuypers and colleagues determined that MDMA increased emotional empathy while oxytocin did not produce similar effects on measures of empathy and social interaction [569]. Studies examining the prosocial effects of MDMA, in relation to oxytocin, should be considered in the context of previous findings that showed no discernable subjective effects were found for intranasal oxytocin [612]. A single nucleotide polymorphism in the oxytocin receptor gene was found to predict subjective responses to MDMA, suggesting that this question remains worthy of further study [613]. Two studies have found that MDMA increased AVP [251, 561]. Neither study reported analysis or findings concerning any relationship between AVP levels and the subjective, emotional or social effects of MDMA.

Studies in healthy controls comparing doses between 0.75 and 1 mg/kg and 1.5 to 2 mg/kg suggest that the higher dose produces greater prosocial effects than the lower dose, while the lower dose may increase self-reported loneliness and use of empathy-related language [35, 39, 596, 609]. However, higher doses also produce a greater degree of stimulation and anxiety. It is notable that the first study investigating the impact of variation in an oxytocin receptor gene reported that those with one variation did not exhibit an increase in sociability after 1.5 mg/kg without a statistically significant difference in response at 0.75 mg/kg [613].

#### 5.3.8.4 Emotional Effects

MDMA increases positive mood and anxiety [8, 10-12] on measures of alteration in consciousness and subjective effects. There is evidence that increases in positive mood and anxiety increase with dose [8, 12, 35, 614]. MDMA users report feeling more talkative and friendly after receiving MDMA. Self-reported interpersonal closeness was noted during a study in healthy volunteers [13]. Subsequent research confirmed the occurrence of increased interpersonal closeness after MDMA [29, 30, 35, 91, 558]. Researchers using two items within an instrument designed to assess drug effects and a visual analog scale rating closeness to others failed to detect increased feelings of empathy after 1.5 mg/kg MDMA [12], possibly due to the low sensitivity of these measures. In another investigation, the SSRI paroxetine was pre-administered to healthy volunteers before administering MDMA. The researchers found that MDMA increased feelings of being social and closeness to others, and paroxetine reduced these effects, indicating a significant role of the serotonergic system for the prosocial effects of MDMA [91]. People have reported feeling anxious or experiencing negative derealization while under the influence of MDMA, including increased anxiety related to loss of control and experiences of racing or blocked thoughts [8, 10, 13].

People receiving active doses of MDMA experience euphoria, positive mood, vigor, and positively experienced derealization, consonant with early retrospective reports, but also report experiencing anxiety, tension, and dysphoria, as well as concern over losing control over the self [8, 10-12]. More surprisingly, subjects report increased positive mood even after a dose of 25 mg [614]. It is uncertain whether the increases in positive and negative mood occur simultaneously or at different times throughout the duration of MDMA effects; evidence from two different teams suggests that peaks in negative mood may precede peaks in positive mood [11, 563]. MDMA may have a greater impact on mood in women than in men. Women report greater elevation in negative mood despite reaching plasma concentrations of MDMA and metabolites similar to those of men [552]. A second dose of MDMA 2 hours after the first does not increase subjective effects beyond that of an initial dose, interpreted by Peiro and colleagues as indications of tolerance to these effects [328]. When two 100 mg doses are given 4 hours apart, most subjective effects are comparable to those after a single dose, despite there being double the amount of plasma MDMA [546]. It is notable that the second dose in this study was identical to the first



dose, in contrast to sponsor-supported studies, wherein the second dose is half the size of the initial dose.

### 5.3.8.5 Suicidal Ideation, Behavior, and Depression

There is high incidence of positive suicidal ideation and behavior in populations of people with PTSD, especially those suffering from chronic, treatment-resistant PTSD [615, 616]. The FDA has responded to concerns over the occurrence of treatment emergent suicidal ideation or behavior by requiring clinical trials of psychiatric drugs to measure suicidality via the C-SSRS, a clinician-administered guided interview [617]. A score of 4 or 5 on the suicidal ideation category is considered serious, as well as a score of 1 or greater on the behavior category, and individuals with serious ideation or behavior are closely followed until levels return to normal or additional interventions are recommended. In order to determine if suicidal ideation and behavior worsens or improves after treatment in ongoing MAPS-sponsored trials (MP-4, MP-8, MP-9, MP-12, MAA-1, MDA-1, and MT-1), the C-SSRS is given repeatedly throughout a study, including lifetime incidence, baseline, before/during/after drug administration, endpoints when other measures are administered, and follow-up visits. Findings concerning suicidal ideation or behavior have not been formally measured in the first two sponsor-supported studies or reported in studies of healthy volunteers. Due to the nature of the therapeutic method, wherein a person may re-experience emotions associated with the traumatic event in order to reprocess the memory in a new, less detrimental way, thoughts of ending one's life may surface during this process. However, evidence from clinical studies indicates that these thoughts are most often transient, returning to normal, or even improve during the acute period following MDMA treatment. C-SSRS scores have also escalated during the preparatory sessions (before any drug administration), which is thought to be either a result of discussing traumatic experiences, or subjects tapering off long-prescribed medications, such as SSRIs and benzodiazepines, which have been documented elsewhere to induce suicidal ideation or behavior during withdrawal [618-620]. During both non-drug and MDMA-assisted psychotherapy sessions, subjects are asked to think about and discuss their experiences, thoughts, and emotions related to their condition. They may experience intense emotional responses to recalling and speaking about this material. As MDMA is only administered in combination with psychotherapy, the distress associated with psychotherapy is unavoidable, and is considered a necessary part of the therapeutic process that requires proper facilitation and support from therapists.

In Tables 13 through 17 below, suicidal ideation and behavior are summarized for subjects in MP-4, MP-8, MP-9, MP-12, MDA-1, and MAA-1 according to suggestions made in the C-SSRS Scoring and Data Analysis Guide [621]. A positive response for suicidal ideation is counted when a subject responds "yes" to any one of the five suicidal ideation questions (Categories 1 to 5) on the C-SSRS (i.e. a score >0 for suicidal ideation score). Serious suicidal ideation is a suicidal ideation score of 4 or 5. A positive response for suicidal behavior occurs when a subject responds "yes" to any one of the five suicidal behavior questions (Categories 6 to 10) on the C-SSRS (i.e. a score >0 for suicidal behavior score). Lifetime scores account for all suicidal ideation and behavior prior to enrollment according to subject recall and medical records. Pre-drug exposure represents measures collected on the Since Last Visit C-SSRS after enrollment during preparatory sessions and before first drug administration in experimental session 1 upon completion of tapering off psychiatric medications. Frequencies are event-based, calculated based on percentage of observations in which subjects would have the opportunity to report, as the C-SSRS is collected multiple times with each exposure to MDMA.



**Table 13: Summary of Baseline Positive and Serious Responses on C-SSRS for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015**

Condition		Lifetime <sup>A</sup> N (%)	Pre-drug Exposure <sup>B</sup> N (%)
<b>PTSD</b>			
Blinded Placebo (0 mg)	PI	3 (75%)	3 (38%)
	SI	2 (50%)	0 (0)
	PB	2 (50%)	0 (0)
	O	4	8
	N	4	4
Blinded Comparator Doses (25-40 mg)	PI	11 (79%)	5 (15%)
	SI	3 (21%)	0 (0)
	PB	6 (43%)	0 (0)
	O	14	33
	N	14	13
Blinded Active Doses (75-125 mg)	PI	41 (93%)	28 (29%)
	SI	19 (43%)	0 (0)
	PB	19 (43%)	2 (2%)
	O	44	97
	N	44	44
<b>Social Anxiety in Autistic Adults</b>			
Blinded Placebo (0 mg)	PI	2 (100%)	0 (0)
	SI	0 (0)	0 (0)
	PB	1 (50%)	0 (0)
	O	2	3
	N	2	2
Blinded Active Doses (75-125 mg)	PI	2 (100%)	0 (0)
	SI	1 (50%)	0 (0)
	PB	1 (50%)	0 (0)
	O	2	6
	N	2	3
<b>Anxiety Associated with a Life-threatening Illness</b>			
Blinded Placebo (0 mg)	PI	1 (50%)	3 (38%)
	SI	1 (50%)	0 (0)
	PB	0 (0)	1 (13%)
	O	2	8
	N	2	2
Blinded Active Dose (125 mg)	PI	1 (50%)	0 (0)
	SI	0 (0)	0 (0)
	PB	1 (50%)	0 (0)
	O	2	8
	N	2	2

PI=Positive Ideation, SI=Serious Ideation, PB=Positive Behavior, O=Observations, N=Number of Subjects

<sup>A</sup> Lifetime accounts for all suicidal ideation and behavior prior to study Visit 1, according to participant recall and medical records

<sup>B</sup> Pre-drug exposure represents measures taken during Preparatory Sessions and before drug administration in Experimental Session 1

Based on lifetime results, most subjects across populations and dose groups had a history of suicidal ideation. In the PTSD sample, 39% of subjects had a history of serious ideation and 43% had positive behavior, which is consistent with the literature. Although samples were small, non-PTSD samples also have evidence of suicidal ideation and behavior, although prevalence may change as these studies enroll more subjects. Two PTSD subjects randomized to active dose and one autistic subject randomized to placebo exhibited suicidal behavior prior to any MDMA administration.

**Table 14: C-SSRS Positive and Serious Responses During Experimental Sessions and 1-Day Post-Drug for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015**

Condition		Session 1 N (%)			Session 2 N (%)			Session 3 N (%)		
		Pre-drug <sup>A</sup>	During-drug <sup>B</sup>	Integration Day 1	Pre-drug <sup>A</sup>	During-drug <sup>B</sup>	Integration Day 1	Pre-drug <sup>A</sup>	During-drug <sup>B</sup>	Integration Day 1
<b>PTSD</b>										
Blinded	PI	1 (25%)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	---	---	---
Placebo	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
(0 mg)	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	N	4	4	4	4	4	4			
Blinded	PI	1 (7%)	1 (8%)	0 (0)	0 (0)	0 (0)	1 (8%)	1 (33%)	0 (0)	0 (0)
Comparator	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Doses	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
(25-40 mg)	N	14	13	13	12	11	12	3	2	2
Blinded	PI	9 (21%)	3 (7%)	5 (12%)	9 (21%)	7 (17%)	2 (5%)	4 (11%)	6 (17%)	3 (9%)
Active Doses	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
(75-125 mg)	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	44	43	43	42	42	42	35	35	35
Open-label	PI	0 (0)	1 (6%)	2 (11%)	1 (5%)	0 (0)	0 (0)	1 (6%)	0 (0)	0 (0)
Stage 2	SI	0 (0)	1 (6%)	1 (5%)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Active Dose	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
(100-125 mg)	N	19	18	19	19	17	19	18	17	17

PI=Positive Ideation, SI=Serious Ideation, PB=Positive Behavior, N=Number of Subjects

<sup>A</sup> Pre-drug measurement taken day of experimental session prior to drug administration.

<sup>B</sup> During-drug observation measured at experimental session endpoint, approximately 6 hours after drug administration.

In a PTSD sample with prevalent lifetime history of suicidal ideation, subjects randomized to either dose group reported pre-drug suicidal ideation in blinded experimental sessions. More active dose subjects reported pre-drug positive ideation, likely due to oversampling. During blinded session 1, numbers of comparator and active dose subjects reporting positive ideation were equivalent, with only active dose subjects reporting positive ideation the next day and none serious. As active dose subjects went deeper in the therapeutic process, reports of positive ideation 6 hours post-drug increased to 17% in the blinded session 2, with no reports from comparator subjects. No subjects reported positive suicidal behavior 6 hours post-drug as they were under continuous clinical observation during treatment and for 24 hours after. 5% of active dose subjects and 8% of comparator dose subjects experienced positive ideation the next day, with none serious. Active dose session 3 was similar to the second. Interestingly, open-label experimental sessions had fewer reports of positive and serious ideation, suggesting a protective effect of receiving comparator dose sessions prior to active dose, which could be attributed to developing the therapeutic alliance.

**Table 15: C-SSRS Positive and Serious Responses During Experimental Sessions and 1-Day Post-Drug for Studies MAA-1 and MDA-1 as of 01 October 2015**

Condition		Session 1 N (%)			Session 2 N (%)			Session 3 N (%)		
		Pre- drug <sup>A</sup>	During- drug <sup>B</sup>	Integration Day 1	Pre- drug <sup>A</sup>	During- drug <sup>B</sup>	Integration Day 1	Pre- drug <sup>A</sup>	During- drug <sup>B</sup>	Integration Day 1
<b>Social Anxiety in Autistic Adults</b>										
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	---	---	---
Placebo (0 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	N	2	2	2	2	2	2			
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	---	---	---
Active Doses (75-125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	N	3	3	3	3	3	3			
Open-label Stage 2	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	---	---	---
Active Dose (100-125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	N	2	2	2	2	2	1			
<b>Anxiety Associated with a Life-threatening Illness</b>										
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	---	---	---
Placebo (0 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)			
	N	2	2	2	1	1	1			
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Active Dose (125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	2	2	2	1	2	2	1	1	1

PI=Positive Ideation, SI=Serious Ideation, PB=Positive Behavior, N=Number of Subjects

<sup>A</sup> Pre-drug measurement taken day of experimental session prior to drug administration.

<sup>B</sup> During-drug observation measured at experimental session endpoint, approximately 6 hours after drug administration.

Despite prevalent lifetime history of suicidal ideation with 50% of subjects reporting serious ideation, five autistic subjects in MAA-1 reported no suicidal ideation or behavior before, during, or after experimental sessions regardless of MDMA dose. Although prevalence was about half of the PTSD and autistic subject samples, subjects with anxiety associated with a life-threatening illness also did not report suicidal ideation or behavior before, during, or after experimental sessions. These results may vary as more subjects are treated, but appear encouraging.

**Table 16: C-SSRS Positive Responses During Telephone Contact Following Experimental Sessions for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015**

Condition		Session 1		Session 2		Session 3	
		N (%)		N (%)		N (%)	
		Day 2	Day 7	Day 2	Day 7	Day 2	Day 7
<b>PTSD</b>							
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	---	---
Placebo (0 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)		
	PB	0 (0)	0 (0)	0 (0)	0 (0)		
	N	4	4	4	4		
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Comparator Doses (25-40 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	14	14	12	12	2	2
Blinded	PI	5 (12%)	6 (15%)	8 (20%)	6 (15%)	4 (12%)	4 (12%)
Active Doses (75-125 mg)	SI	0 (0)	0 (0)	0 (0)	1 (3%)	0 (0)	1 (3%)
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	42	41	42	39	34	33
Open-label Stage 2	PI	3 (16%)	0 (0)	1 (5%)	0 (0)	0 (0)	0 (0)
Active Doses (100-125 mg)	SI	2 (11%)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	19	18	19	19	17	18
<b>Social Anxiety in Autistic Adults</b>							
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	---	---
Placebo (0 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)		
	PB	0 (0)	0 (0)	0 (0)	0 (0)		
	N	2	2	2	2		
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	---	---
Active Doses (75-125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)		
	PB	0 (0)	0 (0)	0 (0)	0 (0)		
	N	3	3	3	3		
Open-label Stage 2	PI	0 (0)	0 (0)	0 (0)	0 (0)	---	---
Active Dose (100-125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)		
	PB	0 (0)	0 (0)	0 (0)	0 (0)		
	N	2	2	2	2		
<b>Anxiety Associated with a Life-threatening Illness</b>							
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	---	---
Placebo (0 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)		
	PB	0 (0)	0 (0)	0 (0)	0 (0)		
	N	2	2	1	1		
Blinded	PI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
Active Dose (125 mg)	SI	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	PB	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)	0 (0)
	N	2	2	2	2	1	1

PI=Positive Ideation, SI=Serious Ideation, PB=Positive Behavior, N=Number of Subjects

Reports of positive ideation during treatment continued during the week after experimental sessions in 12% to 20% of subjects randomized to active dose MDMA. Lack of reports in comparator dose subjects suggests a dose-dependent effect. Prevalence increased after the second experimental session as seen during experimental sessions, likely due to enhancement of the therapeutic process with each exposure bringing up disturbing traumatic thoughts. As MDMA is only administered in the context of psychotherapy, and PTSD subjects have a lifetime history of suicidal ideation, these effects are expected. In contrast, autistic adults and those with anxiety associated with a life-threatening illness reported no suicidal ideation or behavior.

**Table 17: C-SSRS Positive Responses at Endpoints After Treatment for Studies MP-4, MP-8, MP-9, MP-12, MAA-1, and MDA-1 as of 01 October 2015**

Condition		Primary/ Secondary Endpoint N (%)	End of Stage 1/ End of Stage 2 N (%)	Long-term Follow-up N (%)
<b>PTSD</b>				
Blinded Placebo (0 mg)	PI	0 (0)	---	---
	SI	0 (0)		
	PB	0 (0)		
	N	2		
Blinded Comparator Doses (25-40 mg)	PI	1 (8%)	0 (0)	1 (14%)
	SI	0 (0)	0 (0)	0 (0)
	PB	0 (0)	0 (0)	0 (0)
	N	12	2	7
Blinded Active Doses (75-125 mg)	PI	13 (36%)	7 (23%)	5 (31%)
	SI	1 (3%)	0 (0)	1 (6%)
	PB	0 (0)	0 (0)	0 (0)
	N	36	30	16
Open-label Stage 2 Active Doses (100-125 mg)	PI	0 (0)	1 (8%)	---
	SI	0 (0)	0 (0)	
	PB	0 (0)	0 (0)	
	N	7	13	
<b>Social Anxiety in Autistic Adults</b>				
Blinded Placebo (0 mg)	PI	---	0 (0)	0 (0)
	SI		0 (0)	0 (0)
	PB		0 (0)	0 (0)
	N		2	2
Blinded Active Doses (75-125 mg)	PI	---	1 (17%)	0 (0)
	SI		0 (0)	0 (0)
	PB		0 (0)	0 (0)
	N		6	3
<b>Anxiety Associated with a Life-threatening Illness</b>				
Blinded Placebo (0 mg)	PI	0 (0)	---	---
	SI	0 (0)		
	PB	0 (0)		
	N	1		
Blinded Active Dose (125 mg)	PI	0 (0)	0 (0)	---
	SI	0 (0)	0 (0)	
	PB	0 (0)	0 (0)	
	N	1	1	

PI=Positive Ideation, SI=Serious Ideation, PB=Positive Behavior, N=Number of Subjects

About a third of active dose PTSD subjects and one comparator dose PTSD subject continued to experience suicidal ideation at the primary endpoint 1 month after treatment, but this was only serious in one case. The prevalence of suicidal ideation remained consistent at long-term follow-up, and was comparable to the pre-drug preparatory period after medication washout. Only one



autistic subject reported positive ideation during the study as a result of ending the therapeutic relationship to date.

Only five cases of suicidal ideation have been considered clinically significant across sponsor-supported studies in 122 people. Two AEs were rated serious and were not related to study drug. One SAE was reported 12 days after treatment with 30 mg MDMA and lasted 6 days, concurrent with a major depressive episode that was triggered by external trauma cues, and was treated with prescription medication and hospitalization. The other SAE was reported 9 months after treatment during the long-term follow-up period, lasted 3 days and resulted in hospitalization. Three AEs of suicidal ideation were reported during the treatment period (2 in MP-12, 1 in MAA-1), one moderate AE started on the day of an active dose experimental session and lasted 1 week, one mild AE started 1 month after the last active dose experimental session and lasted 2 days, one moderate AE started 27 days after active dose treatment, lasted 12 days and resolved after treatment with prescription medication and therapy. All cases resolved without development of suicidal behavior.

Overall the incidence of serious suicidal ideation or behavior in sponsor-supported studies is low, occurring in only a few subjects post-MDMA treatment, and returning to non-life-threatening scores while subjects were closely monitored. Given that severe PTSD sufferers are known to experience suicidal ideation and behavior, it is difficult to identify a single cause of the increase in suicidal thinking or behavior (i.e. exacerbation of PTSD symptoms or from MDMA-stimulated effects). A large percentage of people enrolled in the studies reported suicidal ideation and behavior during sometime in their lives prior to study enrollment, which may reflect a manifestation of PTSD or co-morbid affective disorders. When positive serious ideation or behavior occurred after enrollment, the investigators made follow-up observations of C-SSRS to ensure subject safety, and tracked scores until they returned to non-serious levels.

The Beck Depression Inventory-II (BDI-II) is a widely used self-administered measure of depression and includes an item on suicidal ideation. Subjects’ depression levels were evaluated at baseline and at endpoints throughout the study, as a secondary measure of effectiveness of treatment. Tables 18 through 20 below show mean BDI-II scores for subjects in MP-4, MP-8, MP-9, MP-12, and MAA-1. Scores of 13 or lower indicate minimal, 14 to 19 mild, 20 to 28 moderate, 29 and above indicate severe depression symptoms.

**Table 18: Mean BDI-II Scores at Baseline, Primary Endpoint, and End of Stage 1 by Dose for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015**

Condition	Baseline	Primary Endpoint	End of Stage 1
0 mg	32.5 (6.4) N=2	28.5 (9.2) N=2	---
25 mg	17.0 (0.0) N=2	15.5 (5.0) N=2	---
30 mg	30.4 (13.7) N=7	25.8 (12.2) N=6	20.0 (15.9) N=3
40 mg	23.8 (6.2) N=6	12.8 (6.9) N=5	---
75 mg	24.7 (12.6) N=7	10.3 (6.7) N=6	7.0 (5.7) N=2
100 mg	29.3 (14.2) N=8	21.9 (16.7) N=7	12.7 (9.1) N=7
125 mg	32.2 (11.0) N=30	15.8 (13.5) N=30	10.4 (12.0) N=19

As depression is not the primary indication in sponsor-supported studies, only a subset of subjects presented with clinically significant co-morbid depression at baseline, which contributes to variation within each dose group. Statistical tests have yet to be conducted, but scores appear to be trending downward in most active MDMA dose groups, indicating an improvement in depression symptoms on average.

**Table 19: Mean BDI-II Scores at Secondary Endpoint, End of Stage 2, and 12-month Follow-up for Studies MP-4, MP-8, MP-9, and MP-12 as of 01 October 2015**

Condition	Secondary Endpoint	End of Stage 2	12-month Follow-up
<b>Stage 1/Stage 2</b>			
0 mg/125 mg	---	---	---
30 mg/125 mg	16.5 (11.1) N=6	19.5 (13.0) N=6	17.2 (13.9) N=5
40 mg/125 mg	6.8 (6.2) N=5	9.6 (9.0) N=5	0.5 (0.7) N=2
75 mg/125 mg	9.8 (11.4) N=6	6.0 (6.3) N=5	11.2 (10.1) N=5
125 mg	---	---	12.3 (11.5) N=13

Stage 2 crossover data after initial treatment with placebo or comparator shows that depression scores are in the minimal to mild range on average after active dose treatment. Most subjects receive active dose treatments in either Stage 1 or Stage 2 and continue to long-term follow-up in PTSD studies. Depression scores remain in the minimal to mild range at 12-month follow-up, suggesting that improvements in depression observed during treatment are durable on average.

**Table 20: Mean BDI-II Scores After MDMA or Placebo in MAA-1 as of 01 October 2015**

Condition	Baseline Mean (SD)	1 Day Post Session 1 Mean (SD)	2 Weeks Post Session 1 Mean (SD)	1 Month Post Session 1 Mean (SD)
Placebo	3.0 (1.4) N=2	1.5 (2.1) N=2	0.0 (0.0) N=2	0.0 (0.0) N=2
MDMA (75-125 mg)	25.0 (18.1) N=3	4.7 (3.5) N=3	6.7 (9.8) N=3	11.7 (17.6) N=3
Condition		1 Day Post Session 2 Mean (SD)	2 Weeks Post Session 2 Mean (SD)	1 Month Post Session 2 Mean (SD)
Placebo		0.0 (0.0) N=2	0.0 (0.0) N=2	0.0 (0.0) N=2
MDMA (75-125 mg)		7.0 (3.5) N=3	5.0 (6.3) N=3	2.0 (2.8) N=2

MDMA does not worsen symptoms of depression in people exhibiting moderate to severe co-morbid depression, and may have an acute antidepressant effect in this sub-group. In most cases, symptom scores declined or remained at similar levels after MDMA-assisted psychotherapy. Some subjects experienced transient positive suicidal ideation during treatment, with these scores declining throughout the course of psychotherapy, as discussed in [Section 5.3.8.5 Suicidal Ideation, Behavior, and Depression](#) above. Taken together with C-SSRS findings that do not suggest a general increase in suicidality, improvements in depression scores indicate that MDMA-assisted psychotherapy does not exacerbate or provoke symptoms of suicidality or depression.

### 5.3.9 Adverse Events

#### 5.3.9.1 Commonly Reported Adverse Events

Common AEs of MDMA reported in non-sponsor supported Phase 1 studies in healthy volunteers include elevation in blood pressure and heart rate, increased anxiety or dysphoria, and dilated pupils [8-10, 12]. Some reports indicated decreased rather than increased alertness [8]. Other common AEs reported in controlled studies of MDMA include reduced appetite, dizziness, tight jaw, bruxism (tooth-grinding), disturbance in attention, impaired gait or balance, dry mouth, and thirst. Subjects in some studies also reported or exhibited changes in cognition, such as increases in speed of thought or thought blocking, facilitated imagination or recall [13], and unusual thoughts or ideas [12]. Other less commonly reported events include paresthesia (unusual body sensations) such as tingling, or feeling hot or cold. MDMA can produce anxiety in healthy volunteers [10, 12, 13]. These effects are transient and recede as drug effects wane. One study found that women were more likely than men to experience the most commonly reported adverse effects of MDMA, though men were more likely than women to experience the specific AEs of nausea and sweating [10]. Kirkpatrick and colleagues examined a pooled sample of 220 healthy volunteers from three laboratories and failed to find gender differences in subjective or cardiovascular effects [589].

The most commonly reported AEs from Phase 1 studies published between 1986 and 2012 were used to develop a list of common reactions, or Spontaneously Reported Reactions, to record daily occurrence, duration and severity [12, 13, 28, 44, 62, 205, 208, 527, 556, 559, 560, 563]. Based on the reports summarized in Table 18, 24 reactions were identified to be tracked during sponsor-supported studies MP-1 and MP-2, and three were added after examining data from the first sponsor-supported study in a PTSD sample (MP-1). The investigators noted that subjects in MP-1 reported greater incidence of Diarrhea and Muscle Tightness, which were added to the list, and further observation led to the addition of Judgment Impaired. The following 27 reactions listed by preferred terms were tracked in MP-4, MP-8, MP-9, MP-12, MAA-1, MDA-1, and MT-1: decreased appetite, diarrhea, dry mouth, judgment impaired, muscle tightness (jaw), muscle tightness, disturbance in attention, thirst, restlessness, disturbed gait, depressed mood, dizziness, hyperhidrosis, feeling cold, obsessive ruminations, sensation of heaviness, somnolence, nystagmus, paresthesia, nausea, anxiety, irritability, insomnia, asthenia, fatigue, hypersomnia, and headache.

**Table 21: Mean Percentage of Subjects Reporting Commonly Reported Reactions During MDMA or Placebo Treatment Collected from 12 Phase 1 Studies Conducted Outside of Sponsor Support**

Treatment Group		Placebo	MDMA		
Subjects		57	174		
Reaction	Preferred Term	Mean%	Mean%	Min%	Max%
Anxiety	Anxiety	0%	19%	14%	50%
Difficulty concentrating	Disturbance in attention	16%	53%	3%	88%
Dizziness	Dizziness	2%	43%	21%	75%
Drowsiness	Somnolence	50%	26%	14%	50%
Dry mouth	Dry mouth	N/A	64%	57%	88%
Fatigue	Fatigue	26%	15%	7%	50%
Feeling cold	Feeling cold	4%	43%	23%	75%
Weakness	Asthenia	0%	16%	3%	36%
Headache	Headache	0%	11%	0%	50%
Heavy legs	Sensation of heaviness	0%	38%	38%	38%
Impaired balance/gait	Disturbed gait	0%	44%	10%	71%
Insomnia	Insomnia	0%	17%	0%	31%
Jaw clenching/tight	Muscle tightness (jaw)	0%	60%	44%	76%
Lack of appetite	Decreased appetite	2%	68%	50%	97%
Lack of energy	Decreased energy	14%	14%	3%	50%
Muscle ache/tension	Muscle tightness	N/A	20%	0%	50%
Nausea	Nausea	4%	21%	8%	36%
Nystagmus	Nystagmus	N/A	23%	3%	80%
Parasthesia	Parasthesia	0%	22%	3%	75%
Ruminations	Obsessive ruminations	23%	38%	38%	38%
Perspiration	Hyperhidrosis	0%	40%	0%	50%
Restlessness	Restlessness	0%	46%	29%	69%
Sensitivity to cold	Feeling cold	7%	38%	38%	38%
Thirst	Thirst	4%	48%	38%	63%
Restless legs	Restless legs syndrome	0%	45%	44%	46%
Palpitations	Palpitations	0%	37%	21%	63%
Hot flashes	Feeling hot	0%	23%	23%	23%
Trismus	Trismus	N/A	21%	3%	57%
Inner tension	Tension	0%	18%	3%	50%
Urge to urinate	Micturition urgency	8%	15%	15%	15%
Tremor	Tremor	0%	22%	3%	56%
Forgetfulness	Memory impairment	0%	15%	3%	38%
Brooding	Obsessive rumination	0%	12%	3%	29%

In sponsor-supported Phase 2 studies, researchers record any spontaneous (unsolicited) report of common reactions on the day of each experimental session and 7 days after. The same severity coding system for AEs was employed throughout all studies, based on limitation in daily function. [Table 19](#) and [Table 20](#) above display data from 342 experimental sessions, with each subject receiving between one and six experimental sessions at different doses across Stage 1 and Stage 2. More subjects received the 100 mg to 125 mg initial dose due to additional open-label experimental sessions offered to subjects randomized to comparator and medium dose in blinded experimental sessions.

**Table 22: Percentage of Observations of Most Commonly Reported Spontaneously Reported Reactions During Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015**

Dose Subjects	Placebo (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart
<b>Sessions</b>	24	19	15	11	14	255	4
<b>Muscle Tightness (jaw)</b>							
Mild	2 (8%)	1 (5%)	---	1 (9%)	4 (29%)	62 (24%)	---
Moderate	3 (13%)	---	---	2 (18%)	1 (7%)	73 (29%)	1 (25%)
Severe	---	---	---	---	---	6 (2%)	1 (25%)
<b>Total</b>	5 (21%)	1 (5%)	0 (0%)	3 (27%)	5 (36%)	141 (55%)	2 (50%)
<b>Anxiety</b>							
Mild	1 (4%)	---	---	2 (18%)	4 (29%)	48 (19%)	1 (25%)
Moderate	9 (38%)	---	7 (47%)	2 (18%)	1 (7%)	61 (24%)	---
Severe	4 (17%)	---	1 (7%)	---	1 (7%)	13 (5%)	---
<b>Total</b>	14 (58%)	0 (0%)	8 (53%)	4 (36%)	6 (43%)	122 (48%)	1 (25%)
<b>Decreased Appetite</b>							
Mild	1 (4%)	2 (11%)	3 (20%)	---	4 (29%)	63 (25%)	---
Moderate	1 (4%)	1 (5%)	---	---	---	40 (16%)	1 (25%)
Severe	---	1 (5%)	---	---	---	3 (1%)	---
<b>Total</b>	2 (8%)	4 (21%)	3 (20%)	0 (0%)	4 (29%)	106 (42%)	1 (25%)
<b>Headache</b>							
Mild	5 (21%)	1 (5%)	6 (40%)	3 (27%)	10 (71%)	57 (22%)	1 (25%)
Moderate	7 (29%)	---	1 (7%)	1 (9%)	---	37 (15%)	---
Severe	---	---	---	---	---	1 (<1%)	---
<b>Total</b>	12 (50%)	1 (5%)	7 (47%)	4 (36%)	10 (71%)	95 (37%)	1 (25%)
<b>Fatigue</b>							
Mild	3 (13%)	4 (21%)	6 (40%)	---	2 (14%)	32 (13%)	---
Moderate	7 (29%)	---	2 (13%)	3 (27%)	2 (14%)	52 (20%)	---
Severe	---	---	---	---	---	3 (1%)	---
<b>Total</b>	10 (42%)	4 (21%)	8 (53%)	3 (27%)	4 (28%)	87 (34%)	0 (0%)
<b>Muscle Tightness</b>							
Mild	1 (4%)	---	5 (33%)	1 (9%)	1 (7%)	44 (17%)	---
Moderate	2 (8%)	---	2 (13%)	2 (18%)	2 (14%)	25 (10%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	3 (13%)	0 (0%)	7 (47%)	3 (27%)	3 (21%)	69 (27%)	0 (0%)
<b>Nausea</b>							
Mild	2 (8%)	2 (11%)	1 (7%)	---	2 (14%)	34 (13%)	1 (25%)
Moderate	1 (4%)	---	2 (13%)	---	---	28 (11%)	---
Severe	---	---	---	---	---	6 (2%)	---
<b>Total</b>	3 (13%)	2 (11%)	3 (20%)	0 (0%)	2 (14%)	68 (27%)	1 (25%)
<b>Feeling Cold</b>							
Mild	2 (8%)	1 (5%)	7 (47%)	---	4 (29%)	47 (18%)	---
Moderate	1 (4%)	---	1 (7%)	---	2 (14%)	20 (8%)	---
Severe	---	---	---	---	---	1 (<1%)	---
<b>Total</b>	3 (13%)	1 (5%)	8 (53%)	0 (0%)	6 (43%)	68 (27%)	0 (0%)

Source: [Table 31](#)

The sponsor has analyzed the cumulative frequency of AEs commonly reported during each experimental session, collected as Spontaneously Reported Reactions. Most spontaneously reported reactions were rated as mild in studies across populations. The most frequently reported

acute and sub-acute reactions related to 255 experimental sessions with 100 mg to 125 mg initial dose of MDMA in 100 people across Phase 2 studies were muscle tightness (jaw) (141 reports during 55% of sessions at any severity, 2% severe), anxiety (122 reports during 48% of sessions at any severity, 5% severe), decreased appetite (106 reports during 42% of sessions at any severity, 1% severe), headache (95 reports during 37% of sessions at any severity, <1% severe), fatigue (87 reports during 34% of sessions at any severity, 1% severe), muscle tightness (69 reports during 27% of sessions at any severity, none severe), and nausea (68 reports during 27% of sessions at any severity, 2% severe). The next most common reactions reported during 9% to 25% of experimental sessions with a 100 mg to 125 mg initial dose of MDMA in order of frequency are: hyperhidrosis, restlessness, dizziness, insomnia, thirst, disturbed gait, dry mouth, disturbance in attention, depressed mood, and nystagmus, described in [Table 31](#). The highest initial dose of 150 mg MDMA was only administered during four experimental sessions in MP-2, and was associated with reports of insomnia (3), muscle tightness (jaw) (2), dizziness (2), disturbed gait (2), dry mouth (2), and thirst (2), but it should be noted that this group is small. The following reactions were reported during less than 9% of experimental sessions with 100 mg to 125 mg initial dose of MDMA on the day of drug administration: feeling cold, obsessive ruminations, sensation of heaviness, somnolence, paresthesia, diarrhea, judgment impaired, irritability, asthenia, and hypersomnia. These reactions may be of less concern than previously proposed in the scientific literature on MDMA.

In studies where a low dose of MDMA (25 mg to 40 mg) was administered in 20 subjects across 45 sessions, infrequent reports of fatigue (15), anxiety (12), headache (12), muscle tightness (10), and feeling cold (9) were observed. In comparison, 12 subjects who received an inactive placebo in 24 experimental sessions reported anxiety (14), insomnia (12), headache (12), fatigue (10), and muscle tightness (jaw) (5) during experimental sessions. Taking into consideration that the 100 mg to 125 mg MDMA dose has been administered by far the most frequently, the sponsor concludes that the frequency of spontaneously reported reactions are likely to be most accurate in the 100 mg to 125 mg dose experimental sessions that have been administered to date in these studies. The higher number of experimental sessions at this dose meant that there was greater opportunity to report reactions. While 100 mg to 125 mg MDMA was associated with more reactions overall, these reactions were self-limiting and generally did not persist beyond the 7-day window after experimental sessions, unless associated with medical history.



**Table 23: Percentage of Observations of Most Commonly Reported Spontaneously Reported Reactions During Telephone Contact on Day 1-7 After Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015**

Dose Subjects	Placebo (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart
<b>Observations</b>	168	133	105	77	98	1785	21
<b>Anxiety</b>							
Mild	26 (15%)	4 (3%)	25 (24%)	8 (10%)	5 (5%)	239 (13%)	---
Moderate	35 (21%)	---	10 (10%)	3 (4%)	1 (1%)	172 (10%)	---
Severe	2 (1%)	---	---	---	---	24 (1%)	---
Total	63 (38%)	4 (3%)	35 (33%)	11 (14%)	6 (6%)	435 (24%)	0 (0%)
<b>Fatigue</b>							
Mild	23 (14%)	16 (12%)	20 (19%)	3 (4%)	27 (28%)	245 (14%)	2 (10%)
Moderate	27 (16%)	8 (6%)	8 (8%)	4 (5%)	1 (1%)	163 (9%)	14 (67%)
Severe	1 (1%)	1 (1%)	---	---	---	13 (1%)	2 (10%)
Total	51 (30%)	25 (19%)	28 (27%)	7 (9%)	28 (29%)	421 (24%)	18 (86%)
<b>Insomnia</b>							
Mild	19 (11%)	17 (13%)	8 (8%)	5 (6%)	6 (6%)	158 (9%)	---
Moderate	29 (17%)	13 (10%)	9 (9%)	5 (6%)	2 (2%)	88 (5%)	---
Severe	1 (1%)	8 (6%)	1 (1%)	5 (6%)	---	8 (<1%)	---
Total	49 (29%)	38 (29%)	18 (17%)	15 (19%)	8 (8%)	254 (14%)	0 (0%)
<b>Depressed Mood</b>							
Mild	13 (8%)	14 (11%)	4 (4%)	3 (4%)	---	114 (6%)	3 (14%)
Moderate	8 (5%)	11 (8%)	4 (4%)	---	---	101 (6%)	---
Severe	---	---	---	---	---	17 (1%)	---
Total	21 (13%)	25 (19%)	8 (8%)	3 (4%)	0 (0%)	232 (13%)	3 (14%)
<b>Hypersomnia</b>							
Mild	15 (9%)	8 (6%)	9 (9%)	3 (4%)	17 (17%)	145 (8%)	2 (10%)
Moderate	9 (5%)	7 (5%)	1 (1%)	3 (4%)	---	63 (4%)	---
Severe	---	---	1 (1%)	---	---	---	---
Total	24 (14%)	15 (11%)	11 (10%)	6 (8%)	17 (17%)	208 (12%)	2 (10%)
<b>Disturbance in Attention</b>							
Mild	11 (7%)	---	2 (2%)	2 (3%)	---	141 (8%)	---
Moderate	11 (7%)	---	2 (2%)	2 (3%)	---	38 (2%)	---
Severe	---	---	---	---	---	3 (<1%)	---
Total	22 (13%)	0 (0%)	4 (4%)	4 (5%)	0 (0%)	182 (10%)	0 (0%)
<b>Decreased Appetite</b>							
Mild	---	10 (8%)	2 (2%)	1 (1%)	2 (2%)	89 (5%)	---
Moderate	---	2 (2%)	---	1 (1%)	---	67 (4%)	---
Severe	---	3 (2%)	---	---	---	1 (<1%)	---
Total	0 (0%)	15 (11%)	2 (2%)	2 (3%)	2 (2%)	157 (9%)	0 (0%)
<b>Dizziness</b>							
Mild	5 (3%)	6 (5%)	---	---	1 (1%)	122 (7%)	---
Moderate	---	1 (1%)	---	---	---	19 (1%)	---
Severe	---	---	---	---	---	2 (<1%)	---
Total	5 (3%)	7 (5%)	0 (0%)	0 (0%)	1 (1%)	143 (8%)	0 (0%)

<b>Dose Subjects</b>	<b>Placebo (N=12)</b>	<b>25 mg (N=7)</b>	<b>30 mg (N=7)</b>	<b>40 mg (N=6)</b>	<b>75 mg (N=7)</b>	<b>100-125 mg (N=100)</b>	<b>150 mg (N=3)</b>
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-3 doses 3-5 weeks apart	1-2 doses 3-5 weeks apart
<b>Observations</b>	168	133	105	77	98	1785	21
<b>Irritability</b>							
Mild	12 (7%)	3 (2%)	5 (5%)	---	4 (4%)	69 (4%)	---
Moderate	9 (5%)	---	3 (3%)	3 (4%)	---	48 (3%)	---
Severe	---	---	---	---	---	2 (<1%)	---
<b>Total</b>	<b>21 (13%)</b>	<b>3 (2%)</b>	<b>8 (8%)</b>	<b>3 (4%)</b>	<b>4 (4%)</b>	<b>119 (7%)</b>	<b>0 (0%)</b>
<b>Headache</b>							
Mild	9 (5%)	13 (10%)	1 (1%)	3 (4%)	4 (4%)	76 (4%)	---
Moderate	5 (3%)	13 (10%)	---	3 (4%)	---	33 (2%)	---
Severe	---	---	---	---	---	3 (<1%)	---
<b>Total</b>	<b>14 (8%)</b>	<b>26 (20%)</b>	<b>1 (1%)</b>	<b>6 (8%)</b>	<b>4 (4%)</b>	<b>112 (6%)</b>	<b>0 (0%)</b>
<b>Muscle Tightness (jaw)</b>							
Mild	2 (1%)	---	---	6 (8%)	2 (2%)	90 (5%)	---
Moderate	1 (1%)	---	---	4 (5%)	---	20 (1%)	---
Severe	---	---	---	---	---	2 (<1%)	---
<b>Total</b>	<b>3 (2%)</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>10 (13%)</b>	<b>2 (2%)</b>	<b>112 (6%)</b>	<b>0 (0%)</b>

Source: [Table 32](#)

The sponsor has analyzed the cumulative frequency of Spontaneously Reported Reactions reported during 7 days following each experimental session. The most frequently reported reactions related to 255 experimental sessions with 100 mg to 125 mg initial dose of MDMA in 100 people across Phase 2 studies were anxiety (24% of observations, 1% severe), fatigue (24% of observations, 1% severe), insomnia (14% of observations, <1% severe), depressed mood (13% of observations, 1% severe), hypersomnia (12% of observations, none severe), disturbance in attention (10% of observations, <1% severe), decreased appetite (9% of observations, <1% severe), dizziness (8% of observations, <1% severe), irritability (7% of observations, <1% severe), headache (6% of observations, <1% severe), muscle tightness (jaw and elsewhere) (6% of observations, <1% severe). The next most common reactions reported during the week after experimental sessions with 100 mg to 125 mg initial dose of MDMA in less than 6% of daily telephone contacts, in order of frequency, are: muscle tightness, nausea, obsessive ruminations, restlessness, asthenia, feeling cold, diarrhea, dry mouth, judgment impaired, disturbed gait, hyperhidrosis, sensation of heaviness, somnolence, nystagmus, parasthesia, and thirst, as described in [Table 32](#). The highest initial dose of 150 mg MDMA was only administered during four experimental sessions in MP-2, and was associated with reports of fatigue (86% of observations), depressed mood (14% of observations), hypersomnia (10% of observations), and dry mouth (5% of observations) during the week following experimental sessions.

In studies where a low dose of 25 mg to 40 mg initial dose of MDMA was administered, infrequent reports of insomnia (22% of observations), fatigue (19% of observations), anxiety (16% of observations), depressed mood (11% of observations), headache (10% of observations), hypersomnia (10% of observations), muscle tightness (6% of observations), nausea (6% of observations), and decreased appetite (6% of observations). The following reactions were observed in 4% or less of daily telephone contact observations, in order of frequency: obsessive ruminations, irritability, muscle tightness (jaw), disturbance in attention, restlessness, dizziness, feeling cold, diarrhea, somnolence, judgment impaired, asthenia, thirst, dry mouth, hyperhidrosis, and sensation of heaviness. In comparison, 12 subjects who received an inactive placebo in 24 experimental sessions reported anxiety (38% of observations), fatigue (30% of observations),

insomnia (29% of observations), hypersomnia (14% of observations), depressed mood (13% of observations), disturbance in attention (13% of observations), and irritability (13% of observations) were reported during daily contact for 1 week following each experimental session. Headache, nausea, muscle tightness, somnolence, obsessive rumination, dizziness, muscle tightness (jaw), diarrhea, disturbed gait, and hyperhidrosis were reported in less than 10% of observations during daily telephone contact. While 100 mg to 125 mg MDMA was associated with more reactions overall, these reactions were self-limiting and generally did not persist beyond the 7-day window after experimental sessions.

Any reactions that continued beyond the 7-day window were tracked as unexpected AEs until they returned to baseline levels. In all studies to date, 18 severe reactions lasted beyond the 7-day window: insomnia (2 lasting up to 26 days), anxiety (6 lasting up to 53 days), restlessness (2 lasting up to 18 days), obsessive rumination (1 lasting 4 days) and depressed mood (4 lasting up to 51 days), headache (1 lasting 13 days), muscle tightness (jaw) (1 lasting 20 days), and muscle tightness (1 lasting 20 days). These reactions were tracked as AEs until resolution and subjects experiencing them were provided with prescription medication and additional therapy. Among the subset of AEs collected as commonly reported severe reactions, severe anxiety, insomnia, fatigue, nausea, muscle tightness, and depressed mood were reported in 4% or more subjects. Severe anxiety was reported the most during both inactive placebo (22%) and MDMA experimental sessions (5% to 10%, depending on dose). These reactions also overlap with symptoms of pre-existing conditions in medical history associated with PTSD (depression, somatic symptoms, insomnia, anxiety) and anxiety disorders, which can be exacerbated by processing traumatic content and may influence the frequency and duration of reactions observed in sponsored-supported clinical trials of MDMA-assisted psychotherapy.

### 5.3.9.2 Adverse Events

The sponsor has analyzed cumulative frequency of AEs through the data cut-off period, which included 122 subjects treated with MDMA at any dose in 10 MAPS-sponsored studies conducted under U.S. IND. See [Table 24](#) below for distribution by severity and relationship. There have been no Safety Reports to date under this IND.

**Table 24: Overview of All Adverse Events Post-Drug by Severity and Relationship in MAPS-Sponsored Studies Across Populations as of 01 October 2015**

	Related AEs (% by severity)	All AEs (% by severity)
Mild AEs	58 (34%)	117 (36%)
Moderate AEs	100 (59%)	181 (56%)
Severe AEs	12 (7%)	28 (9%)
Any Severity	170 (100%)	326 (100%)

In data collected from all studies described above, there were 170 possibly or probably related AEs out of 326 at any severity after MDMA administration. Since MDMA is administered as an adjunct to psychotherapy, judging relationship to study drug is a known challenge for this combined therapy. In the context of complex medical histories associated with the PTSD diagnosis, somatic symptoms may wax and wane independent of treatment. In addition, it is known that processing trauma during psychotherapy for PTSD, with or without concomitant pharmacological treatment, can temporarily increase symptoms as an expected aspect of the therapeutic process. This is borne out by the high incidence of spontaneously reported reactions and AEs in the placebo group. Possibly or probably related AEs were more often moderate than mild or severe. Multiple severe AEs were rarely reported by the same subject. In [Table 25](#) below,

body systems of AEs reported by 2% or more of subjects are displayed, with distribution and frequency of severe AEs by body system.

**Table 25: Body Systems of All Adverse Events Post-Drug Reported by 2% or More of Subjects in MAPS-Sponsored Studies Across Populations**

Dose	Placebo (0 mg)		Comparator Dose (25-40 mg)		Active Dose (75-150 mg)		Any MDMA Dose (25-150 mg)	
Schedule	1-2 doses 3-5 weeks apart		1-3 doses 3-5 weeks apart		1-6 doses 3-5 weeks apart		1-6 doses 3-5 weeks apart	
Subjects	14		24		141		122	
Sessions	27		49		306		355	
Relationship to Drug	PR	NR	PR	NR	PR	NR	PR	NR
<b>Cardiac Disorders</b>	---	---	---	---	2 (1%)	1 (1%)	2 (2%)	1 (1%)
Severe					(0)	(0)	(0)	(0)
<b>Ear and Labyrinth Disorders</b>	---	---	1 (4%)	---	1 (1%)	1 (1%)	2 (2%)	1 (1%)
Severe			(0)		(0)	(0)	(0)	(0)
<b>Eye Disorders</b>	---	---	---	---	9 (6%)	1 (1%)	9 (7%)	1 (0%)
Severe					(0)	(0)	(0)	(0)
<b>Gastrointestinal Disorders</b>	1 (7%)	---	1 (4%)	---	17 (12%)	13 (9%)	18 (15%)	13 (11%)
Severe	(0)		(0)		1 (1%)	(0)	1 (1%)	(0)
<b>General Disorders and Administration Site Conditions</b>	4 (29%)	---	4 (17%)	4 (17%)	21 (15%)	10 (7%)	25 (20%)	14 (11%)
Severe	(0)		(0)	1 (4%)	(0)	(0)	(0)	1 (1%)
<b>Infections and Infestations</b>	1 (7%)	4 (29%)	---	5 (21%)	6 (4%)	12 (9%)	6 (5%)	17 (14%)
Severe	(0)	(0)		(0)	(0)	2 (1%)	(0)	2 (1%)
<b>Injury, Poisoning, and Procedural Complications</b>	---	1 (7%)	---	---	2 (1%)	7 (5%)	2 (2%)	7 (6%)
Severe		(0)			(0)	1 (1%)	(0)	1 (1%)
<b>Metabolism and Nutrition Disorders</b>	---	---	---	1 (4%)	2 (1%)	2 (1%)	2 (2%)	3 (2%)
Severe				(0)	(0)	(0)	(0)	(0)
<b>Musculoskeletal and Connective Tissue Disorders</b>	8 (57%)	2 (14%)	1 (4%)	---	20 (14%)	13 (9%)	21 (17%)	13 (11%)
Severe	1 (7%)	(0)	(0)		(0)	1 (1%)	(0)	1 (1%)
<b>Neoplasms Benign, Malignant, and Unspecified</b>	---	---	---	---	---	2 (1%)	---	2 (2%)
Severe						2 (1%)		2 (2%)
<b>Nervous System Disorders</b>	1 (7%)	3 (21%)	3 (13%)	1 (4%)	12 (9%)	9 (6%)	15 (12%)	10 (8%)
Severe	(0)	1 (7%)	2 (8%)	(0)	(0)	(0)	2 (2%)	(0)
<b>Psychiatric Disorders</b>	9 (64%)	2 (14%)	11 (46%)	7 (29%)	47 (33%)	44 (31%)	58 (48%)	51 (42%)
Severe	1 (7%)	(0)	2 (8%)	3 (13%)	7 (5%)	5 (4%)	9 (7%)	8 (7%)
<b>Renal and Urinary Disorders</b>	---	---	---	---	2 (1%)	---	2 (2%)	---
Severe					(0)		(0)	

<b>Dose</b>	<b>Placebo (0 mg)</b>		<b>Comparator Dose (25-40 mg)</b>		<b>Active Dose (75-150 mg)</b>		<b>Any MDMA Dose (25-150 mg)</b>	
<b>Schedule</b>	1-2 doses 3-5 weeks apart		1-3 doses 3-5 weeks apart		1-6 doses 3-5 weeks apart		1-6 doses 3-5 weeks apart	
<b>Subjects</b>	14		24		141		122	
<b>Sessions</b>	27		49		306		355	
<b>Reproductive System and Breast Disorders</b>	---	---	---	---	---	4 (3%)	---	4 (3%)
Severe						1 (1%)		1 (1%)
<b>Respiratory, Thoracic, and Mediastinal Disorders</b>	1 (7%)	---	2 (8%)	---	2 (1%)	9 (6%)	4 (3%)	9 (7%)
Severe	(0)		(0)		(0)	(0)	(0)	(0)
<b>Skin and Subcutaneous Tissue Disorders</b>	1 (7%)	1 (7%)	---	1 (4%)	4 (3%)	3 (2%)	4 (3%)	4 (3%)
Severe	(0)	(0)		(0)	(0)	(0)	(0)	(0)
<b>Vascular Disorders</b>	---	---	---	---	---	3 (2%)	---	3 (2%)
Severe						(0)		(0)

PR=Possibly or probably related, NR=Not related



Across all body systems, most related AEs reported at any dose of MDMA were psychiatric disorders (48% of MDMA versus 64% of placebo subjects), followed by general disorders and administration site conditions (20% of MDMA versus 29% of placebo subjects), musculoskeletal and connective tissue disorders (17% of MDMA versus 57% of placebo subjects), gastrointestinal disorders (15% of MDMA versus 7% of placebo subjects), nervous system disorders (12% of MDMA versus 7% of placebo subjects), eye disorders (7% of MDMA versus none of placebo subjects), and infections and infestations (5% of MDMA versus 7% of placebo subjects). See [Table 26](#) below for details of related AEs under each system organ class. Based on comparison of frequencies, taking into account that sample sizes are heavily weighted towards active dose MDMA due to study design, gastrointestinal disorders, nervous system disorders, and eye disorders appear to be associated with more MDMA subjects over placebo. Based on the elimination half-life of 7 to 9 hours for active doses of MDMA, it is difficult to judge relationship of AEs reported after the 7-day safety window as they may also be related to the therapeutic process or medical history. Investigators tended to be more conservative and judged events to be related based on known pharmacodynamics of MDMA, for example with gastrointestinal disorders and the distribution of serotonin receptors in the gut [622].

A majority of the AEs were psychiatric disorders. Given study inclusion criteria requiring a pre-existing diagnosis of chronic anxiety or PTSD and the fact that subjects were receiving MDMA, a drug that is known to increase general anxiety in an average of 19% healthy volunteers across multiple Phase 1 studies, these AEs are expected. However, the frequency of psychiatric disorders in the small group of subjects who received inactive placebo was even higher than the active doses, suggesting that these AEs may, at least in part, be related to exacerbation of medical history diagnoses during the study independent of MDMA administration.

Related AEs reported in 3% or less of MDMA subjects were: respiratory, thoracic and mediastinal disorders, skin and subcutaneous tissue disorders, cardiac, ear and labyrinth, injury, poisoning and procedural complications, metabolism and nutrition disorders, renal and urinary disorders. It is noteworthy that, although there was one related, moderate, expected cardiac AE that was deemed serious because it led to overnight monitoring of increased ventricular extrasystoles, no severe cardiac, renal and urinary, or vascular disorders were reported, and they were also the least frequently reported types of AEs after any MDMA dose, in contrast to reports of cardiovascular toxicity, hyperthermia, ARF, hyponatremia, and neurotoxicity in epidemiological settings, as described in [Section 4.5 Serious Reports, Mortality, and Morbidity in Animals and Epidemiological Settings](#) and in [Table 2](#). The difference in frequency suggests that AEs in these body systems are likely to be rare in a controlled clinical setting with proper medical screening, and that they may receive disproportionate coverage in the scientific literature on epidemiological studies due to significant impact on the body. In [Table 26](#) below, preferred terms of AEs possibly or probably related to comparator and active doses of study drug are presented for a more detailed view.

**Table 26: Related Adverse Events in Sponsor Supported Studies of MDMA-Assisted Psychotherapy Across Populations Organized by Body System as of 01 October 2015**

<b>Dose</b>	<b>Comparator Dose (25-40 mg)</b>	<b>Active Dose (75-150 mg)</b>	<b>Any MDMA Dose (25-150 mg)</b>
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart
<b>Subjects</b>	22	121	122
<b>Sessions</b>	45	306	355
<b>Cardiac Disorders</b>			
Palpitations	---	1 (1%)	1 (1%)
Ventricular extrasystoles	---	1 (1%)	1 (1%)
<b>Ear and Labyrinth Disorders</b>			
Tinnitus	1 (5%)	---	1 (1%)
<b>Eye Disorders</b>			
Visual impairment	---	5 (4%)	5 (4%)
Vitreous floaters	---	1 (1%)	1 (1%)
Dry eyes/abnormal sensation in eye	---	1 (1%)	1 (1%)
Vision blurred	---	2 (2%)	2 (2%)
<b>Gastrointestinal Disorders</b>			
Diarrhea	1 (5%)	4 (3%)	5 (4%)
Dyspepsia	---	3 (2%)	3 (2%)
Abdominal pain	---	3 (2%)	3 (2%)
Nausea	---	2 (2%)	2 (2%)
Oropharyngeal blistering	---	1 (1%)	1 (1%)
Vomiting	---	4 (3%)	4 (3%)
<b>General Disorders and Administration Site</b>			
Asthenia	---	1 (1%)	1 (1%)
Fatigue	4 (18%)	9 (7%)	13 (11%)
Feeling abnormal	---	1 (1%)	1 (1%)
Feeling hot	---	2 (2%)	2 (2%)
Irritability	---	1 (1%)	1 (1%)
Pain (body aching, body tension)	---	4 (3%)	4 (3%)
Pyrexia	---	1 (1%)	1 (1%)
Chills	---	1 (1%)	1 (1%)
<b>Infections and Infestations</b>			
Pharyngitis streptococcal	---	1 (1%)	1 (1%)
Upper respiratory infection	---	3 (2%)	3 (2%)
Urinary tract Infection	---	2 (2%)	2 (2%)
<b>Injury, Poisoning, and Procedural Complications</b>			
Contusion	---	1 (1%)	1 (1%)
Skin abrasion	---	1 (1%)	1 (1%)
<b>Metabolism and Nutrition Disorders</b>			
Anorexia	---	1 (1%)	1 (1%)
Decreased appetite	---	1 (1%)	1 (1%)

<b>Dose</b>	<b>Comparator Dose (25-40 mg)</b>	<b>Active Dose (75-150 mg)</b>	<b>Any MDMA Dose (25-150 mg)</b>
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart
<b>Subjects</b>	22	121	122
<b>Sessions</b>	45	306	355
<b>Musculoskeletal and Connective Tissue Disorders</b>			
Arthralgia (joint)	---	1 (1%)	1 (1%)
Back pain	---	2 (2%)	2 (2%)
Joint stiffness	---	1 (1%)	1 (1%)
Muscle spasms	---	1 (1%)	1 (1%)
Muscle tightness	1 (5%)	7 (6%)	8 (7%)
Muscle twitches	---	1 (1%)	1 (1%)
Musculoskeletal pain (shoulder)	---	3 (2%)	3 (2%)
Myalgia	---	3 (2%)	3 (2%)
Neck pain	---	1 (1%)	1 (1%)
<b>Nervous System Disorders</b>			
Burning sensation (fingers, thighs)	---	2 (2%)	2 (2%)
Dizziness	---	1 (1%)	1 (1%)
Hangover (feeling hungover)	---	1 (1%)	1 (1%)
Headache	1 (5%)	3 (2%)	4 (3%)
Hypersomnia	1 (5%)	3 (2%)	4 (3%)
Hypoaesthesia facial	---	1 (1%)	1 (1%)
Migraine headache	1 (5%)	---	1 (1%)
Myoclonus	---	1 (1%)	1 (1%)
Tension headache	---	1 (1%)	1 (1%)
<b>Psychiatric Disorders</b>			
Agitation	---	1 (1%)	1 (1%)
Anxiety	4 (18%)	22 (18%)	26 (21%)
Bruxism	---	2 (2%)	2 (2%)
Depressed mood	3 (14%)	4 (3%)	7 (6%)
Derealization	---	1 (1%)	1 (1%)
Dissociation	---	1 (1%)	1 (1%)
Disturbance in attention	---	1 (1%)	1 (1%)
Flashback	---	1 (1%)	1 (1%)
Hypnagogic hallucination	---	1 (1%)	1 (1%)
Hypnopompic hallucination	---	1 (1%)	1 (1%)
Intentional self-injury	---	1 (1%)	1 (1%)
Irritability	---	2 (2%)	2 (2%)
Negative thoughts	1 (5%)	---	1 (1%)
Obsessive Rumination	1 (5%)	1 (1%)	1 (1%)
Panic attack	---	4 (3%)	1 (1%)
Restlessness	---	1 (1%)	1 (1%)
Suicidal ideation	---	1 (1%)	1 (1%)
Tic (teeth tapping)	---	1 (1%)	1 (1%)
Time perception altered	1 (5%)	---	1 (1%)
Trichotillomania	1 (5%)	---	1 (1%)
<b>Renal and Urinary Disorders</b>			
Nocturia	---	1 (1%)	1 (1%)
Dysuria	---	1 (1%)	1 (1%)

Dose	Comparator Dose (25-40 mg)	Active Dose (75-150 mg)	Any MDMA Dose (25-150 mg)
<b>Schedule</b>	1-2 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart	1-6 doses 3-5 weeks apart
<b>Subjects</b>	22	121	122
<b>Sessions</b>	45	306	355
<b>Respiratory, Thoracic, and Mediastinal Disorders</b>			
Cough	1 (5%)	---	1 (1%)
Nasal congestion	1 (5%)	2 (2%)	3 (2%)
Sinus headache	---	1 (1%)	1 (1%)
<b>Skin and Subcutaneous Tissue Disorders</b>			
Petechiae	---	2 (2%)	2 (2%)
Pruritis	---	2 (2%)	2 (2%)

The most frequently reported possibly or probably related AEs were anxiety (18% in active versus 18% comparator subjects), fatigue (7% active versus 18% comparator subjects), muscle tightness (6% active versus 5% comparator subjects), and visual impairment (4% active versus none of comparator subjects). Subjective effects of MDMA are known to have a dose response relationship, so AEs with equivalent frequencies across comparator versus active doses of MDMA suggest an absence of dose response and possible relationship to medical history or therapeutic process. Since more individuals received active dose MDMA due to unequal group sizes weighted towards the active MDMA doses, and the partial crossover offered to comparator dose and placebo subjects, a greater number of AEs is expected with active doses of MDMA beyond any expected dose response relationship. In addition, frequencies in comparator dose subjects are based on a small sample of 22 subjects who received 45 experimental sessions.

Muscle tightness in the body as well as specific to the jaw was frequently reported as an unsolicited reaction during experimental sessions, as described in [Table 22](#). During the 7-day safety window, these reactions were much less frequently reported, as described in [Table 23](#) and [Table 32](#). Among related AEs reported during and after drug administration, somatic symptoms were more frequently experienced in active dose subjects, such as pain associated with body tension (3% of active dose subjects versus none of comparator subjects), muscle tightness (6% of active dose versus 5% of comparator dose subjects), musculoskeletal pain in the shoulder (2% of active dose subjects versus none of comparator), back pain (2% of active dose subjects versus none of comparator), and myalgia (2% of active dose subjects versus none of comparator). As previously discussed in [Section 5.3.9.2 Adverse Events](#), it is difficult to judge relationship between study drug and conditions associated with medical history diagnoses. Pain and somatic symptoms can be directly related to traumatic events, such as physical or sexual assault, a motor vehicle accident, or combat [\[623\]](#). A meta-analytic review and several large studies have found a robust association between PTSD and somatic symptoms, suggesting that PTSD itself may be a contributing factor beyond combat exposure, sexual, or physical abuse that lead to the PTSD [\[624-627\]](#).

Although MDMA is not a classic hallucinogen, as classified by chemical structure and mechanism of action, data from sponsor-supported studies suggest that in a clinical population mild psychoactive effects, such as hypnagogic and hypnopompic hallucinations and visual distortions may be observed in some individuals. Hallucinogenic subjective effects were not actively solicited during therapy sessions, as was done in Phase 1 studies of healthy volunteers [\[8, 10, 11, 556\]](#). Any unsolicited reports were collected as spontaneously reported reactions or AEs in sponsor-supported studies.

**Table 27: Severe Related Adverse Events in Sponsor Supported Studies of MDMA-Assisted Psychotherapy Across Populations as of 01 October 2015**

<b>Indication</b>	PTSD	Healthy	Anxiety	Social Anxiety	Total
<b>Population</b>	All	Therapists	Life-Threatening Illness	Autistic Adults	
<b>Subjects</b>	107	7	4	9	127
<b>Sessions</b>	365	7	8	19	399
<b>Psychiatric</b>					
Re-experiencing Episode	1	---	---	---	1 (1%)
Panic Attack	2	---	---	---	2 (2%)
Depressed Mood	2	---	---	---	2 (2%)
Obsessive Rumination	1	---	---	---	1 (1%)
Anxiety	3	---	---	---	3 (2%)
<b>Nervous System</b>					
Headache	1	---	---	---	1 (1%)
<b>Gastrointestinal</b>					
Abdominal Cramps/Pain	1	---	---	---	1 (1%)
<b>General</b>					
Restlessness	1	---	---	---	1 (1%)
<b>Musculoskeletal &amp; Connective Tissue</b>					
Musculoskeletal Chest Pain	1	---	---	---	1 (1%)

The sponsor has analyzed the cumulative frequency of AEs and found the most frequent severe possibly or probably related AEs to be anxiety or distress (N=3, 2% of subjects), depressed mood (N=2, 2% of subjects), and panic attacks (N=2, 2% of subjects) in sponsor-supported PTSD studies. The following severe related AEs were observed in 1% of subjects: re-experiencing episode, obsessive rumination, restlessness, headache, abdominal cramps/pain, and musculoskeletal chest pain. Severe related AEs were treated with prescription medications and followed by additional phone contact and psychotherapy to ensure that the subjects returned to baseline or were stabilized. It is noteworthy that no severe related AEs were reported in non-PTSD populations in sponsor-supported studies, which could also be attributed to small sample sizes.

### 5.3.9.3 Serious Adverse Events

Eleven SAEs have occurred across five sponsor-supported studies. These include one expected related SAE and 10 unrelated SAEs after drug administration. See [Table 28](#) below for a summary of these SAEs.

**Table 28: Serious Adverse Events in Sponsor-Supported Studies of MDMA-Assisted Psychotherapy Across Populations as of 01 October 2015**

Dose	Comparator Dose (30 mg)	Active Dose (100 mg)	Active Dose (125 mg)
System Organ Class Preferred Term	Relationship		
<b>Gastrointestinal Disorders</b>			
Appendicitis	None		1
<b>Injury, Poisoning, and Procedural Complications</b>			
Fractured Clavicle (auto accident)	None		1
Lower Limb Fracture	None	1	
<b>Nervous System Disorder</b>			
Vasovagal Syncope	None		1
<b>Neoplasms Benign, Malignant, and Unspecified</b>			
Brain Metastasis (frontal brain syndrome)	None		1
Breast Cancer	None		1
<b>Reproductive System and Breast Disorders</b>			
Ovarian Cyst Ruptured	None	1	
<b>Psychiatric Disorders</b>			
Suicidal Ideation	None	1	1
Major Depressive Episode	None		1
<b>Cardiac Disorders</b>			
Increase in Ventricular Extrasystoles	Probably		1

One related serious adverse reaction has occurred within all sponsor-supported studies to date. Subject 0811 experienced an increase in frequency of ventricular extrasystoles (PVC’s), a form of arrhythmia, on the day of his third and final experimental session with open-label 125 mg MDMA. The subject had no other signs and no symptoms of cardiac distress. In the absence of any symptoms of coronary insufficiency, the investigator judged the only medical measure necessary to be withholding the supplemental dose of MDMA. This was the final drug administration in Stage 2. No similar events occurred during the first two 125 mg experimental sessions, nor the two blinded experimental sessions with 30 mg MDMA in Stage 1. There was no evidence of acute cardiac damage or ischemia or underlying heart disease. At baseline during screening, the subject had one PVC on baseline electrocardiogram (EKG), but the EKG was otherwise normal. The subject had a family history of his father having had a coronary artery bypass graft, which had prompted the subject to consult a cardiologist several years before study enrollment, and the cardiologist’s note indicated that he did not suspect cardiovascular disease or see the need for further workup. Based on the medical history and clinical presentation of this subject, the investigator judged the SAE to be a moderate exacerbation probably related to drug administration. The event required overnight monitoring in the hospital, but did not lead to any adverse sequelae. He was given one dose of 25 mg metoprolol by the hospital physician but did not require any ongoing treatment. Serial cardiac isoenzymes, an echocardiogram and a nuclear stress test performed during the overnight hospital admission failed to show evidence of cardiovascular or other cardiac disease. Full recovery occurred 1 day after MDMA administration. Arrhythmia is described in [sections 4.5](#) and [5.3.4](#) as an expected adverse effect of MDMA.

### 5.3.10 Abuse Potential

When reviewing the effects of MDMA in a sample of 74 largely drug-naïve subjects in a study conducted outside of sponsor support, Liechti and colleagues stated that “none of the subjects expressed any interest in taking MDMA as a recreational drug” after receiving MDMA in a



controlled research setting [10]. When assessed in terms of willingness to choose money over receiving the drug, subjects previously experienced with Ecstasy provided similar responses to 2 mg/kg MDMA and 20 mg d-amphetamine, a sign of having reinforcing effects [596]. A study that enrolled subjects with a history of Ecstasy use (4 to 40 occasions) found that only self-reported feelings of playfulness were associated with subjects' desire to take MDMA in a controlled research setting [39].

The sponsor has assessed abuse potential of MDMA in Phase 2 clinical trials with collection of self-report information on Ecstasy use during long-term follow-up in all studies. In addition, one study (MP-2) incorporated random and scheduled drug testing during long-term follow-up. One subject in MP-1 who had received two experimental sessions with active dose MDMA reported the use of Ecstasy in an attempt to recreate the therapeutic setting but found the experience unsatisfactory, and after this experience indicated no desire to repeat it. No other subjects in this study reported using Ecstasy after completing the study [42]. In sponsor-supported study MP-2, drug screens specific for MDMA performed 2 months, 6 months, and 12 months after the final experimental session were negative, suggesting that study subjects did not seek out MDMA or Ecstasy after taking part in the study. Although MDMA does not demonstrate signals associated with known abuse liability patterns, the drug will only be administered in a clinic setting under continuous observation on an intermittent schedule, which further limits abuse potential.

## 5.4 Efficacy of MDMA Across Populations

### 5.4.1 PTSD

Ongoing and completed sponsor-supported studies of MDMA-assisted therapies employ recognized clinician-administered gold-standard measures of the condition or symptoms. The primary outcome measure of efficacy for studies of MDMA-assisted psychotherapy for PTSD to date is the Clinician Administered PTSD Scale (CAPS) following DSM-IV, an established semi-structured interview conducted by a trained clinician [628-630]. The Global Severity CAPS score encompasses frequency and intensity scores for three symptom domains; re-experiencing, avoidance and hyperarousal. An independent rater that does not see the subjects during any of the psychotherapy sessions administers the CAPS at baseline and at the primary endpoint, 1 or 2 months after blinded MDMA-assisted psychotherapy sessions. Secondary endpoints include an assessment 1 to 2 months after a third experimental session and 12 months after the last treatment.

Analyses of the CAPS at the primary endpoint after two experimental sessions in MP-1 found subjects receiving MDMA-assisted psychotherapy experienced a clinically and statistically significant decline in PTSD symptoms compared to placebo-assisted psychotherapy [41]. Global CAPS scores declined for all subjects over time (overall baseline mean Global CAPS=79.1±21.7, and 2 months after the second experimental session, mean Global CAPS=38.2±30.3), indicating a clinically significant drop of 40.9 points, and a 52% reduction in symptoms. People in the MDMA and placebo conditions began the study with similar CAPS scores, while CAPS scores after experimental sessions were lower for people in the MDMA condition through 2 months after the second experimental session (Placebo=59.1±28.9 versus MDMA, 25.4±23.95). Placebo subject scores dropped 20.5 points 2 months after the second experimental session while MDMA subject CAPS scores dropped 53.3 points, or a 26% drop in PTSD symptoms for controls versus a 68% drop in PTSD symptoms for MDMA subjects.

The second study of MDMA-assisted psychotherapy (MP-2) found results similar to the MP-1 study, but improvement after three blinded experimental sessions with 125 mg MDMA was numerically but not statistically superior to the 25 mg MDMA comparator dose [43]. CAPS

scores declined over time for the eight subjects given 125 mg MDMA (baseline mean=66.4±13.6 versus 3 weeks after the third experimental session mean=50.7±19.7), indicating a drop of 15.7 points, or a 23.5% decrease in scores. On the other hand, CAPS scores increased slightly over time for the four subjects given comparator dose (baseline mean=63.2±7.9 versus 3 weeks after the third experimental session mean=66.5±7.5), indicating an increase of 2.3 points, or a 5.2% increase in CAPS scores.

Table 29 and Table 30 below show pooled mean Global CAPS Scores for completed (MP-1, MP-2) and ongoing sponsor-supported studies (MP-4, MP-8, MP-9, MP-12). Since data collection is still in progress, formal analyses have yet to be executed, but data trends appear similar to published reports, with a medium to large effect size of active dose MDMA-assisted psychotherapy depending on number of experimental sessions completed. Table 29 below depicts mean Global CAPS scores for each condition at Baseline, 1 to 2 months after the second experimental session (Primary Endpoint), and 1 to 2 months after the third experimental session (End of Stage 1). Placebo and comparator groups cross over to Stage 2 after the Primary Endpoint, therefore CAPS is not administered at the End of Stage 1 for these groups. Active dose groups (100 mg and 125 mg) do not crossover, hence no data for Stage 2 endpoints. Long-term follow-up data collection is ongoing.

**Table 29: Mean Global CAPS Scores in Stage 1 of Sponsor-Supported Studies of MDMA-Assisted Psychotherapy for PTSD as of 01 October 2015**

Dose	Baseline Mean (SD) N	Primary Endpoint Mean (SD) N	End of Stage 1 Mean (SD) N
0 mg	83.6 (21.11) N=10	62.9 (27.04) N=10	---
25 mg	70.4 (10.01) N=8	61.2 (8.18) N=6	66.5 (7.55) N=4
30 mg	87.4 (14.12) N=7	73.5 (24.58) N=6	62.7 (36.12) N=3
40 mg	91.0 (17.89) N=7	80.6 (18.81) N=5	---
75 mg	82.4 (17.32) N=7	24.0 (18.79) N=6	18.5 (9.19) N=2
100 mg	94.4 (20.17) N=9	71.0 (30.85) N=7	40.9 (20.92) N=7
125 mg	84.13 (19.01) N=56	46.0 (31.46) N=53	42.4 (27.21) N=34

Across studies, CAPS scores are downward trending at the primary endpoint after two experimental sessions of MDMA-assisted psychotherapy. Formal pooled analyses to determine statistical significance have not been conducted as data collection is ongoing. Primary endpoint results after active doses of 75 mg to 125 mg initial dose, with an optional supplemental half-dose administered 1.5 to 2.5 hours later, appear lower than placebo or comparator dose results after two experimental sessions. Two-month follow-up results at the End of Stage 1 after a blinded or open-label third experimental session demonstrate signals of efficacy that should be further explored in a blinded three session treatment model of MDMA-assisted psychotherapy.

**Table 30: Mean Global CAPS Scores in Stage 2 and Long-term Follow-up of Sponsor-Supported Studies of MDMA-Assisted Psychotherapy for PTSD as of 01 October 2015**

Condition Stage 1/Stage 2	Last Stage 1 Observation Mean (SD) N	Secondary Endpoint Mean (SD) N	End of Stage 2 Mean (SD) N	12-month Follow-up Mean (SD) N
0 mg/125 mg	62.9 (27.04) N=10	33.9 (12.8) N=7	33.6 (18.6) N=5	18.7 (7.6) N=6
25 mg/125 mg	61.2 (8.18) N=6	42.5 (25.3) N=4	36.8 (13.6) N=4	31.5 (19.2) N=4
30 mg/125 mg	73.5 (24.58) N=6	46.5 (20.5) N=6	46.2 (30.5) N=6	59.0 (42.6) N=5
40 mg/125 mg	80.6 (18.81) N=5	38.6 (29.2) N=5	35.2 (31.1) N=5	14.0 (19.8) N=2
75 mg/125 mg	24.0 (18.79) N=6	22.3 (18.9) N=6	22.2 (20.5) N=5	26.8 (21.2) N=5
100 mg	40.9 (20.92) N=7	---	---	37.0 N=1
125 mg	42.4 (27.21) N=34	---	---	34.6 (28.1) N=30

Across studies, CAPS scores are also downward trending at the secondary endpoint after two open-label experimental sessions of MDMA-assisted psychotherapy and are consistent with Stage 1 results. Secondary endpoint results in the crossover set receiving an active dose of 125 mg MDMA after receiving comparator dose or placebo in Stage 1 are in range with subjects receiving 100 mg or 125 mg in Stage 1. Comparison between the 75 mg MDMA results in Stage 1 and the Stage 2 results suggest that this dose is also active and receiving additional 125 mg MDMA sessions does not lead to further improvement in this small sample. Formal analyses to determine statistical significance within-subjects have not been conducted as data collection is ongoing. Twelve-month follow-up results after all subjects have received active dose MDMA in either Stage 1 or Stage 2 suggest that the integration process may continue and lead to further improvement of PTSD symptoms in some subjects.

#### 5.4.2 Social Anxiety in Autistic Adults

The primary outcome measure for the study of social anxiety in people on the autism spectrum is the Liebowitz Social Anxiety Scale (LSAS). This observer-blind measure is an established clinician-administered measure of social anxiety, assessing fear and avoidance in different situations. The LSAS consists of 24 items, with each item rated on a four-point scale (from 0 to 3), with subscales for performance fear, performance avoidance, social fear, and social avoidance. The study is ongoing and efficacy findings will not be presented in this version of the IB.

Data is being collected on the effects of two sessions of MDMA-assisted therapy in people on the autism spectrum with social anxiety symptoms. The study is still blinded; therefore, efficacy data is not presented.

#### 5.4.3 Anxiety Associated with Life-Threatening Illness

MAPS is studying a new indication, the effects of MDMA-assisted psychotherapy on people experiencing anxiety as they face of a potentially life-threatening illness. No data is available at this time.

## 6.0 Summary of Data and Guidance for the Investigator

MDMA is a psychoactive compound that affects mood, perception, and increases prosocial feelings. The sponsor is investigating use of this compound as an adjunct to psychotherapy for treating PTSD, social anxiety in people on the autism spectrum, and anxiety related to a life-threatening illness. Researchers with and without sponsor support have conducted *in vitro* and *in vivo* non-clinical and clinical studies with MDMA, and additional clinical trials are ongoing. At this time, MDMA is listed as a Schedule I controlled substance in the U.S. and is not permitted for medical use outside of research settings. Psychotherapists in the U.S. began to use MDMA as an adjunct to psychotherapy in the mid to late 1970s, and narrative accounts describe therapeutic use prior to its scheduling. MDMA was administered to thousands of people in a therapeutic setting prior to scheduling, and has been administered to approximately 1180 people in controlled research settings as of 01 October 2015. These studies have demonstrated that MDMA can be safely administered to people with PTSD in a controlled clinical setting.

In comparison to anxiolytics, antidepressants, and atypical antipsychotics, MDMA does not require steady state levels in the blood to function as a catalyst to psychotherapy with rapid onset in some subjects. A limited number of exposures to MDMA, spaced approximately 1 month apart at moderate doses, are sufficient to obtain therapeutic outcomes. This intermittent dosing mitigates AE frequency and improves the risk/benefit ratio of MDMA, which may provide a significant advantage over medications that require daily dosing. Based on the current state of scientific knowledge and the risk/benefit profile of therapeutic doses of MDMA, the sponsor concludes that it appears favorable to pursue the research of MDMA as a medicine used as an adjunct to psychotherapy.

### 6.1 Pharmacology

The pharmacology of MDMA is complex as it activates multiple signaling cascades in the body. The formulation of the investigational product consists of a gelatin capsule consisting of racemic white crystalline MDMA, at doses ranging from 12.5 mg to 150 mg, compounded with alpha-lactose, and administered orally. Due to a wide range of responses to identical mg/kg dosing between individuals, possibly as a result of inconsistent relationship between body weight and pharmacodynamic activity, the sponsor's human trials use fixed doses between approximately 1 and 4 mg/kg (active fixed doses range from 75 mg to 225 mg cumulative with supplemental dosing, assuming a 60 kg individual) to achieve a more consistent response between subjects. In humans, onset of effects occurs approximately 30 to 60 minutes after administration, and peak effects occur 75 to 120 minutes after administration. Duration of effects lasts 3 to 6 hours, which extends to 6 to 8 hours with supplemental dosing.

The pharmacokinetics of MDMA in humans has been characterized using oral doses of up to 150 mg MDMA in humans. MDMA disposition in the body follows nonlinear pharmacokinetics. MDMA is metabolized in the liver by several enzymes. It is likely that active doses of MDMA saturate CYP2D6 function for an extended period, with function normalizing up to 10 days post-MDMA. The enzymes CYP1A2, COMT, and MAO may also be involved in the metabolism of MDMA. MDMA is metabolized by *N*-demethylation to MDA. The parent compound and MDA are further *O*-demethylated to HHMA and HHA, respectively. Both HHMA and HHA are subsequently *O*-methylated mainly to HMMA and HMA. These four metabolites, particularly HMMA and HMA, are known to be excreted in the urine as conjugated glucuronide or sulfate metabolites. The elimination half-life of active MDMA doses is 7 to 9 hours. This window should be considered when evaluating relationship of AEs to MDMA.

MDMA is a triple monoamine reuptake inhibitor, which concomitantly promotes carrier-mediated release, inhibits reuptake, and extends duration of serotonin, norepinephrine, and dopamine in the synaptic cleft to increase serotonergic, noradrenergic, and dopaminergic neurotransmission. MDMA appears to alter the conformation of the transporters, enabling monoamines to diffuse out of the neuron rather than being actively transported into the presynaptic neuron. MDMA was found to compete with monoamines for sites on the VMAT2, suggesting MDMA also promotes active release of monoamines from vesicular stores, in addition to inhibiting reuptake. MDMA extends the presence of monoamines in the synaptic cleft by inhibiting MAO-A, an enzyme that breaks down monoamines in the synapse. MDMA has self-limiting subjective and physiological effects. MDMA administration is contraindicated in subjects requiring MAOI medications. Fatalities have been reported after the combination of MAOIs and MDMA in Ecstasy users. Co-administration with an SSRI may eliminate or greatly attenuate the effects of MDMA, and these medications should be tapered in line with the investigator's clinical judgment and an approved study protocol.

MDMA has been shown to acutely decrease activity in the left amygdala and increase blood flow to the PFC in the human brain. The chief mechanism behind its therapeutic effects is likely to be serotonergic, along with some norepinephrine and to a minor extent dopamine-mediated effects. Indirect, but potentially significant effects of MDMA include the release of the hormones cortisol, oxytocin, prolactin, and AVP. MDMA likely stimulates secretion of oxytocin into peripheral blood via indirect activation of 5HT<sub>1A</sub>, 5HT<sub>2C</sub>, and 5HT<sub>4</sub> receptor subtypes, as well as AVP secretion via activation of 5HT<sub>2C</sub>, 5HT<sub>4</sub>, and 5HT<sub>7</sub> receptor subtypes. Both oxytocin and AVP are implicated in the widespread regulation of behavioral aspects of mood and also act on different target organs to modulate physiological functions in the body. Taken together, MDMA has been shown to have a diverse array of pharmacodynamic effects in animals and humans.

## 6.2 Toxicology

The toxicity of MDMA has been investigated in numerous animal and *in vitro* studies published in peer-reviewed journals. Intravenous MDMA doses that cause lethality in 50% of the cases, known as the LD50, are 97 mg/kg in mice, 49 mg/kg in rats, 14 to 18 mg/kg in dogs, and 22 mg/kg in monkeys. LD50 varies between different strains of the same animal species, across the sexes, housing conditions, environmental conditions, social interactions with co-habiting individuals, exercise levels, and water supply. Most preclinical toxicology data is derived from repeated dose studies. Preclinical researchers typically selected doses through use of interspecies scaling, a method of modeling human-equivalent doses in other species, however pharmacokinetic and pharmacodynamic data show this conversion is not appropriate for MDMA. As a result, most research in rodents and primates used doses of MDMA much higher than those consumed by humans, thus translation to human recreational and therapeutic use is limited. Many published epidemiological studies of Ecstasy effects in humans are also subject to the limitations in interpretation due to unknown purity, dose, and quantity of MDMA existing in Ecstasy tablets used in naturalistic settings.

Extensive preclinical toxicological studies report that high or repeated doses of MDMA can increase locomotor activity and signs of serotonin syndrome, which can damage serotonergic axons originating in the brainstem dorsal raphe nuclei, probably as a result of oxidative stress, and this damage is associated with decreases in serotonin production, serotonin metabolites, and SERT site densities. While these findings are consistent across studies, studies in low to moderate Ecstasy users do not report an increase in a biological marker of neuronal injury, and only one of three studies of this marker in humans detected it in heavy users. Retrospective studies in Ecstasy users have found contradictory effects on visual and verbal memory, planning and making decisions, and some types of visual processing. An uncontrolled prospective study of moderate

Ecstasy users failed to find changes in SERT sites or signs of neuronal injury; slight changes in cerebral blood flow in the dorsolateral PFC were found. In the same study, Ecstasy users showed less improvement on a memory task than non-users. Taken together, these findings suggest possible indications of cumulative toxicity in chronic high dose dosing regimens.

MDMA has not been demonstrated to be genotoxic. Consistent with this, despite very high doses of MDMA being tested in preclinical studies, none have reported carcinogenic effects. Risks posed to pregnant women by MDMA are not known. Two of three studies of Ecstasy users suggest that use of Ecstasy and other drugs during pregnancy may be associated with some abnormalities at birth, delays in mental and motor development, but not language or emotional development. Rodent fertility, reproductive, and developmental toxicity studies with MDMA have generally found no abnormalities in gestational duration, neonatal birth weights, or physical appearance when exposure occurs during organogenesis through lactation. However, one study of fertility and developmental toxicity in mice found evidence of toxicity at doses 5 mg/kg s.c. and above when exposure occurred in both genders of a breeding pair at some point between spermatogenesis/ovulation through closure of the hard palate. The results of several behavioral tests indicate that developmental MDMA exposure combined with adult exposure in rats may interfere with some aspects of learning, including visual-spatial memory, and time spent with a novel object. MDMA exposure *in utero* exacerbated hyperthermic response to a subsequent dose to MDMA. A study in neonatal rats suggests two distinct critical periods wherein repeated MDMA doses affected learning versus acoustic startle. In conclusion, MDMA might possess weak reproductive or developmental toxicity with a daily toxic chronic dosing regimen, in contrast to six or less exposures, spaced 1 month apart, tested in clinical trials. All sponsor-supported trials of MDMA exclude pregnant and lactating women, and women who are able to become pregnant must have a negative pregnancy screen before undergoing each experimental session and must agree to use birth control during the period of the protocol. If any subject becomes pregnant during study participation, the sponsor and clinical investigator will follow the pregnancy to outcome.

There have been a number of reports of morbidity and mortality in individuals who use Ecstasy (material represented as containing MDMA, as defined above) around the world in unsupervised and uncontrolled settings, usually involving poly-drug use. These events are relatively rare given the prevalence of Ecstasy use, estimated to be in the millions worldwide. The most common adverse effects in Ecstasy and poly-drug use include hyperthermia, psychiatric problems, hepatotoxicity secondary to hyperthermia, and hyponatremia (see [Section 4.4 Toxicology in Animals and Epidemiological Settings](#) and [4.5 Serious Reports, Morbidity, and Mortality in Epidemiological Settings](#)). Fatal dysrhythmias have been reported following heavy MDMA use, resulting in ventricular fibrillation and asystole. Individuals with underlying cardiac and/or pulmonary disease and preexisting conditions such as Wolff-Parkinson-White syndrome are especially at risk for heart failure and fatal arrhythmias when using MDMA. Set and setting likely play a role in the development of some Ecstasy-related adverse reports, such as vigorous exercise, lack of attention to somatic cues, and too little or too much hydration combined with pharmacological action on AVP, resulting in hyperthermia or hyponatremia. Even if ambient temperature does less to moderate the effects of MDMA on body temperature than originally believed based on animal studies, other environmental and behavioral factors, as those related to vigorous exercise, may be involved. Overall, the risks of serious reports appear to be minimal in controlled settings with adequate screening according to eligibility criteria defined in study protocols. None of these events have occurred within the context of human clinical studies with MDMA, likely due to careful screening for pre-existing risk factors and limited exposure in a controlled clinical setting.



### 6.3 Physiological Effects

MDMA is responsible for a series of dose dependent physiological effects due to enhanced neurochemical release of serotonin, norepinephrine, and dopamine, and for indirect effects on hormone secretion, including oxytocin and AVP, which act on different target organs to modulate physiological functions in the body. Active doses of MDMA (75 mg to 150 mg), alone or followed by a supplemental half-dose 1.5 to 2.5 hours later, are expected to produce statistically significant but transient, self-limited increases in blood pressure, heart rate, and body temperature that are likely to be well tolerated by healthy individuals. The elevation of blood pressure and increased heart rate produced by MDMA, like that produced by other sympathomimetic drugs, can lead to additional complications in people with pre-existing medical conditions that increase risk. In combination with clinical signs and symptoms, elevations in pulse and blood pressure can also lead to cardiac events, such as arrhythmias. No clinical studies have reported clinically important changes in physiological parameters.

Subjects enrolled in controlled Phase 1 single dose MDMA trials conducted without sponsor support had elevations above a pre-determined cut-off of at least 140/90 mmHg (approximately 5% per trial). All subjects in a subsequent trial in a separate sample given a regimen of 50 mg followed by 100 mg 2 hours later had blood pressure elevations above 140/90 mmHg. Based on the literature, effects of the initial dose of MDMA on blood pressure and heart rate are expected to have a linear dose-response relationship, and the supplemental dose may have an effect on SBP elevation. In sponsor-supported studies, SBP above 160 mmHg was detected in 27% of experimental sessions where MDMA was administered at any dose, and in 35% of subjects in sponsor-supported trials overall. The majority of these instances occurred with the 125 mg MDMA dose group. Both peak and longest duration of blood pressure elevation were also observed in the 125 mg MDMA group. Maximum duration of SBP above 160 mmHg was 6 hours in two subjects with peak values of 172 and 174, respectively. MDMA doses of 40 mg and greater were associated with SBP above 160 mmHg, supporting a dose dependent effect of MDMA on blood pressure. DBP above 110 mmHg was observed in only 5% of experimental sessions with MDMA at any dose in 7% of subjects. The majority of these instances occurred with the 125 mg MDMA dose. Maximum duration of DBP above 110 mmHg was 5 hours. Heart rate above 110 bpm was detected in 31% experimental sessions where MDMA was administered at any dose, in 39% of subjects in sponsor-supported trials. Both peak and maximum duration above 110 bpm were observed in 125 mg MDMA sessions. The highest pulse observed was 160 bpm for 1 hour. Maximum duration above 110 bpm was 9.5 hours. A comparison of subjects receiving the supplemental dose to those who only received the initial dose in MP-1 indicate that the supplemental dose did not cause further elevation in blood pressure and heart rate beyond the initial dose.

Candidates with controlled hypertension are excluded from participation in all but one of sponsor-supported studies to limit cardiovascular risk during treatments. In MP-8, the only study that did enroll a sub-group of subjects with controlled hypertension, SBP above 160 mmHg was detected in 75% (3 of 4) of subjects and 67% (8 of 12) of experimental sessions where MDMA was administered to this sub-group. The prevalence of these elevations appears higher in this sub-group than the overall sample, although the prevalence could decrease in a larger group. Pre-drug SBP was typically higher in this sub-group, and peak SBP of these subjects was typically at the upper end of the range of the overall sample. Final SBP readings remained 11 to 14 mmHg higher than pre-drug SBP readings in one of three subjects. The single subject with extended duration of SBP elevation had a medical history of both hypertension and hyperlipidemia. The same subject had DBP above 110 mmHg in each experimental session, suggesting that pre-existing cardiovascular risk factors beyond hypertension itself may be associated with further elevations in

blood pressure, though a larger sample would be needed to establish this. None of the subjects with controlled hypertension experienced AEs of the cardiovascular system.

Literature on epidemiological studies suggest a relationship between Ecstasy dose and likelihood of hyperthermia. Hyperthermia has occurred in people using Ecstasy in unsupervised and non-medical conditions, and though rare, is one of the most frequently reported serious adverse reports occurring in Ecstasy users. Environmental and behavioral factors, as well as thyroid dysregulation, may contribute to case reports and preclinical findings of hyperthermia. Findings from previous Phase 1 trials indicate that MDMA administered in a controlled setting produces a statistically but not clinically significant increase in body temperature (mean elevation of 0.6°C). The supplemental dose may limit elevations in body temperature, since it inhibits metabolism of MDMA to its bioactive metabolite MDA. MDA levels have been demonstrated to correlate with elevation in temperature in rodents. Unlike rodents, ambient temperature does not effect elevation in core temperature in humans. Controlled clinical settings have been sufficient to manage body temperature in humans.

Body temperature greater than 1°C above baseline was detected in 33% of experimental sessions in which MDMA was administered at any dose, in 46% of subjects in sponsor-supported trials, with most of these cases observed in sessions with 125 mg MDMA. In contrast, in 7% of experimental sessions in which inactive placebo was administered, and in 14% of subjects receiving inactive placebo, elevation of body temperature above cut-off was observed. Both peak and longest duration of body temperature elevation were observed in the 125 mg MDMA group. Maximum peak in all sessions was 38.7°C lasting 3 hours, and maximum duration of elevation in all sessions was 9.2 hours, in separate subjects. Vital signs in sponsor-supported Phase 2 studies presented above suggest a dose-dependent action on SBP and pulse, which is consistent with the literature on healthy volunteers. Body temperature and DBP do not appear to be strongly related to MDMA dose. No subjects receiving MDMA in sponsor-supported clinical trials have required any clinical interventions for elevated vital signs, as all values returned to normal as the effects of MDMA diminish.

### **6.3.1 Immunological Effects**

Humans exhibit transient immunological changes after a dose of 100 mg, including reduced numbers of CD4 cells, increased numbers of NK cells, and an increase in levels of immunosuppressive and anti-inflammatory cytokines compared with levels of pro-inflammatory and immunostimulating cytokines. In several respects, these effects are similar to those that occur with other psychoactive substances, so are not unique to MDMA. Immunological effects last for approximately 24 hours after administration, and most arise indirectly from serotonin release. The significance of these immunological effects remains unclear. Previous reports did not show increases in infections after MDMA and data from the study of MDMA-assisted psychotherapy has reported only instances of infection (upper respiratory or urinary tract) within 7 days of MDMA administration. Based on results from trials conducted by the sponsor, the impact of these effects is expected to be modest. The investigators may exclude subjects that might face additional risks from immunosuppression.

### **6.3.2 Hepatic Effects**

Phase 1 studies conducted outside of sponsor support involving administration of MDMA to healthy volunteers have not published any results of liver function after MDMA administration. There have been no reported adverse effects on the liver from these studies. The first two sponsor-supported Phase 2 studies (MP-1, MP-2) assessed liver function after completion of two or three blinded experimental sessions. No clinically or statistically significant changes in liver

function occurred in MP-1. Values for laboratory tests were within the normal range in MP-1. No AEs related to liver function have been reported in subsequent sponsor-supported studies. Only two subjects in the MP-2 study reported two clinically significant hepatic abnormalities, with one likely due to hereditary factors and the other indicating inflammation in a subject with a medical history of breast cancer 3 months after the last administration of MDMA as an AE unrelated to the study drug.

#### **6.4 Suicidal Ideation, Behavior, and Depression**

There is high incidence of suicidal ideation and behavior in populations of people with PTSD, especially those suffering from chronic, treatment resistant PTSD. In order to determine if suicidal ideation and behavior worsens or improves after treatment in MAPS-sponsored trials, the C-SSRS is administered repeatedly throughout the study. Due to the nature of the therapeutic method, wherein a person may re-experience emotions associated with the traumatic event in order to reprocess the memory in a new, therapeutic way, thoughts of ending one's life may surface during this process. However, evidence from ongoing studies indicates that these thoughts are most often transient, returning to baseline, or even improving during the acute period following MDMA treatment. C-SSRS scores have escalated during the preparatory sessions (before any drug administration), which is thought to be a result of preparatory discussion of traumatic experiences, and/or of subjects tapering off long-prescribed medications, such as SSRIs and benzodiazepines. Withdrawal of these drugs is known induce suicidal ideation or behavior in some people. During both non-drug and MDMA-assisted psychotherapy sessions, subjects are asked to think about and discuss their experiences, thoughts, and emotions related to their condition. They may experience intense emotional responses to recalling and speaking about this material. As MDMA is only administered in combination with psychotherapy, the distress associated with psychotherapy is unavoidable, and is considered a necessary part of the therapeutic process that requires proper facilitation and support from therapists.

Overall the incidence of serious suicidal ideation or behavior in sponsor-supported studies is low, occurring in only a few subjects post-MDMA treatment, and returning to non-life-threatening scores while subjects are closely monitored. Given that people suffering from severe PTSD are known to experience suicidal ideation and behavior, it is difficult to identify a single cause of the increase in suicidal thinking or behavior (i.e. exacerbation of PTSD symptoms related to medication withdrawal or to the psychotherapeutic process, or from MDMA effects). A large percentage of people enrolled in the studies reported suicidal ideation and behavior during sometime in their lives prior to study enrollment, which may reflect a manifestation of PTSD or co-morbid affective disorders. When positive serious ideation or behavior occurred after study enrollment, the investigators made follow-up observations of C-SSRS to ensure subject safety, and tracked scores until they returned to non-serious levels. Only two incidences of suicidal ideation have been considered clinically significant and tracked as severe AEs, but they were reported during the long-term follow-up period and were not related to study drug.

#### **6.5 Adverse Events**

Overall, adverse effects of MDMA are modest and generally have not been associated with serious discomfort in healthy volunteers or in people with PTSD. Risks posed by sympathomimetic effects of MDMA treatments are addressed in MAPS' clinical trials by excluding people with pre-existing cardiovascular disease, cerebrovascular disease or uncontrolled hypertension and by monitoring blood pressure, body temperature, and pulse. Common reactions reported in clinical trials are transient and diminish as drug effects wane during the MDMA session and over the next 24 hours. Once the drug leaves the body, 3 to 4 days post-treatment, most reactions diminish. Reactions are monitored daily for 1 week after each

treatment and followed until resolution. The most common acute reactions at any severity include muscle tension in the jaw, exacerbation of anxiety, decreased appetite, muscle tension, nausea, and feeling cold. Headache and fatigue are commonly reported across MDMA and placebo groups, and are likely to be background events. During the week after each experimental session, the most commonly reported reactions at any severity were anxiety, fatigue, insomnia, depressed mood, hypersomnia, difficulty concentrating, decreased appetite, and dizziness in the active dose MDMA groups across studies, with PTSD studies overrepresented. Of these reactions, only decreased appetite and dizziness were appreciably elevated above the placebo group, and the remaining reactions are likely to be background events. Severe unexpected AEs included abdominal cramps, panic attacks, and the following reactions lasting longer than 7 days: anxiety, headache, low mood, rumination, and restlessness, all reported in studies of MDMA-assisted psychotherapy for PTSD. All subjects fully recovered from these events.

Unexpected and expected SAEs related to administration of MDMA in MAPS-sponsored clinical trials have been rare and none have been life threatening. One probably drug-related expected SAE has occurred to date in this clinical development program. This event was an increase in frequency of ventricular extrasystoles experienced during treatment with 125 mg MDMA, which resolved with full recovery to baseline after the study drug's effects ceased. The subject was hospitalized for observation and recovered fully after the event, with no cardiac damage. Excluding people with cerebrovascular or cardiovascular disease will reduce the likelihood of risks arising from the cardiovascular effects of MDMA.

## **6.6 Risk Mitigation in MDMA-Assisted Clinical Trials**

Investigators must establish subject eligibility prior to enrollment in trials with MDMA, with eligibility established through medical history, physical examination, vital signs, clinical laboratory tests, EKG, psychiatric interview, and assessment of relevant psychiatric symptoms. Additional procedures may be used as indicated, such as exercise tests and carotid ultrasound imaging. If the study is investigating use of MDMA in people with a specific psychiatric condition, then the investigators must also determine whether an individual has the condition and that specified exclusion criteria are absent.

MDMA-assisted psychotherapy clinical trials use questionnaire-based measures and clinical interviews that can cause testing fatigue and/or emotional reactions stemming from discussing trauma or other psychological stressors. Investigators should be experienced in treating the condition under investigation and they should seek to minimize testing fatigue and emotional stress during screening and participation in the study. Subjects enrolled in studies of MDMA-assisted psychotherapy should be prepared to engage in processing their trauma, which requires proper facilitation and support from study therapists. MDMA-assisted psychotherapy will always be performed in controlled clinical settings to mitigate risk. It is best to ensure that the controlled setting for treatments with MDMA-assisted psychotherapy has the capacity to control ambient temperature for subject comfort, though there is no evidence that this will significantly influence or is needed for control of core body temperature. Cardiovascular risk is primarily mitigated through rigorous screening to exclude subjects with uncontrolled cardiovascular risk. During experimental sessions, therapists should monitor for clinical signs and symptoms (severe headache, confusion or focal neurologic signs, vision problems, chest pain, difficulty breathing, or palpitations) and add more frequent vitals measurements only if clinically indicated. Investigators conducting trials of MDMA should be prepared to treat elevated blood pressure with medications if needed, and either to provide appropriate care related to these effects or to transport individuals to an emergency department, if necessary.

Discontinuing pre-study medications and the acute/sub-acute effects of MDMA-assisted psychotherapy can produce shifts in mood and activation, which may transiently increase likelihood of suicidal ideation or behavior. In addition, during treatment of subjects with prevalent lifetime history of suicidal ideation, the active dose of MDMA, which catalyzes the therapeutic process, can be associated with suicidal ideation as a result of processing trauma. To mitigate risk, subjects are kept under continuous clinical observation during experimental sessions. Experimental sessions are followed by an overnight residential stay at the study site to allow the integration process to begin, followed by an integrative psychotherapy session on the following day, daily phone contact for 1 week, with channels of easy access to the treating therapists maintained during the studies. The need for additional support in these studies is continually assessed with the General Well Being and AE monitoring. Due to the psychotherapeutic setting in which MDMA is provided in these studies, exacerbations of symptoms often appear to be related to the therapeutic process rather than directly to the MDMA itself. When assessing potential AEs, investigators should consider baseline severity of conditions and symptoms, therapeutic process, and potential relationship to drug administration throughout the study.

In sponsor-supported studies, 18% of people across active dose and comparator dose groups experienced periods of increased anxiety (2% severe) and 3% experienced panic attacks, all in the active dose MDMA group (2% severe). Psychological distress may arise at any time during an MDMA-assisted psychotherapy session from the time of first drug effects until effects have dissipated approximately 3 to 5 hours after administration. Anxiety or distress during the session may last for as little as 15 minutes or for as long as 5 hours or more and may be related to the therapeutic process itself. In addition, psychological distress could arise following an MDMA session as a result of subjects having difficulty integrating their experience after the effects of MDMA have subsided. In previous Phase 1 and Phase 2 studies, these symptoms have been modest and self-limiting, and have responded well to reassurance from the investigator, with occasional use of benzodiazepines for anxiety after the experimental session. In clinical trials of PTSD treatment, subjects are informed that experimental sessions are intended to include periods of confronting and working through traumatic experiences. Hence signs of psychological distress, anxiety, or other unpleasant psychological reactions are to be expected and may be considered an element of the psychotherapeutic process. In Phase 1 trials with normal healthy volunteers, mild anxiety and depressed mood were reported by some subjects 1 to 3 days after MDMA administration. It is not known whether these reactions resulted from direct effects of MDMA, or from psychological content that may have been accessed during the MDMA experience.

The potential for destabilizing psychological distress can be minimized by:

- Exclusion of people who might be more vulnerable to psychological destabilization if tapered off other psychiatric drugs, such as people diagnosed with bipolar affective disorder-1 or those with psychotic disorders.
- Preparatory non-drug psychotherapy sessions before the experimental session
- An atmosphere of trust during the experimental session
- Close monitoring of the subject
- Daily contact with subjects for the period of 1 week after the experimental session, and availability of therapists at other times as needed.
- Non-drug integrative psychotherapy sessions
- Having subjects remain at the study site for the night of each experimental session to provide an optimal opportunity for rest and reflection following MDMA-assisted sessions, as part of the integration process.
- Availability of qualified personnel, such as a trained attendant during the overnight stay to support rest and integration of the experience.

Every effort is made to help subjects resolve difficult symptoms and to arrive at a more comfortable and relaxed state by the conclusion of the session. Such efforts include empathic listening on the part of the investigators and affect management techniques, such as diaphragmatic breathing by subjects.

At the end of any experimental session, if the subject is still severely agitated or experiencing any other severe psychological distress, the following measures should be taken:

1. If the subject is anxious, agitated, and/or in danger of any self-harm or is suicidal at the end of the MDMA session, the investigators are available to remain with the subject for at least two more hours. During this time, the investigators use affect management techniques reviewed during the introductory sessions and talk with the subject to help him or her gain cognitive perspective about their experience. If this situation should occur during an integrative therapy session, the same approach should be used, and at least one of the investigators will remain available to stay with the subject for at least two additional hours.
2. If a subject remains severely anxious, agitated or in danger of self-harm or suicide, or is otherwise psychologically unstable at the end of this 2-hour stabilization period, the clinical investigator decides between the following options:
  - a. A psychiatric nurse, therapeutic assistant, or therapist will stay with the subject until the time of his or her appointment with investigators the next day. The investigators then meet with the subject daily until the period of destabilization has passed.
  - b. If a subject experiences severe, persisting emotional distress, such as panic attacks, severe generalized anxiety, or insomnia following an MDMA session, the investigator may prescribe a drug with a short half-life as a "rescue medication." Investigators should not prescribe an SSRI, SNRI, or MAOI in this context unless it is determined that such treatment is clinically necessary and the subject will be terminated from study participation. Residual symptoms are addressed during the frequent follow-up psychotherapy visits with the investigators.
  - c. Hospitalization for stabilization. If a subject should become psychotic, or if for any reason the investigators deem it necessary for safety, arrangements are made to stabilize and transfer him or her to the study site inpatient unit or the nearest appropriate inpatient psychiatric facility.

Subjects hospitalized after a severe panic reaction or other adverse psychological reaction would be suspended from further participation in the trial until after recovery or stabilization, at which time the investigator would carefully evaluate the subject's emotional status and decide whether or not the subject may continue the study. For those subjects engaged in an ongoing therapeutic relationship with a psychotherapist or psychiatrist, the subject's outside therapists are involved in the management of any psychiatric complications.

## **6.7 Abuse Potential**

Despite its classification as a Schedule I drug, an examination of findings in humans and animals suggests that MDMA possesses moderate abuse potential that is higher than that reported for "classic hallucinogens" like psilocybin, but lower than that reported for psychostimulants, such as cocaine or methamphetamine. Studies assessing prevalence of problematic Ecstasy use or dependence suggest that a small percentage of individuals, especially those with prior psychological difficulties, may develop problematic Ecstasy use or dependence. In two published studies of MDMA-assisted psychotherapy for people with PTSD, only one of 32 subjects reported



using Ecstasy subsequent to study participation, and several subjects volunteered that they would not seek out Ecstasy outside of a psychotherapeutic setting. Diversion is not an issue for sponsor-supported studies because MDMA will only be administered under the supervision of the clinical investigator and no take-home doses are permitted. MDMA is handled following all regulations pertaining to the handling and dispensing of controlled substances within research studies.

## **7.0 Conclusion**

Based on the current state of scientific knowledge, the risk for subjects meeting criteria for clinical studies who are exposed to MDMA at the single intermittent dosing schedule used in sponsor-supported studies appears to be low. The overall rates of AEs and reactions across phase 2 studies are low and the reactions and AEs are self-limiting. A number of the AEs and expected reactions reported in the studies are likely related to background events representing the underlying illness being treated, or the expected result of psychotherapy addressing traumatic experiences.

Future studies conducted by the sponsor are intended to further develop the safety profile of MDMA in the PTSD subject population. In addition, the sponsor is examining the use of MDMA-assisted psychotherapy in the treatment of anxiety, including social anxiety in people on the autistic spectrum and anxiety resulting from a life-threatening illness. MDMA-assisted psychotherapy appears to be a promising treatment method for chronic PTSD. More clinical trials in larger subject populations are warranted. It is hoped that MDMA, with its unique pharmacological mechanisms combined with a novel mode of administration in conjunction with psychotherapy, can improve upon first line PTSD and anxiety treatments in terms of side effect profiles, efficacy and duration of effect.

## 8.0 References

### References

1. Beck, J. and M. Rosenbaum, *In Pursuit of Ecstasy: The MDMA Experience*. 1994, Albany, NY: SUNY Press.
2. Solowij, N., W. Hall, and N. Lee, *Recreational MDMA use in Sydney: a profile of 'Ecstasy' users and their experiences with the drug*. *Br J Addict*, 1992. **87**(8): p. 1161-72.
3. Carlson, R.G., et al., *Drug use practices among MDMA/ecstasy users in Ohio: a latent class analysis*. *Drug Alcohol Depend*, 2005. **79**(2): p. 167-79.
4. Sumnall, H.R., J.C. Cole, and L. Jerome, *The varieties of ecstatic experience: an exploration of the subjective experiences of ecstasy*. *J Psychopharmacol*, 2006. **20**(5): p. 670-82.
5. Cole, J.C. and H.R. Sumnall, *Altered states: the clinical effects of Ecstasy*. *Pharmacol Ther*, 2003. **98**(1): p. 35-58.
6. Shulgin, A.T., *The background and chemistry of MDMA*. *J Psychoactive Drugs*, 1986. **18**(4): p. 291-304.
7. Kolbrich, E.A., et al., *Physiological and subjective responses to controlled oral 3,4-methylenedioxymethamphetamine administration*. *J Clin Psychopharmacol*, 2008. **28**(4): p. 432-40.
8. Cami, J., et al., *Human pharmacology of 3,4-methylenedioxymethamphetamine ("ecstasy"): psychomotor performance and subjective effects*. *J Clin Psychopharmacol*, 2000. **20**(4): p. 455-66.
9. Mas, M., et al., *Cardiovascular and neuroendocrine effects and pharmacokinetics of 3, 4-methylenedioxymethamphetamine in humans*. *J Pharmacol Exp Ther*, 1999. **290**(1): p. 136-45.
10. Liechti, M.E., A. Gamma, and F.X. Vollenweider, *Gender differences in the subjective effects of MDMA*. *Psychopharmacology (Berl)*, 2001. **154**(2): p. 161-8.
11. Tancer, M. and C.E. Johanson, *Reinforcing, subjective, and physiological effects of MDMA in humans: a comparison with d-amphetamine and mCPP*. *Drug Alcohol Depend*, 2003. **72**(1): p. 33-44.
12. Harris, D.S., et al., *Subjective and hormonal effects of 3,4-methylenedioxymethamphetamine (MDMA) in humans*. *Psychopharmacology (Berl)*, 2002. **162**(4): p. 396-405.
13. Vollenweider, F.X., et al., *Psychological and cardiovascular effects and short-term sequelae of MDMA ("ecstasy") in MDMA-naive healthy volunteers*. *Neuropsychopharmacology*, 1998. **19**(4): p. 241-51.
14. de la Torre, R., et al., *Non-linear pharmacokinetics of MDMA ('ecstasy') in humans*. *Br J Clin Pharmacol*, 2000. **49**(2): p. 104-9.
15. Marks, D.M., C.U. Pae, and A.A. Patkar, *Triple reuptake inhibitors: a premise and promise*. *Psychiatry Investig*, 2008. **5**(3): p. 142-7.
16. Millan, M.J., *Dual- and triple-acting agents for treating core and co-morbid symptoms of major depression: novel concepts, new drugs*. *Neurotherapeutics*, 2009. **6**(1): p. 53-77.
17. Bonkale, W.L. and M.C. Austin, *3,4-Methylenedioxymethamphetamine induces differential regulation of tryptophan hydroxylase 2 protein and mRNA levels in the rat dorsal raphe nucleus*. *Neuroscience*, 2008. **155**(1): p. 270-6.
18. Schenk, S., et al., *Effects of repeated exposure to MDMA on 5HT1a autoreceptor function: behavioral and neurochemical responses to 8-OHDPAT*. *Psychopharmacology (Berl)*, 2013. **227**(2): p. 355-61.
19. Hysek, C.M., et al., *MDMA enhances emotional empathy and prosocial behavior*. *Soc Cogn Affect Neurosci*, 2014. **9**(11): p. 1645-52.
20. Adori, C., et al., *Recovery and aging of serotonergic fibers after single and intermittent MDMA treatment in Dark Agouti rat*. *J Comp Neurol*, 2011. **519**(12): p. 2353-78.

21. Han, D.D. and H.H. Gu, *Comparison of the monoamine transporters from human and mouse in their sensitivities to psychostimulant drugs*. BMC Pharmacol, 2006. **6**: p. 6.
22. Liechti, M.E. and F.X. Vollenweider, *Which neuroreceptors mediate the subjective effects of MDMA in humans? A summary of mechanistic studies*. Hum Psychopharmacol, 2001. **16**(8): p. 589-598.
23. Murnane, K.S., et al., *Endocrine and neurochemical effects of 3,4-methylenedioxymethamphetamine and its stereoisomers in rhesus monkeys*. J Pharmacol Exp Ther, 2010. **334**(2): p. 642-50.
24. Setola, V., et al., *3,4-methylenedioxymethamphetamine (MDMA, "Ecstasy") induces fenfluramine-like proliferative actions on human cardiac valvular interstitial cells in vitro*. Mol Pharmacol, 2003. **63**(6): p. 1223-9.
25. Simmler, L.D., et al., *Monoamine transporter and receptor interaction profiles of a new series of designer cathinones*. Neuropharmacology, 2014. **79**: p. 152-60.
26. Bedi, G., et al., *Effects of MDMA on sociability and neural response to social threat and social reward*. Psychopharmacology (Berl), 2009. **207**(1): p. 73-83.
27. Carhart-Harris, R.L., et al., *The Effects of Acutely Administered 3,4-Methylenedioxymethamphetamine on Spontaneous Brain Function in Healthy Volunteers Measured with Arterial Spin Labeling and Blood Oxygen Level-Dependent Resting State Functional Connectivity*. Biol Psychiatry, 2015. **78**(8): p. 554-62.
28. Gamma, A., et al., *3,4-Methylenedioxymethamphetamine (MDMA) modulates cortical and limbic brain activity as measured by [H(2)(15)O]-PET in healthy humans*. Neuropsychopharmacology, 2000. **23**(4): p. 388-95.
29. Dumont, G.J., et al., *Increased oxytocin concentrations and prosocial feelings in humans after ecstasy (3,4-methylenedioxymethamphetamine) administration*. Soc Neurosci, 2009. **4**(4): p. 359-66.
30. Hysek, C.M., G. Domes, and M.E. Liechti, *MDMA enhances "mind reading" of positive emotions and impairs "mind reading" of negative emotions*. Psychopharmacology (Berl), 2012. **222**(2): p. 293-302.
31. Kirkpatrick, M.G., et al., *Plasma oxytocin concentrations following MDMA or intranasal oxytocin in humans*. Psychoneuroendocrinology, 2014. **46**: p. 23-31.
32. Wolff, K., et al., *Vasopressin and oxytocin secretion in response to the consumption of ecstasy in a clubbing population*. J Psychopharmacol, 2006. **20**(3): p. 400-10.
33. Fallon, J.K., et al., *Action of MDMA (ecstasy) and its metabolites on arginine vasopressin release*. Ann N Y Acad Sci, 2002. **965**: p. 399-409.
34. Schmid, Y., et al., *Differential effects of MDMA and methylphenidate on social cognition*. J Psychopharmacol, 2014. **28**(9): p. 847-56.
35. Bedi, G., D. Hyman, and H. de Wit, *Is ecstasy an "empathogen"? Effects of +/-3,4-methylenedioxymethamphetamine on prosocial feelings and identification of emotional states in others*. Biol Psychiatry, 2010. **68**(12): p. 1134-40.
36. Carhart-Harris, R.L., et al., *The effect of acutely administered MDMA on subjective and BOLD-fMRI responses to favourite and worst autobiographical memories*. Int J Neuropsychopharmacol, 2014. **17**(4): p. 527-40.
37. Bedi, G., et al., *A window into the intoxicated mind? Speech as an index of psychoactive drug effects*. Neuropsychopharmacology, 2014. **39**(10): p. 2340-8.
38. Wardle, M.C., M.G. Kirkpatrick, and H. de Wit, *'Ecstasy' as a social drug: MDMA preferentially affects responses to emotional stimuli with social content*. Soc Cogn Affect Neurosci, 2014. **9**(8): p. 1076-81.
39. Wardle, M.C. and H. de Wit, *MDMA alters emotional processing and facilitates positive social interaction*. Psychopharmacology (Berl), 2014. **231**(21): p. 4219-29.
40. Danforth, A., *Findings from a collective case study of the MDMA/Ecstasy experiences of adults on the autism spectrum: Transcripts from Psychedelic Science 2013*. MAPS Bulletin, 2013. **23**(3): p. 30-35.

41. Mithoefer, M.C., et al., *The safety and efficacy of {+/-}3,4-methylenedioxymethamphetamine-assisted psychotherapy in subjects with chronic, treatment-resistant posttraumatic stress disorder: the first randomized controlled pilot study.* J Psychopharmacol, 2011. **25**(4): p. 439-52.
42. Mithoefer, M.C., et al., *Durability of improvement in post-traumatic stress disorder symptoms and absence of harmful effects or drug dependency after 3,4-methylenedioxymethamphetamine-assisted psychotherapy: a prospective long-term follow-up study.* J Psychopharmacol, 2013. **27**(1): p. 28-39.
43. Oehen, P., et al., *A randomized, controlled pilot study of MDMA (+/- 3,4-Methylenedioxymethamphetamine)-assisted psychotherapy for treatment of resistant, chronic Post-Traumatic Stress Disorder (PTSD).* J Psychopharmacol, 2013. **27**(1): p. 40-52.
44. Downing, J., *The psychological and physiological effects of MDMA on normal volunteers.* J Psychoactive Drugs, 1986. **18**(4): p. 335-40.
45. Lester, S.J., et al., *Cardiovascular effects of 3,4-methylenedioxymethamphetamine. A double-blind, placebo-controlled trial.* Ann Intern Med, 2000. **133**(12): p. 969-73.
46. Kirkpatrick, M.G., et al., *A direct comparison of the behavioral and physiological effects of methamphetamine and 3,4-methylenedioxymethamphetamine (MDMA) in humans.* Psychopharmacology (Berl), 2012. **219**(1): p. 109-22.
47. Grob, C.S., et al., *Psychobiologic effects of 3,4-methylenedioxymethamphetamine in humans: methodological considerations and preliminary observations.* Behav Brain Res, 1996. **73**(1-2): p. 103-7.
48. Pacifici, R., et al., *[Immunomodulator properties of ecstasy (MDMA)].* Ann Ist Super Sanita, 2000. **36**(1): p. 69-75.
49. Baggott, M.J., *Preventing problems in Ecstasy users: reduce use to reduce harm.* J Psychoactive Drugs, 2002. **34**(2): p. 145-62.
50. Gore, S.M., *Fatal uncertainty: death-rate from use of ecstasy or heroin.* Lancet, 1999. **354**(9186): p. 1265-6.
51. Hall, A.P. and J.A. Henry, *Acute toxic effects of 'Ecstasy' (MDMA) and related compounds: overview of pathophysiology and clinical management.* Br J Anaesth, 2006. **96**(6): p. 678-85.
52. Liechti, M.E., I. Kunz, and H. Kupferschmidt, *Acute medical problems due to Ecstasy use. Case-series of emergency department visits.* Swiss Med Wkly, 2005. **135**(43-44): p. 652-7.
53. Rogers, G., et al., *The harmful health effects of recreational ecstasy: a systematic review of observational evidence.* Health Technol Assess, 2009. **13**(6): p. iii-iv, ix-xii, 1-315.
54. Rosenson, J., et al., *Patterns of ecstasy-associated hyponatremia in California.* Ann Emerg Med, 2007. **49**(2): p. 164-71, 171 e1.
55. Williams, H., et al., *"Saturday night fever": ecstasy related problems in a London accident and emergency department.* J Accid Emerg Med, 1998. **15**(5): p. 322-6.
56. Freudenmann, R.W., F. Oxler, and S. Bernschneider-Reif, *The origin of MDMA (ecstasy) revisited: the true story reconstructed from the original documents.* Addiction, 2006. **101**(9): p. 1241-5.
57. Shulgin, A. and A. Shulgin, *Pihkal: A Chemical Love Story.* 1st ed. 1991, Berkeley, CA: Transform Press. 1-978.
58. Anderson, G.M.d., et al., *Absolute configuration and psychotomimetic activity.* NIDA Res Monogr, 1978. **22**: p. 8-15.
59. Shulgin, A.T. and D.E. Nichols, *Characterization of three new psychotomimetics,* in *The Pharmacology of Hallucinogens*, R.C. Stillman and R.E. Willette, Editors. 1978, Pergamon: New York.

60. Rudnick, G. and S.C. Wall, *The molecular mechanism of "ecstasy" [3,4-methylenedioxy-methamphetamine (MDMA)]: serotonin transporters are targets for MDMA-induced serotonin release*. Proc Natl Acad Sci U S A, 1992. **89**(5): p. 1817-21.
61. O'Neil, M.J., *The Merck Index: An Encyclopedia of chemicals, drugs and biologicals*. Merck Research Laboratories, Merck and Co. Inc, Whitehouse station, New Jersey, 2006. **319**.
62. Greer, G. and R. Tolbert, *Subjective reports of the effects of MDMA in a clinical setting*. J Psychoactive Drugs, 1986. **18**(4): p. 319-27.
63. Kirkpatrick, M.G., et al., *Effects of MDMA and Intranasal oxytocin on social and emotional processing*. Neuropsychopharmacology, 2014. **39**(7): p. 1654-63.
64. Adamson, S., *Through the gateway of the heart: Accounts of experiences With MDMA and other empathogenic substances*. 1985, San Francisco CA: Four Trees Publications.
65. Grinspoon, L. and J.B. Bakalar, *Can drugs be used to enhance the psychotherapeutic process?* Am J Psychother, 1986. **40**(3): p. 393-404.
66. Nichols, D.E., *Differences between the mechanism of action of MDMA, MBDB, and the classic hallucinogens. Identification of a new therapeutic class: entactogens*. J Psychoactive Drugs, 1986. **18**(4): p. 305-13.
67. Greer, G. and R.J. Strassman, *Information on "Ecstasy"*. Am J Psychiatry, 1985. **142**(11): p. 1391.
68. Greer, G.R. and R. Tolbert, *A method of conducting therapeutic sessions with MDMA*. J Psychoactive Drugs, 1998. **30**(4): p. 371-379.
69. Metzner, R. and S. Adamson, *Using MDMA in healing, psychotherapy and spiritual practice*, in *Ecstasy, A Complete Guide: A Comprehensive Look at the Risks and Benefits of MDMA.*, J. Holland, Editor. 2001, Inner Traditions: Rochester VT. p. 182-207.
70. Greer, G. and R. Tolbert, *The therapeutic use of MDMA.*, in *Ecstasy: The clinical, pharmacological and neurotoxicological effects of the drug MDMA.*, S.J. Peroutka, Editor. 1990, Kluver Academic: Boston, MA. p. 21-35.
71. Carlson, R.G., et al., *MDMA/Ecstasy use among young people in Ohio: perceived risk and barriers to intervention*. J Psychoactive Drugs, 2004. **36**(2): p. 181-9.
72. Substance Abuse and Mental Health Services Administration, *Results from the 2013 National Survey on Drug Use and Health: Summary of National Findings , NSDUH Series H-48*. 2014, Substance Abuse and Mental Health Services Administration,: Rockville, MD.
73. Drugs, E.M.C.f. and D. Addiction, *European Drug Report 2014: Trends and Developments*. 2015.
74. Drug Enforcement Agency, *Scheduling of Controlled Substances: Scheduling of 3,4-methylenedioxymethamphetamine (MDMA) into Schedule I of the Controlled Substances Act: Remand*, DEA, Editor. 1988, Federal Register: Washington, DC. p. 5156-5160.
75. Lawn, J.C., *On the Matter of MDMA Scheduling*, in *Docket No. 84-48*, Drug\_Enforcement\_Administration, Editor. 1986: Washington, DC.
76. Seiden, L.S. and K.E. Sabol, *Methamphetamine and methylenedioxy-methamphetamine neurotoxicity: possible mechanisms of cell destruction*. NIDA Res Monogr, 1996. **163**: p. 251-76.
77. Baumann, M.H., X. Wang, and R.B. Rothman, *3,4-Methylenedioxy-methamphetamine (MDMA) neurotoxicity in rats: a reappraisal of past and present findings*. Psychopharmacology (Berl), 2007. **189**(4): p. 407-24.
78. Baumann, M.H., et al., *Effects of dose and route of administration on pharmacokinetics of (+ or -)-3,4-methylenedioxy-methamphetamine in the rat*. Drug Metab Dispos, 2009. **37**(11): p. 2163-70.
79. Mueller, F., et al., *Neuroimaging in moderate MDMA use: A systematic review*. Neurosci Biobehav Rev, 2015.

80. Henry, J.A., K.J. Jeffreys, and S. Dawling, *Toxicity and deaths from 3,4-methylenedioxymethamphetamine ("ecstasy")*. *Lancet*, 1992. **340**(8816): p. 384-7.
81. Cohen, R.S. and J. Cocores, *Neuropsychiatric manifestations following the use of 3,4-methylenedioxymethamphetamine (MDMA: "Ecstasy")*. *Prog Neuropsychopharmacol Biol Psychiatry*, 1997. **21**(4): p. 727-34.
82. Williamson, S., et al., *Adverse effects of stimulant drugs in a community sample of drug users*. *Drug Alcohol Depend*, 1997. **44**(2-3): p. 87-94.
83. Krystal, J.H., et al., *Chronic 3,4-methylenedioxymethamphetamine (MDMA) use: effects on mood and neuropsychological function?* *Am J Drug Alcohol Abuse*, 1992. **18**(3): p. 331-41.
84. McCann, U.D., et al., *Serotonin neurotoxicity after (+/-)3,4-methylenedioxymethamphetamine (MDMA; "Ecstasy"): a controlled study in humans*. *Neuropsychopharmacology*, 1994. **10**(2): p. 129-38.
85. McCann, U.D., et al., *Cognitive performance in (+/-) 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") users: a controlled study*. *Psychopharmacology (Berl)*, 1999. **143**(4): p. 417-25.
86. Parrott, A.C. and J. Lasky, *Ecstasy (MDMA) effects upon mood and cognition: before, during and after a Saturday night dance*. *Psychopharmacology (Berl)*, 1998. **139**(3): p. 261-8.
87. Ricaurte, G.A., et al., *Toxicodynamics and long-term toxicity of the recreational drug, 3,4-methylenedioxymethamphetamine (MDMA, 'Ecstasy')*. *Toxicol Lett*, 2000. **112-113**: p. 143-6.
88. Biezonski, D.K. and J.S. Meyer, *The Nature of 3, 4-Methylenedioxymethamphetamine (MDMA)-Induced Serotonergic Dysfunction: Evidence for and Against the Neurodegeneration Hypothesis*. *Curr Neuropharmacol*, 2011. **9**(1): p. 84-90.
89. Gouzoulis-Mayfrank, E. and J. Daumann, *The confounding problem of polydrug use in recreational ecstasy/MDMA users: a brief overview*. *J Psychopharmacol*, 2006. **20**(2): p. 188-93.
90. Bouso, J.C., *MDMA/PTSD research in Spain: An update*. *MAPS Bulletin*, 2003. **13**(1): p. 7-8.
91. Farre, M., et al., *Pharmacological interaction between 3,4-methylenedioxymethamphetamine (ecstasy) and paroxetine: pharmacological effects and pharmacokinetics*. *J Pharmacol Exp Ther*, 2007. **323**(3): p. 954-62.
92. van Wel, J.H., et al., *Effects of Acute MDMA Intoxication on Mood and Impulsivity: Role of the 5-HT(2) and 5-HT(1) Receptors*. *PLoS One*, 2012. **7**(7): p. e40187.
93. Lyon, R.A., R.A. Glennon, and M. Titeler, *3,4-Methylenedioxymethamphetamine (MDMA): stereoselective interactions at brain 5-HT1 and 5-HT2 receptors*. *Psychopharmacology (Berl)*, 1986. **88**(4): p. 525-6.
94. Fantegrossi, W.E., et al., *Discriminative stimulus effects of 3,4-methylenedioxymethamphetamine and its enantiomers in mice: pharmacokinetic considerations*. *J Pharmacol Exp Ther*, 2009. **329**(3): p. 1006-15.
95. Yarosh, H.L., et al., *MDMA-like behavioral effects of N-substituted piperazines in the mouse*. *Pharmacol Biochem Behav*, 2007. **88**(1): p. 18-27.
96. Fantegrossi, W.E., et al., *3,4-Methylenedioxymethamphetamine (MDMA, "ecstasy") and its stereoisomers as reinforcers in rhesus monkeys: serotonergic involvement*. *Psychopharmacology (Berl)*, 2002. **161**(4): p. 356-64.
97. Fantegrossi, W.E., et al., *Role of dopamine transporters in the behavioral effects of 3,4-methylenedioxymethamphetamine (MDMA) in nonhuman primates*. *Psychopharmacology (Berl)*, 2009.
98. McClung, J., W. Fantegrossi, and L.L. Howell, *Reinstatement of extinguished amphetamine self-administration by 3,4-methylenedioxymethamphetamine (MDMA) and its enantiomers in rhesus monkeys*. *Psychopharmacology (Berl)*, 2010. **210**(1): p. 75-83.



99. Murnane, K.S., et al., *The neuropharmacology of prolactin secretion elicited by 3,4-methylenedioxymethamphetamine ("ecstasy"): a concurrent microdialysis and plasma analysis study*. *Horm Behav*, 2012. **61**(2): p. 181-90.
100. Spitzer, M., et al., *Enantio-selective cognitive and brain activation effects of N-ethyl-3,4-methylenedioxymethamphetamine in humans*. *Neuropharmacology*, 2001. **41**(2): p. 263-71.
101. Acquas, E., et al., *Differential effects of intravenous R,S-(+/-)-3,4-methylenedioxymethamphetamine (MDMA, Ecstasy) and its S(+)- and R(-)-enantiomers on dopamine transmission and extracellular signal regulated kinase phosphorylation (pERK) in the rat nucleus accumbens shell and core*. *J Neurochem*, 2007. **102**(1): p. 121-32.
102. Llabres, S., et al., *Molecular basis of the selective binding of MDMA enantiomers to the alpha4beta2 nicotinic receptor subtype: synthesis, pharmacological evaluation and mechanistic studies*. *Eur J Med Chem*, 2014. **81**: p. 35-46.
103. Navarro, M., et al., *Usefulness of saliva for measurement of 3,4-methylenedioxymethamphetamine and its metabolites: correlation with plasma drug concentrations and effect of salivary pH*. *Clin Chem*, 2001. **47**(10): p. 1788-95.
104. Kousik, S.M., T.C. Napier, and P.M. Carvey, *The effects of psychostimulant drugs on blood brain barrier function and neuroinflammation*. *Front Pharmacol*, 2012. **3**: p. 121.
105. Nichols, D.E., *Chromatographic Purity Of 3,4-Methylenedioxymethamphetamine Hydrochloride (MDMA Hydrochloride), Lot 5810-09*. 2006, Purdue University: Lafayette IN. p. 1-6.
106. Hardman, H.F., C.O. Haavik, and M.H. Seevers, *Relationship of the structure of mescaline and seven analogs to toxicity and behavior in five species of laboratory animals*. *Toxicol Appl Pharmacol*, 1973. **25**(2): p. 299-309.
107. Mehan, A., et al., *Pharmacokinetic profile of single and repeated oral doses of MDMA in squirrel monkeys: relationship to lasting effects on brain serotonin neurons*. *Neuropsychopharmacology*, 2006. **31**(2): p. 339-50.
108. Wang, X., et al., *(+/-)-3,4-Methylenedioxymethamphetamine administration to rats does not decrease levels of the serotonin transporter protein or alter its distribution between endosomes and the plasma membrane*. *J Pharmacol Exp Ther*, 2005. **314**(3): p. 1002-12.
109. Gaston, T.R., Rasmussen, G. T., *Identification of 3,4-methylenedioxy-n-methylamphetamine*. *Microgram*, 1972. **5**: p. 60-63.
110. Baggott, M., et al., *Chemical analysis of ecstasy pills*. *Jama*, 2000. **284**(17): p. 2190.
111. Cole, J.C., et al., *The content of ecstasy tablets: implications for the study of their long-term effects*. *Addiction*, 2002. **97**(12): p. 1531-6.
112. Tanner-Smith, E.E., *Pharmacological content of tablets sold as "ecstasy": results from an online testing service*. *Drug Alcohol Depend*, 2006. **83**(3): p. 247-54.
113. de la Torre, R. and M. Farre, *Neurotoxicity of MDMA (ecstasy): the limitations of scaling from animals to humans*. *Trends Pharmacol Sci*, 2004. **25**(10): p. 505-8.
114. Concheiro, M., et al., *Nonlinear pharmacokinetics of (+/-)3,4-methylenedioxymethamphetamine (MDMA) and its pharmacodynamic consequences in the rat*. *Drug Metab Dispos*, 2014. **42**(1): p. 119-25.
115. Scheidweiler, K.B., et al., *(+/-)-3,4-methylenedioxymethamphetamine and metabolite disposition in plasma and striatum of wild-type and multidrug resistance protein 1a knock-out mice*. *J Anal Toxicol*, 2011. **35**(7): p. 470-80.
116. Kolbrich, E.A., et al., *Plasma pharmacokinetics of 3,4-methylenedioxymethamphetamine after controlled oral administration to young adults*. *Ther Drug Monit*, 2008. **30**(3): p. 320-32.
117. Pacifici, R., et al., *Acute effects of 3,4-methylenedioxymethamphetamine alone and in combination with ethanol on the immune system in humans*. *J Pharmacol Exp Ther*, 2001. **296**(1): p. 207-15.

118. Farre, M., et al., *Repeated doses administration of MDMA in humans: pharmacological effects and pharmacokinetics*. Psychopharmacology (Berl), 2004. **173**(3-4): p. 364-75.
119. Mueller, M., et al., *Direct comparison of (+/-) 3,4-methylenedioxymethamphetamine ("ecstasy") disposition and metabolism in squirrel monkeys and humans*. Ther Drug Monit, 2009. **31**(3): p. 367-73.
120. Mueller, M., et al., *Simultaneous liquid chromatographic-electrospray ionization mass spectrometric quantification of 3,4-methylenedioxymethamphetamine (MDMA, Ecstasy) and its metabolites 3,4-dihydroxymethamphetamine, 4-hydroxy-3-methoxymethamphetamine and 3,4-methylenedioxyamphetamine in squirrel monkey and human plasma after acidic conjugate cleavage*. Forensic Sci Int, 2009. **184**(1-3): p. 64-8.
121. Mueller, M., et al., *Nonlinear pharmacokinetics of (+/-)3,4-methylenedioxymethamphetamine (MDMA, "Ecstasy") and its major metabolites in squirrel monkeys at plasma concentrations of MDMA that develop after typical psychoactive doses*. J Pharmacol Exp Ther, 2008. **327**(1): p. 38-44.
122. Mueller, M., et al., *Metabolism and disposition of 3,4-methylenedioxymethamphetamine ("ecstasy") in baboons after oral administration: comparison with humans reveals marked differences*. J Pharmacol Exp Ther, 2011. **338**(1): p. 310-7.
123. Johnson, M.P., A.J. Hoffman, and D.E. Nichols, *Effects of the enantiomers of MDA, MDMA and related analogues on [3H]serotonin and [3H]dopamine release from superfused rat brain slices*. Eur J Pharmacol, 1986. **132**(2-3): p. 269-76.
124. Cole, J.C. and H.R. Sumnall, *The pre-clinical behavioural pharmacology of 3,4-methylenedioxymethamphetamine (MDMA)*. Neurosci Biobehav Rev, 2003. **27**(3): p. 199-217.
125. Fitzgerald, J.L. and J.J. Reid, *Effects of methylenedioxymethamphetamine on the release of monoamines from rat brain slices*. Eur J Pharmacol, 1990. **191**(2): p. 217-20.
126. Kankaanpaa, A., et al., *The acute effects of amphetamine derivatives on extracellular serotonin and dopamine levels in rat nucleus accumbens*. Pharmacol Biochem Behav, 1998. **59**(4): p. 1003-9.
127. Verrico, C.D., G.M. Miller, and B.K. Madras, *MDMA (Ecstasy) and human dopamine, norepinephrine, and serotonin transporters: implications for MDMA-induced neurotoxicity and treatment*. Psychopharmacology (Berl), 2007. **189**(4): p. 489-503.
128. Cozzi, N.V., et al., *Inhibition of plasma membrane monoamine transporters by beta-ketoamphetamines*. Eur J Pharmacol, 1999. **381**(1): p. 63-9.
129. Partilla, J.S., et al., *Interaction of amphetamines and related compounds at the vesicular monoamine transporter*. J Pharmacol Exp Ther, 2006. **319**(1): p. 237-46.
130. Bogen, I.L., et al., *Short- and long-term effects of MDMA ("ecstasy") on synaptosomal and vesicular uptake of neurotransmitters in vitro and ex vivo*. Neurochem Int, 2003. **43**(4-5): p. 393-400.
131. Gu, X.F. and E.C. Azmitia, *Integrative transporter-mediated release from cytoplasmic and vesicular 5-hydroxytryptamine stores in cultured neurons*. Eur J Pharmacol, 1993. **235**(1): p. 51-7.
132. Leonardi, E.T. and E.C. Azmitia, *MDMA (ecstasy) inhibition of MAO type A and type B: comparisons with fenfluramine and fluoxetine (Prozac)*. Neuropsychopharmacology, 1994. **10**(4): p. 231-8.
133. Pilgrim, J.L., et al., *Serotonin toxicity involving MDMA (ecstasy) and moclobemide*. Forensic Sci Int, 2012. **215**(1-3): p. 184-8.
134. Vuori, E., et al., *Death following ingestion of MDMA (ecstasy) and moclobemide*. Addiction, 2003. **98**(3): p. 365-8.
135. Slikker, W., Jr., et al., *Behavioral and neurochemical effects of orally administered MDMA in the rodent and nonhuman primate*. Neurotoxicology, 1989. **10**(3): p. 529-42.
136. Mlinar, B. and R. Corradetti, *Endogenous 5-HT, released by MDMA through serotonin transporter- and secretory vesicle-dependent mechanisms, reduces hippocampal*

- excitatory synaptic transmission by preferential activation of 5-HT<sub>1B</sub> receptors located on CA1 pyramidal neurons.* Eur J Neurosci, 2003. **18**(6): p. 1559-71.
137. Rothman, R.B. and M.H. Baumann, *Monoamine transporters and psychostimulant drugs.* Eur J Pharmacol, 2003. **479**(1-3): p. 23-40.
138. Gudelsky, G.A. and B.K. Yamamoto, *Actions of 3,4-methylenedioxymethamphetamine (MDMA) on cerebral dopaminergic, serotonergic and cholinergic neurons.* Pharmacol Biochem Behav, 2008. **90**(2): p. 198-207.
139. Feduccia, A.A., N. Kongovi, and C.L. Duvauchelle, *Heat increases MDMA-enhanced NAcc 5-HT and body temperature, but not MDMA self-administration.* Eur Neuropsychopharmacol, 2010. **20**(12): p. 884-94.
140. Yamamoto, B.K. and L.J. Spanos, *The acute effects of methylenedioxymethamphetamine on dopamine release in the awake-behaving rat.* Eur J Pharmacol, 1988. **148**(2): p. 195-203.
141. Shankaran, M. and G.A. Gudelsky, *Effect of 3,4-methylenedioxymethamphetamine (MDMA) on hippocampal dopamine and serotonin.* Pharmacol Biochem Behav, 1998. **61**(4): p. 361-6.
142. Bradbury, S., et al., *Acquisition of MDMA self-administration: pharmacokinetic factors and MDMA-induced serotonin release.* Addict Biol, 2014. **19**(5): p. 874-84.
143. Acquas, E., et al., *Intravenous administration of ecstasy (3,4-methylenedioxymethamphetamine) enhances cortical and striatal acetylcholine release in vivo.* Eur J Pharmacol, 2001. **418**(3): p. 207-11.
144. Nair, S.G. and G.A. Gudelsky, *3,4-Methylenedioxymethamphetamine enhances the release of acetylcholine in the prefrontal cortex and dorsal hippocampus of the rat.* Psychopharmacology (Berl), 2006. **184**(2): p. 182-9.
145. Battaglia, G., et al., *Pharmacologic profile of MDMA (3,4-methylenedioxymethamphetamine) at various brain recognition sites.* Eur J Pharmacol, 1988. **149**(1-2): p. 159-63.
146. Jones, D.C., S.S. Lau, and T.J. Monks, *Thioether metabolites of 3,4-methylenedioxymethamphetamine and 3,4-methylenedioxymethamphetamine inhibit human serotonin transporter (hSERT) function and simultaneously stimulate dopamine uptake into hSERT-expressing SK-N-MC cells.* J Pharmacol Exp Ther, 2004. **311**(1): p. 298-306.
147. PDSP, *MDMA receptor binding profiles from Psychoactive Drug Screening Program Database Contract # NO1MH32004 (NIMH PDSP), with data accessed between November 12 and November 14, 2007.* 2007, National Institute of Mental Health.
148. Giannaccini, G., et al., *Short-term effects of 3,4-methylen-dioxy-metamphetamine (MDMA) on 5-HT(1A) receptors in the rat hippocampus.* Neurochem Int, 2007. **51**(8): p. 496-506.
149. Brunner, D. and R. Hen, *Insights into the neurobiology of impulsive behavior from serotonin receptor knockout mice.* Ann N Y Acad Sci, 1997. **836**: p. 81-105.
150. Graeff, F.G., et al., *Role of 5-HT in stress, anxiety, and depression.* Pharmacol Biochem Behav, 1996. **54**(1): p. 129-41.
151. Glennon, R.A. and R. Young, *MDMA stimulus generalization to the 5-HT(1A) serotonin agonist 8-hydroxy-2-(di-n-propylamino)tetralin.* 2000. **66**(3): p. 483-488.
152. Glennon, R.A., et al., *N-Methyl-1-(4-methoxyphenyl)-2-aminopropane (PMMA) and N-Methyl-1-(3,4-methylenedioxyphenyl)-2-aminopropane (MDMA) produce non-identical discriminative stimuli in rats.* Pharmacol Biochem Behav, 2007. **86**(3): p. 477-84.
153. Schechter, M.D., *Discriminative profile of MDMA.* Pharmacol Biochem Behav, 1986. **24**(6): p. 1533-7.
154. Nair, S.G. and G.A. Gudelsky, *3,4-Methylenedioxymethamphetamine (MDMA) enhances the release of acetylcholine by 5-HT<sub>4</sub> and D<sub>1</sub> receptor mechanisms in the rat prefrontal cortex.* Synapse, 2005. **58**(4): p. 229-35.

155. Kuteykin-Teplyakov, K. and R. Maldonado, *Looking for prosocial genes: ITRAQ analysis of proteins involved in MDMA-induced sociability in mice*. Eur Neuropsychopharmacol, 2014. **24**(11): p. 1773-83.
156. Bankson, M.G. and B.K. Yamamoto, *Serotonin-GABA interactions modulate MDMA-induced mesolimbic dopamine release*. J Neurochem, 2004. **91**(4): p. 852-9.
157. Cordero-Erausquin, M., et al., *Nicotinic receptor function: new perspectives from knockout mice*. Trends Pharmacol Sci, 2000. **21**(6): p. 211-7.
158. Jorgensen, H., et al., *Serotonin receptors involved in vasopressin and oxytocin secretion*. J Neuroendocrinol, 2003. **15**(3): p. 242-9.
159. Forsling, M.L., et al., *The effect of 3,4-methylenedioxymethamphetamine (MDMA, 'ecstasy') and its metabolites on neurohypophysial hormone release from the isolated rat hypothalamus*. Br J Pharmacol, 2002. **135**(3): p. 649-56.
160. Thompson, M.R., et al., *A role for oxytocin and 5-HT(1A) receptors in the prosocial effects of 3,4 methylenedioxymethamphetamine ("ecstasy")*. Neuroscience, 2007. **146**(2): p. 509-14.
161. Morley, K.C., J.C. Arnold, and I.S. McGregor, *Serotonin (1A) receptor involvement in acute 3,4-methylenedioxymethamphetamine (MDMA) facilitation of social interaction in the rat*. Prog Neuropsychopharmacol Biol Psychiatry, 2005. **29**(5): p. 648-57.
162. Broadbear, J.H., et al., *Oxytocinergic regulation of endogenous as well as drug-induced mood*. Pharmacol Biochem Behav, 2014. **119**: p. 61-71.
163. Miller, C.A. and J.D. Sweatt, *Covalent modification of DNA regulates memory formation*. Neuron, 2007. **53**(6): p. 857-69.
164. Unternaehrer, E., et al., *Childhood maternal care is associated with DNA methylation of the genes for brain-derived neurotrophic factor (BDNF) and oxytocin receptor (OXTR) in peripheral blood cells in adult men and women*. Stress, 2015. **18**(4): p. 451-61.
165. Koczor, C.A., et al., *Ecstasy (MDMA) Alters Cardiac Gene Expression and DNA Methylation: Implications for Circadian Rhythm Dysfunction in the Heart*. Toxicol Sci, 2015. **148**(1): p. 183-91.
166. Peng, W., et al., *Synaptotagmin I and IV are differentially regulated in the brain by the recreational drug 3,4-methylenedioxymethamphetamine (MDMA)*. Brain Res Mol Brain Res, 2002. **108**(1-2): p. 94-101.
167. Pompei, P., et al., *Preprotachykinin A gene expression after administration of 3,4-methylene dioxymethamphetamine (Ecstasy)*. Eur J Pharmacol, 2002. **450**(3): p. 245.
168. Sprague, J.E., et al. *Effects of antisense oligonucleotide alteration of MAO-B gene expression on MDMA-induced serotonergic neurotoxicity*. in Society for Neuroscience. 1999. Miami, FL.
169. Thiriet, N., et al., *Analysis of ecstasy (MDMA)-induced transcriptional responses in the rat cortex*. Faseb J, 2002. **16**(14): p. 1887-94.
170. Fernandez-Castillo, N., et al., *Active and passive MDMA ('ecstasy') intake induces differential transcriptional changes in the mouse brain*. Genes Brain Behav, 2012. **11**(1): p. 38-51.
171. Martinez-Turrillas, R., et al., *Differential effects of 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") on BDNF mRNA expression in rat frontal cortex and hippocampus*. Neurosci Lett, 2006. **402**(1-2): p. 126-30.
172. Sandrini, M., et al., *Causal role of prefrontal cortex in strengthening of episodic memories through reconsolidation*. Curr Biol, 2013. **23**(21): p. 2181-4.
173. Llado-Pelfort, L., et al., *5-HT1A receptor agonists enhance pyramidal cell firing in prefrontal cortex through a preferential action on GABA interneurons*. Cereb Cortex, 2012. **22**(7): p. 1487-97.
174. Biezonski, D.K. and J.S. Meyer, *Effects of 3,4-methylenedioxymethamphetamine (MDMA) on serotonin transporter and vesicular monoamine transporter 2 protein and*

- gene expression in rats: implications for MDMA neurotoxicity.* J Neurochem, 2010. **112**(4): p. 951-62.
175. Kindlundh-Hogberg, A.M., P. Svenningsson, and H.B. Schioth, *Quantitative mapping shows that serotonin rather than dopamine receptor mRNA expressions are affected after repeated intermittent administration of MDMA in rat brain.* Neuropharmacology, 2006. **51**(4): p. 838-47.
176. Salzman, J., et al., *Analysis of transcriptional responses in the mouse dorsal striatum following acute 3,4-methylenedioxymethamphetamine (ecstasy): identification of extracellular signal-regulated kinase-controlled genes.* Neuroscience, 2006. **137**(2): p. 473-82.
177. Karpova, N.N., *Role of BDNF epigenetics in activity-dependent neuronal plasticity.* Neuropharmacology, 2014. **76 Pt C**: p. 709-18.
178. Young, M.B., et al., *3,4-Methylenedioxymethamphetamine facilitates fear extinction learning.* Transl Psychiatry, 2015. **5**: p. e634.
179. Connor, T.J., et al., *Methylenedioxymethamphetamine (MDMA; Ecstasy) suppresses IL-1beta and TNF-alpha secretion following an in vivo lipopolysaccharide challenge.* Life Sci, 2000. **67**(13): p. 1601-12.
180. Pennock, J.W., et al., *3,4-Methylenedioxymethamphetamine increases susceptibility to genital herpes simplex virus infection in mice.* J Infect Dis, 2009. **200**(8): p. 1247-50.
181. de Paula, V.F., et al., *Methylenedioxymethamphetamine (Ecstasy) decreases neutrophil activity and alters leukocyte distribution in bone marrow, spleen and blood.* Neuroimmunomodulation, 2009. **16**(3): p. 191-200.
182. Boyle, N.T. and T.J. Connor, *MDMA ("Ecstasy") suppresses the innate IFN-gamma response in vivo: A critical role for the anti-inflammatory cytokine IL-10.* Eur J Pharmacol, 2007.
183. Stankevicius, D., et al., *3,4-methylenedioxymethamphetamine (ecstasy) decreases inflammation and airway reactivity in a murine model of asthma.* Neuroimmunomodulation, 2012. **19**(4): p. 209-19.
184. Passos, I.C., et al., *Inflammatory markers in post-traumatic stress disorder: a systematic review, meta-analysis, and meta-regression.* Lancet Psychiatry, 2015. **2**(11): p. 1002-12.
185. Gordon, C.J., *Thermophysiological responses to hyperthermic drugs: extrapolating from rodent to human.* Prog Brain Res, 2007. **162**: p. 63-79.
186. Reveron, M.E., E.Y. Maier, and C.L. Duvachelle, *Experience-dependent changes in temperature and behavioral activity induced by MDMA.* Physiol Behav, 2006. **89**(3): p. 358-63.
187. Kiyatkin, E.A., et al., *Critical role of peripheral vasoconstriction in fatal brain hyperthermia induced by MDMA (Ecstasy) under conditions that mimic human drug use.* J Neurosci, 2014. **34**(23): p. 7754-62.
188. Dafters, R.I., *Effect of ambient temperature on hyperthermia and hyperkinesia induced by 3,4-methylenedioxymethamphetamine (MDMA or "ecstasy") in rats.* Psychopharmacology (Berl), 1994. **114**(3): p. 505-8.
189. Fantegrossi, W.E., et al., *Pharmacological characterization of the effects of 3,4-methylenedioxymethamphetamine ("ecstasy") and its enantiomers on lethality, core temperature, and locomotor activity in singly housed and crowded mice.* Psychopharmacology (Berl), 2003. **166**(3): p. 202-11.
190. Malberg, J.E. and L.S. Seiden, *Small changes in ambient temperature cause large changes in 3,4-methylenedioxymethamphetamine (MDMA)-induced serotonin neurotoxicity and core body temperature in the rat.* J Neurosci, 1998. **18**(13): p. 5086-94.
191. Gordon, C.J., et al., *Effects of 3,4-methylenedioxymethamphetamine on autonomic thermoregulatory responses of the rat.* Pharmacol Biochem Behav, 1991. **38**(2): p. 339-44.

192. Pedersen, N.P. and W.W. Blessing, *Cutaneous vasoconstriction contributes to hyperthermia induced by 3,4-methylenedioxymethamphetamine (ecstasy) in conscious rabbits*. *J Neurosci*, 2001. **21**(21): p. 8648-54.
193. Von Huben, S.N., et al., *Impact of ambient temperature on hyperthermia induced by (+/-)3,4-methylenedioxymethamphetamine in rhesus macaques*. *Neuropsychopharmacology*, 2007. **32**(3): p. 673-81.
194. Bowyer, J.F., et al., *Plasma levels of parent compound and metabolites after doses of either d-fenfluramine or d-3,4-methylenedioxymethamphetamine (MDMA) that produce long-term serotonergic alterations*. *Neurotoxicology*, 2003. **24**(3): p. 379-90.
195. Freedman, R.R., C.E. Johanson, and M.E. Tancer, *Thermoregulatory effects of 3,4-methylenedioxymethamphetamine (MDMA) in humans*. *Psychopharmacology (Berl)*, 2005. **183**(2): p. 248-56.
196. Crean, R.D., S.A. Davis, and M.A. Taffe, *Oral administration of (+/-)3,4-methylenedioxymethamphetamine and (+)methamphetamine alters temperature and activity in rhesus macaques*. *Pharmacol Biochem Behav*, 2007. **87**(1): p. 11-9.
197. Crean, R.D., et al., *Effects of (+/-)3,4-methylenedioxymethamphetamine, (+/-)3,4-methylenedioxyamphetamine and methamphetamine on temperature and activity in rhesus macaques*. *Neuroscience*, 2006. **142**(2): p. 515-25.
198. Banks, M.L., et al., *Ambient temperature effects on 3,4-methylenedioxymethamphetamine-induced thermodyregulation and pharmacokinetics in male monkeys*. *Drug Metab Dispos*, 2007. **35**(10): p. 1840-5.
199. Schindler, C.W., et al., *Effects of 3,4-methylenedioxymethamphetamine (MDMA) and its main metabolites on cardiovascular function in conscious rats*. *Br J Pharmacol*, 2014. **171**(1): p. 83-91.
200. Tiangco, D.A., et al., *3,4-Methylenedioxymethamphetamine activates nuclear factor-kappaB, increases intracellular calcium, and modulates gene transcription in rat heart cells*. *Cardiovasc Toxicol*, 2005. **5**(3): p. 301-10.
201. Vandeputte, C. and J.R. Docherty, *Vascular actions of 3,4-methylenedioxymethamphetamine in alpha(2A/D)- adrenoceptor knockout mice*. *Eur J Pharmacol*, 2002. **457**(1): p. 45-9.
202. Bexis, S. and J.R. Docherty, *Effects of MDMA, MDA and MDEA on blood pressure, heart rate, locomotor activity and body temperature in the rat involve alpha-adrenoceptors*. *Br J Pharmacol*, 2006. **147**(8): p. 926-34.
203. Bexis, S. and J.R. Docherty, *Role of alpha2A-adrenoceptors in the effects of MDMA on body temperature in the mouse*. *Br J Pharmacol*, 2005. **146**(1): p. 1-6.
204. Quinn, S.T., et al., *Blockade of noradrenaline transport abolishes 4-methylthioamphetamine-induced contraction of the rat aorta in vitro*. *Auton Autacoid Pharmacol*, 2006. **26**(4): p. 335-44.
205. Hysek, C.M., et al., *Effects of the alpha(2)-adrenergic agonist clonidine on the pharmacodynamics and pharmacokinetics of 3,4-methylenedioxymethamphetamine in healthy volunteers*. *J Pharmacol Exp Ther*, 2012. **340**(2): p. 286-94.
206. Hysek, C.M. and M.E. Liechti, *Effects of MDMA alone and after pretreatment with reboxetine, duloxetine, clonidine, carvedilol, and doxazosin on pupillary light reflex*. *Psychopharmacology (Berl)*, 2012. **224**(3): p. 363-76.
207. Hysek, C.M., et al., *Carvedilol inhibits the cardiostimulant and thermogenic effects of MDMA in humans: Lost in translation*. *Br J Pharmacol*, 2013. **170**(6): p. 1273-5.
208. Hysek, C.M., et al., *Duloxetine inhibits effects of MDMA ("ecstasy") in vitro and in humans in a randomized placebo-controlled laboratory study*. *PLoS One*, 2012. **7**(5): p. e36476.
209. Burbach, J., L. Young, and J. Russell, *Oxytocin: synthesis, secretion, and reproductive functions*. *Knobil and Neill's physiology of reproduction*, 2006. **2**: p. 3055-3128.

210. Tasker, J.G., et al., *Cell biology of oxytocin and vasopressin cells*, in *Hormones, brain and behavior*, Pfaff D.W., et al., Editors. 2002, Academic Press: London. p. 811-842.
211. Hargreaves, G.A., et al., *High ambient temperature increases 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy")-induced Fos expression in a region-specific manner*. *Neuroscience*, 2007. **145**(2): p. 764-74.
212. Fantegrossi, W.E., et al., *Nantenine: an antagonist of the behavioral and physiological effects of MDMA in mice*. *Psychopharmacology (Berl)*, 2004. **173**(3-4): p. 270-7.
213. Meyer, J.S., et al., *Neural effects of MDMA as determined by functional magnetic resonance imaging and magnetic resonance spectroscopy in awake marmoset monkeys*. *Ann N Y Acad Sci*, 2006. **1074**: p. 365-76.
214. Karimi, S., M. Jahanshahi, and M.J. Golalipour, *The Effect of MDMA-Induced Anxiety on Neuronal Apoptosis in Adult Male Rats' Hippocampus*. *Folia Biol (Praha)*, 2014. **60**(4): p. 187-91.
215. Isbister, G.K. and N.A. Buckley, *The pathophysiology of serotonin toxicity in animals and humans: implications for diagnosis and treatment*. *Clin Neuropharmacol*, 2005. **28**(5): p. 205-14.
216. Gillman, P.K., *Ecstasy, serotonin syndrome and the treatment of hyperpyrexia*. *Med J Aust*, 1997. **167**(2): p. 109, 111.
217. Mueller, P.D. and W.S. Korey, *Death by "ecstasy": the serotonin syndrome?* *Ann Emerg Med*, 1998. **32**(3 Pt 1): p. 377-80.
218. Ferraz de Paula, V., et al., *Methylenedioxymethamphetamine (Ecstasy) Decreases Neutrophil Activity and Alters Leukocyte Distribution in Bone Marrow, Spleen and Blood*. *Neuroimmunomodulation*, 2009. **16**(3): p. 191-200.
219. Nash, J.F., Jr., H.Y. Meltzer, and G.A. Gudelsky, *Elevation of serum prolactin and corticosterone concentrations in the rat after the administration of 3,4-methylenedioxymethamphetamine*. *J Pharmacol Exp Ther*, 1988. **245**(3): p. 873-9.
220. Connor, T.J., J.P. Kelly, and B.E. Leonard, *An assessment of the acute effects of the serotonin releasers methylenedioxymethamphetamine, methylenedioxyamphetamine and fenfluramine on immunity in rats*. *Immunopharmacology*, 2000. **46**(3): p. 223-35.
221. Balogh, B., et al., *Effects of a single dose of 3,4-methylenedioxymethamphetamine on circadian patterns, motor activity and sleep in drug-naive rats and rats previously exposed to MDMA*. *Psychopharmacology (Berl)*, 2004. **173**(3-4): p. 296-309.
222. Bhattacharya, S.K., A. Bhattacharya, and S. Ghosal, *Anxiogenic activity of methylenedioxymethamphetamine (Ecstasy): an experimental study*. *Biogenic Amines*, 1998. **7**: p. 217-237.
223. Ho, Y.J., et al., *Acute and long-term consequences of single MDMA administration in relation to individual anxiety levels in the rat*. *Behav Brain Res*, 2004. **149**(2): p. 135-44.
224. Selken, J. and D.E. Nichols, *Alpha1-adrenergic receptors mediate the locomotor response to systemic administration of (+/-)-3,4-methylenedioxymethamphetamine (MDMA) in rats*. *Pharmacol Biochem Behav*, 2007. **86**(4): p. 622-30.
225. Mechan, A.O., et al., *A study of the effect of a single neurotoxic dose of 3,4-methylenedioxymethamphetamine (MDMA; "ecstasy") on the subsequent long-term behaviour of rats in the plus maze and open field*. *Psychopharmacology (Berl)*, 2002. **159**(2): p. 167-75.
226. Gurtman, C.G., et al., *Increased anxiety in rats after 3,4-methylenedioxymethamphetamine: association with serotonin depletion*. *Eur J Pharmacol*, 2002. **446**(1-3): p. 89-96.
227. Clemens, K.J., et al., *MDMA ("ecstasy"), methamphetamine and their combination: long-term changes in social interaction and neurochemistry in the rat*. *Psychopharmacology (Berl)*, 2004. **173**(3-4): p. 318-25.



228. Ando, R.D., et al., *Partial lesion of the serotonergic system by a single dose of MDMA results in behavioural disinhibition and enhances acute MDMA-induced social behaviour on the social interaction test*. *Neuropharmacology*, 2006. **50**(7): p. 884-96.
229. Thompson, M.R., G.E. Hunt, and I.S. McGregor, *Neural correlates of MDMA ("Ecstasy")-induced social interaction in rats*. *Soc Neurosci*, 2009. **4**(1): p. 60-72.
230. Hunt, G.E., et al., *MDMA-induced c-Fos expression in oxytocin-containing neurons is blocked by pretreatment with the 5-HT-1A receptor antagonist WAY 100635*. *Brain Res Bull*, 2011. **86**(1-2): p. 65-73.
231. Hasler, F., et al., *Investigation of serotonin-1A receptor function in the human psychopharmacology of MDMA*. *J Psychopharmacol*, 2009. **23**(8): p. 923-35.
232. Hysek, C.M., F.X. Vollenweider, and M.E. Liechti, *Effects of a beta-blocker on the cardiovascular response to MDMA (Ecstasy)*. *Emerg Med J*, 2010. **27**(8): p. 586-9.
233. van Wel, J.H., et al., *Blockade of 5-HT(2) Receptor Selectively Prevents MDMA-Induced Verbal Memory Impairment*. *Neuropsychopharmacology*, 2011. **36**(9): p. 1932-9.
234. Liester, M.B., et al., *Phenomenology and sequelae of 3,4-methylenedioxymethamphetamine use*. *J Nerv Ment Dis*, 1992. **180**(6): p. 345-52; discussion 353-4.
235. Peroutka, S.J., H. Newman, and H. Harris, *Subjective effects of 3,4-methylenedioxymethamphetamine in recreational users*. *Neuropsychopharmacology*, 1988. **1**(4): p. 273-7.
236. Halpern, J.H., et al., *Residual neurocognitive features of long-term ecstasy users with minimal exposure to other drugs*. *Addiction*, 2011. **106**(4): p. 777-86.
237. Hahn, I.-H. *MDMA Toxicity*. Medscape, 2015.
238. Cohen, R.S., *Subjective reports on the effects of the MDMA ('ecstasy') experience in humans*. *Prog Neuropsychopharmacol Biol Psychiatry*, 1995. **19**(7): p. 1137-45.
239. Litjens, R.P., et al., *Hallucinogen persisting perception disorder and the serotonergic system: a comprehensive review including new MDMA-related clinical cases*. *Eur Neuropsychopharmacol*, 2014. **24**(8): p. 1309-23.
240. Baggott, M.J., et al., *Intimate insight: MDMA changes how people talk about significant others*. *J Psychopharmacol*, 2015.
241. Pacifici, R., et al., *Combined immunomodulating properties of 3,4-methylenedioxymethamphetamine (MDMA) and cannabis in humans*. *Addiction*, 2007. **102**(6): p. 931-6.
242. Parrott, A.C., *MDMA and temperature: a review of the thermal effects of 'Ecstasy' in humans*. *Drug Alcohol Depend*, 2012. **121**(1-2): p. 1-9.
243. Dafters, R.I., *Hyperthermia following MDMA administration in rats: effects of ambient temperature, water consumption, and chronic dosing*. *Physiol Behav*, 1995. **58**(5): p. 877-82.
244. Docherty, J.R. and A.R. Green, *The role of monoamines in the changes in body temperature induced by 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) and its derivatives*. *Br J Pharmacol*, 2010. **160**(5): p. 1029-44.
245. Martin, T.L., D.A. Chiasson, and S.J. Kish, *Does hyperthyroidism increase risk of death due to the ingestion of ecstasy?* *J Forensic Sci*, 2007. **52**(4): p. 951-3.
246. Sprague, J.E., et al., *Roles of norepinephrine, free Fatty acids, thyroid status, and skeletal muscle uncoupling protein 3 expression in sympathomimetic-induced thermogenesis*. *J Pharmacol Exp Ther*, 2007. **320**(1): p. 274-80.
247. Droogmans, S., et al., *Possible association between 3,4-methylenedioxymethamphetamine abuse and valvular heart disease*. *Am J Cardiol*, 2007. **100**(9): p. 1442-5.
248. Farah, R. and R. Farah, *Ecstasy (3,4-methylenedioxymethamphetamine)-induced inappropriate antidiuretic hormone secretion*. *Pediatr Emerg Care*, 2008. **24**(9): p. 615-7.

249. Esposito, P., et al., *The syndrome of inappropriate antidiuresis: pathophysiology, clinical management and new therapeutic options*. Nephron Clin Pract, 2011. **119**(1): p. c62-73; discussion c73.
250. Laycock, J., *Introduction to vasopressin*. Perspectives on vasopressin. London: World Scientific Publishing Company, Imperial College Press, 2010: p. 1-16.
251. Forsling, M., et al., *Arginine vasopressin release in response to the administration of 3,4-methylenedioxymethamphetamine ("ecstasy"): is metabolism a contributory factor?* J Pharm Pharmacol, 2001. **53**(10): p. 1357-63.
252. Aitchison, K.J., et al., *Ecstasy (MDMA)-induced hyponatraemia is associated with genetic variants in CYP2D6 and COMT*. J Psychopharmacol, 2012. **26**(3): p. 408-18.
253. Henry, J.A. and I.R. Hill, *Fatal interaction between ritonavir and MDMA*. Lancet, 1998. **352**(9142): p. 1751-2.
254. Semple, D.M., et al., *Reduced in vivo binding to the serotonin transporter in the cerebral cortex of MDMA ('ecstasy') users*. Br J Psychiatry, 1999. **175**: p. 63-9.
255. Gouzoulis-Mayfrank, E. and J. Daumann, *Neurotoxicity of methylenedioxyamphetamines (MDMA; ecstasy) in humans: how strong is the evidence for persistent brain damage?* Addiction, 2006. **101**(3): p. 348-61.
256. Buchert, R., et al., *A voxel-based PET investigation of the long-term effects of "Ecstasy" consumption on brain serotonin transporters*. Am J Psychiatry, 2004. **161**(7): p. 1181-9.
257. Gouzoulis-Mayfrank, E., et al., *Memory impairment suggests hippocampal dysfunction in abstinent ecstasy users*. Prog Neuropsychopharmacol Biol Psychiatry, 2003. **27**(5): p. 819-27.
258. Halpern, J.H., et al., *Residual neuropsychological effects of illicit 3,4-methylenedioxymethamphetamine (MDMA) in individuals with minimal exposure to other drugs*. Drug Alcohol Depend, 2004. **75**(2): p. 135-47.
259. Thomasius, R., et al., *Mood, cognition and serotonin transporter availability in current and former ecstasy (MDMA) users*. Psychopharmacology (Berl), 2003. **167**(1): p. 85-96.
260. Daumann, J., Jr., et al., *Neural mechanisms of working memory in ecstasy (MDMA) users who continue or discontinue ecstasy and amphetamine use: evidence from an 18-month longitudinal functional magnetic resonance imaging study*. Biol Psychiatry, 2004. **56**(5): p. 349-55.
261. Gouzoulis-Mayfrank, E., et al., *Memory performance in polyvalent MDMA (ecstasy) users who continue or discontinue MDMA use*. Drug Alcohol Depend, 2005. **78**(3): p. 317-23.
262. Buchert, R., et al., *Reversibility of ecstasy-induced reduction in serotonin transporter availability in polydrug ecstasy users*. Eur J Nucl Med Mol Imaging, 2006. **33**(2): p. 188-99.
263. Selvaraj, S., et al., *Brain serotonin transporter binding in former users of MDMA ('ecstasy')*. Br J Psychiatry, 2009. **194**(4): p. 355-9.
264. Kish, S.J., et al., *Low striatal serotonin transporter protein in a human polydrug MDMA (ecstasy) user: a case study*. J Psychopharmacol, 2010. **24**(2): p. 281-4.
265. Erritzoe, D., et al., *In vivo imaging of cerebral serotonin transporter and serotonin(2A) receptor binding in 3,4-methylenedioxymethamphetamine (MDMA or "ecstasy") and hallucinogen users*. Arch Gen Psychiatry, 2011. **68**(6): p. 562-76.
266. de Win, M.M., et al., *A Prospective Cohort Study on Sustained Effects of Low-Dose Ecstasy Use on the Brain in New Ecstasy Users*. Neuropsychopharmacology, 2007. **32**(2): p. 458-470.
267. Chang, L., et al., *Cerebral (1)H MRS alterations in recreational 3, 4-methylenedioxymethamphetamine (MDMA, "ecstasy") users*. J Magn Reson Imaging, 1999. **10**(4): p. 521-6.
268. Cowan, R.L., et al., *Occipital cortical proton MRS at 4 Tesla in human moderate MDMA polydrug users*. Psychiatry Res, 2007.

269. Reneman, L., et al., *Reduced N-acetylaspartate levels in the frontal cortex of 3,4-methylenedioxymethamphetamine (Ecstasy) users: preliminary results*. AJNR Am J Neuroradiol, 2002. **23**(2): p. 231-7.
270. De Win, M., Jager, G., Reneman, L., Booij, J., van den Brink, W., Den Heeten, G., et al., *Ecstasy: Is It Safe for the Brain? First Prospective Study on Effects of Low Doses of Ecstasy on the Brain in New Ecstasy Users, Using a Combination of Advanced MR Imaging Techniques and [<sup>123</sup>I]β-CIT SPECT*, in *Radiological Society of North America (RSNA)*. 2006: Chicago, IL.
271. Klomp, A., et al., *The Effects of Ecstasy (MDMA) on Brain Serotonin Transporters Are Dependent on Age-of-First Exposure in Recreational Users and Animals*. PLoS One, 2012. **7**(10): p. e47524.
272. Booij, L., et al., *Brain serotonin synthesis in MDMA (ecstasy) polydrug users: an alpha-[(11) C]methyl-l-tryptophan study*. J Neurochem, 2014. **131**(5): p. 634-44.
273. Bauernfeind, A.L., et al., *Human ecstasy use is associated with increased cortical excitability: an fMRI study*. Neuropsychopharmacology, 2011. **36**(6): p. 1127-41.
274. Salomon, R.M., et al., *MDMA (Ecstasy) association with impaired fMRI BOLD thalamic coherence and functional connectivity*. Drug Alcohol Depend, 2012. **120**(1-3): p. 41-7.
275. Roberts, C.A., et al., *Differences in prefrontal blood oxygenation during an acute multitasking stressor in ecstasy polydrug users*. Psychol Med, 2014: p. 1-12.
276. Jager, G., et al., *Incidental use of ecstasy: no evidence for harmful effects on cognitive brain function in a prospective fMRI study*. Psychopharmacology (Berl), 2007. **193**(3): p. 403-14.
277. Roiser, J.P., et al., *The effect of polymorphism at the serotonin transporter gene on decision-making, memory and executive function in ecstasy users and controls*. Psychopharmacology (Berl), 2006. **188**(2): p. 213-27.
278. Cuyas, E., et al., *The Influence of Genetic and Environmental Factors among MDMA Users in Cognitive Performance*. PLoS One, 2011. **6**(11): p. e27206.
279. Schilt, T., et al., *The effect of Ecstasy on memory is moderated by a functional polymorphism in the catechol-O-methyltransferase (COMT) gene*. Eur Neuropsychopharmacol, 2009. **19**(2): p. 116-24.
280. Parrott, A.C., *MDMA, serotonergic neurotoxicity, and the diverse functional deficits of recreational 'Ecstasy' users*. Neurosci Biobehav Rev, 2013. **37**(8): p. 1466-1484.
281. Randall, S., et al., *Effects of acute 3,4-methylenedioxymethamphetamine on sleep and daytime sleepiness in MDMA users: a preliminary study*. Sleep, 2009. **32**(11): p. 1513-9.
282. Carhart-Harris, R.L., et al., *Equivalent effects of acute tryptophan depletion on REM sleep in ecstasy users and controls*. Psychopharmacology (Berl), 2009. **206**(2): p. 187-96.
283. Allen, R.P., U.D. McCann, and G.A. Ricaurte, *Persistent effects of (+/- )3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") on human sleep*. Sleep, 1993. **16**(6): p. 560-4.
284. McCann, U.D., S.C. Peterson, and G.A. Ricaurte, *The effect of catecholamine depletion by alpha-methyl-para-tyrosine on measures of cognitive performance and sleep in abstinent MDMA users*. Neuropsychopharmacology, 2007. **32**(8): p. 1695-706.
285. McCann, U.D., et al., *Sleep deprivation differentially impairs cognitive performance in abstinent methylenedioxymethamphetamine ("ecstasy") users*. J Neurosci, 2009. **29**(44): p. 14050-6.
286. McCann, U.D., et al., *Sleep apnea in young abstinent recreational MDMA ("ecstasy") consumers*. Neurology, 2009. **73**(23): p. 2011-7.
287. MacInnes, N., S.L. Handley, and G.F. Harding, *Former chronic methylenedioxymethamphetamine (MDMA or ecstasy) users report mild depressive symptoms*. J Psychopharmacol, 2001. **15**(3): p. 181-6.
288. Parrott, A.C., E. Sisk, and J.J. Turner, *Psychobiological problems in heavy 'ecstasy' (MDMA) polydrug users*. Drug Alcohol Depend, 2000. **60**(1): p. 105-10.

289. Sumnall, H.R. and J.C. Cole, *Self-reported depressive symptomatology in community samples of polysubstance misusers who report Ecstasy use: a meta-analysis*. J Psychopharmacol, 2005. **19**(1): p. 84-92.
290. Milani, R.M., et al., *Gender differences in self-reported anxiety, depression, and somatization among ecstasy/MDMA polydrug users, alcohol/tobacco users, and nondrug users*. Addict Behav, 2004. **29**(5): p. 965-71.
291. Sumnall, H.R., G.F. Wagstaff, and J.C. Cole, *Self-reported psychopathology in polydrug users*. J Psychopharmacol, 2004. **18**(1): p. 75-82.
292. Medina, K.L. and P.K. Shear, *Anxiety, depression, and behavioral symptoms of executive dysfunction in ecstasy users: contributions of polydrug use*. Drug Alcohol Depend, 2007. **87**(2-3): p. 303-11.
293. Fisk, J.E., et al., *Modelling the adverse effects associated with ecstasy use*. Addiction, 2011. **106**(4): p. 798-805.
294. Scott, L.A., et al., *The impact of comorbid cannabis and methamphetamine use on mental health among regular ecstasy users*. Addict Behav, 2012. **37**(9): p. 1058-62.
295. Daumann, J., et al., *Self-reported psychopathological symptoms in recreational ecstasy (MDMA) users are mainly associated with regular cannabis use: further evidence from a combined cross-sectional/longitudinal investigation*. Psychopharmacology (Berl), 2004. **173**(3-4): p. 398-404.
296. Whitaker-Azmitia, P.M. and T.A. Aronson, *"Ecstasy" (MDMA)-induced panic*. Am J Psychiatry, 1989. **146**(1): p. 119.
297. McCann, U.D. and G.A. Ricaurte, *Lasting neuropsychiatric sequelae of (+-)-methylenedioxyamphetamine ('ecstasy') in recreational users*. J Clin Psychopharmacol, 1991. **11**(5): p. 302-5.
298. Pallanti, S. and D. Mazzi, *MDMA (Ecstasy) precipitation of panic disorder*. Biol Psychiatry, 1992. **32**(1): p. 91-5.
299. McCann, U.D. and G.A. Ricaurte, *MDMA ("ecstasy") and panic disorder: induction by a single dose*. Biol Psychiatry, 1992. **32**(10): p. 950-3.
300. Allott, K., et al., *Neuroendocrine and subjective responses to pharmacological challenge with citalopram: a controlled study in male and female ecstasy/MDMA users*. J Psychopharmacol, 2008.
301. Wetherell, M.A. and C. Montgomery, *Basal functioning of the hypothalamic-pituitary-adrenal (HPA) axis and psychological distress in recreational ecstasy polydrug users*. Psychopharmacology (Berl), 2014. **231**(7): p. 1365-75.
302. Frokjaer, V.G., et al., *In abstinent MDMA users the cortisol awakening response is off-set but associated with prefrontal serotonin transporter binding as in non-users*. Int J Neuropsychopharmacol, 2014. **17**(8): p. 1119-28.
303. George, A.M., S. Olesen, and R.J. Tait, *Ecstasy use and depression: a 4-year longitudinal study among an Australian general community sample*. Psychopharmacology (Berl), 2013. **229**(4): p. 713-21.
304. Thomasius, R., et al., *Mood, cognition and serotonin transporter availability in current and former ecstasy (MDMA) users: the longitudinal perspective*. J Psychopharmacol, 2006. **20**(2): p. 211-25.
305. Schilt, T., et al., *Cognition in novice ecstasy users with minimal exposure to other drugs: a prospective cohort study*. Arch Gen Psychiatry, 2007. **64**(6): p. 728-36.
306. White, C., J. Brown, and M. Edwards, *Alterations to global but not local motion processing in long-term ecstasy (MDMA) users*. Psychopharmacology (Berl), 2014.
307. Bedi, G. and J. Redman, *Metamemory in recreational ecstasy polydrug users: what do self-reports of memory failures mean?* J Psychopharmacol, 2008.
308. Bedi, G. and J. Redman, *Ecstasy use and higher-level cognitive functions: weak effects of ecstasy after control for potential confounds*. Psychol Med, 2008. **38**(9): p. 1319-30.

309. Hanson, K.L. and M. Luciana, *Neurocognitive impairments in MDMA and other drug users: MDMA alone may not be a cognitive risk factor*. J Clin Exp Neuropsychol, 2010. **32**(4): p. 337-49.
310. Raj, V., et al., *MDMA (ecstasy) use is associated with reduced BOLD signal change during semantic recognition in abstinent human polydrug users: a preliminary fMRI study*. J Psychopharmacol, 2010. **24**(2): p. 187-201.
311. Roiser, J.P., R.D. Rogers, and B.J. Sahakian, *Neuropsychological function in ecstasy users: a study controlling for polydrug use*. Psychopharmacology (Berl), 2007. **189**(4): p. 505-16.
312. Blagrove, M., et al., *Procedural and declarative memory task performance, and the memory consolidation function of sleep, in recent and abstinent ecstasy/MDMA users*. J Psychopharmacol, 2011. **25**(4): p. 465-77.
313. Kish, S.J., et al., *Decreased cerebral cortical serotonin transporter binding in ecstasy users: a positron emission tomography/[<sup>11</sup>C]DASB and structural brain imaging study*. Brain, 2010. **133**(Pt 6): p. 1779-97.
314. Montgomery, C. and J.E. Fisk, *Ecstasy-related deficits in the updating component of executive processes*. Hum Psychopharmacol, 2008.
315. Hadjiefthymoulou, F., et al., *Everyday and prospective memory deficits in ecstasy/polydrug users*. J Psychopharmacol, 2011. **25**(4): p. 453-64.
316. van Holst, R.J. and T. Schilt, *Drug-related decrease in neuropsychological functions of abstinent drug users*. Curr Drug Abuse Rev, 2011. **4**(1): p. 42-56.
317. Jager, G., et al., *Assessment of cognitive brain function in ecstasy users and contributions of other drugs of abuse: results from an FMRI study*. Neuropsychopharmacology, 2008. **33**(2): p. 247-58.
318. Indlekofer, F., et al., *Reduced memory and attention performance in a population-based sample of young adults with a moderate lifetime use of cannabis, ecstasy and alcohol*. J Psychopharmacol, 2009. **23**(5): p. 495-509.
319. Schilt, T., et al., *Long-term neuropsychological effects of ecstasy in middle-aged ecstasy/polydrug users*. Psychopharmacology (Berl), 2010. **207**(4): p. 583-91.
320. Morgan, M.J., *Recreational use of "ecstasy" (MDMA) is associated with elevated impulsivity*. Neuropsychopharmacology, 1998. **19**(4): p. 252-64.
321. Quednow, B.B., et al., *Elevated impulsivity and impaired decision-making cognition in heavy users of MDMA ("Ecstasy")*. Psychopharmacology (Berl), 2007. **189**(4): p. 517-30.
322. Morgan, M.J., et al., *Elevated impulsivity and impaired decision-making in abstinent Ecstasy (MDMA) users compared to polydrug and drug-naïve controls*. Neuropsychopharmacology, 2006. **31**(7): p. 1562-73.
323. Schilt, T., et al., *Decision making as a predictor of first ecstasy use: a prospective study*. Psychopharmacology (Berl), 2009. **203**(3): p. 519-27.
324. Pirona, A. and M.J. Morgan, *An investigation of the subacute effects of ecstasy on neuropsychological performance, sleep and mood in regular ecstasy users*. J Psychopharmacol, 2010. **24**(2): p. 175-85.
325. Hanson, K.L., M. Luciana, and K. Sullwold, *Reward-related decision-making deficits and elevated impulsivity among MDMA and other drug users*. Drug Alcohol Depend, 2008. **96**(1-2): p. 99-110.
326. Mueller, M., et al., *Single oral doses of (+/-) 3,4-methylenedioxymethamphetamine ('Ecstasy') produce lasting serotonergic deficits in non-human primates: relationship to plasma drug and metabolite concentrations*. Int J Neuropsychopharmacol, 2013. **16**(4): p. 791-801.
327. de la Torre, R., et al., *MDMA, methamphetamine, and CYP2D6 pharmacogenetics: what is clinically relevant?* Front Genet, 2012. **3**: p. 235.

328. Peiro, A.M., et al., *Human pharmacology of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) after repeated doses taken 2 h apart*. *Psychopharmacology (Berl)*, 2013. **225**(4): p. 883-93.
329. Mahmood, I. and J.D. Balian, *Interspecies scaling: a comparative study for the prediction of clearance and volume using two or more than two species*. *Life Sci*, 1996. **59**(7): p. 579-85.
330. Frith, C.H., et al., *Toxicity of methylenedioxymethamphetamine (MDMA) in the dog and the rat*. *Fundam Appl Toxicol*, 1987. **9**(1): p. 110-9.
331. Hariri, M., et al., *ASSESSING MUTAGENICITY OF THE DRUG MDMA (ECSTASY) AVAILABLE IN IRAN USING AMES BIOASSAY*. *Jundishapur Journal of Natural Pharmaceutical Products*, 2010. **5**(1): p. 44-49.
332. Yoshioka, H., et al., *Genotoxicity study of illegal drug MDMA and its nitroso derivative N-MDMA by micronucleus and chromosomal aberration tests using Chinese hamster lung fibroblast cell line*. *Environ Health Prev Med*, 2007. **12**(3): p. 129-37.
333. Campbell, N.G., et al., *MDMA administration to pregnant Sprague-Dawley rats results in its passage to the fetal compartment*. *Neurotoxicol Teratol*, 2006. **28**(4): p. 459-65.
334. St Omer, V.E., et al., *Behavioral and neurochemical effects of prenatal methylenedioxymethamphetamine (MDMA) exposure in rats*. *Neurotoxicol Teratol*, 1991. **13**(1): p. 13-20.
335. Plessinger, M.A., *Prenatal exposure to amphetamines. Risks and adverse outcomes in pregnancy*. *Obstet Gynecol Clin North Am*, 1998. **25**(1): p. 119-38.
336. Kwack, S.J., et al., *A One-Generation Reproductive Toxicity Study of 3,4-Methylenedioxy-N-Methamphetamine (MDMA, Ecstasy), an Amphetamine Derivative, in C57BL/6 Mice*. *J Toxicol Environ Health A*, 2014. **77**(22-24): p. 1431-42.
337. Barenys, M., et al., *MDMA (ecstasy) delays pubertal development and alters sperm quality after developmental exposure in the rat*. *Toxicol Lett*, 2010. **197**(2): p. 135-42.
338. Barenys, M., et al., *Chronic exposure to MDMA (ecstasy) increases DNA damage in sperm and alters testes histopathology in male rats*. *Toxicol Lett*, 2009. **191**(1): p. 40-6.
339. Koprach, J.B., N.G. Campbell, and J.W. Lipton, *Neonatal 3,4-methylenedioxymethamphetamine (ecstasy) alters dopamine and serotonin neurochemistry and increases brain-derived neurotrophic factor in the forebrain and brainstem of the rat*. *Brain Res Dev Brain Res*, 2003. **147**(1-2): p. 177-82.
340. Koprach, J.B., et al., *Prenatal 3,4-methylenedioxymethamphetamine (ecstasy) alters exploratory behavior, reduces monoamine metabolism, and increases forebrain tyrosine hydroxylase fiber density of juvenile rats*. *Neurotoxicol Teratol*, 2003. **25**(5): p. 509-17.
341. Cohen, M.A., et al., *Learning and memory after neonatal exposure to 3,4-methylenedioxymethamphetamine (ecstasy) in rats: interaction with exposure in adulthood*. *Synapse*, 2005. **57**(3): p. 148-59.
342. Piper, B.J., J.D. Farelli, and J.S. Meyer, *Dissociation between serotonin neurotoxicity and brain-derived neurotrophic factor induction following neonatal MDMA exposure in rats*. *Dev Neurosci*, 2009. **31**(1-2): p. 90-4.
343. Piper, B.J. and J.S. Meyer, *Increased responsiveness to MDMA in adult rats treated neonatally with MDMA*. *Neurotoxicol Teratol*, 2006. **28**(1): p. 95-102.
344. Green, A.R., et al., *Studies on the effect of MDMA ('ecstasy') on the body temperature of rats housed at different ambient room temperatures*. *Br J Pharmacol*, 2005. **146**(2): p. 306-12.
345. Skelton, M.R., et al., *Distinct periods of developmental sensitivity to the effects of 3,4-(+/-)-methylenedioxymethamphetamine (MDMA) on behaviour and monoamines in rats*. *Int J Neuropsychopharmacol*, 2012. **15**(6): p. 811-24.
346. Schaefer, T.L., et al., *Cognitive impairments from developmental exposure to serotonergic drugs: citalopram and MDMA*. *Int J Neuropsychopharmacol*, 2013. **16**(6): p. 1383-94.

347. McElhatton, P.R., et al., *Congenital anomalies after prenatal ecstasy exposure [letter]*. *Lancet*, 1999. **354**(9188): p. 1441-2.
348. Bateman, D.N., et al., *A case control study to examine the pharmacological factors underlying ventricular septal defects in the North of England*. *Eur J Clin Pharmacol*, 2004. **60**(9): p. 635-41.
349. Ho, E., L. Karimi-Tabesh, and G. Koren, *Characteristics of pregnant women who use Ecstasy (3, 4- methylenedioxymethamphetamine)*. *Neurotoxicol Teratol*, 2001. **23**(6): p. 561-7.
350. Singer, L.T., et al., *Neurobehavioral outcomes of infants exposed to MDMA (Ecstasy) and other recreational drugs during pregnancy*. *Neurotoxicol Teratol*, 2012. **34**(3): p. 303-10.
351. Singer, L.T., et al., *Motor delays in MDMA (ecstasy) exposed infants persist to 2years*. *Neurotoxicol Teratol*, 2016.
352. Greene, S.L., et al., *Multiple toxicity from 3,4-methylenedioxymethamphetamine ("ecstasy")*. *Am J Emerg Med*, 2003. **21**(2): p. 121-4.
353. Mechan, A.O., et al., *The pharmacology of the acute hyperthermic response that follows administration of 3,4-methylenedioxymethamphetamine (MDMA, 'ecstasy') to rats*. *Br J Pharmacol*, 2002. **135**(1): p. 170-80.
354. Colado, M.I., et al., *Studies on the role of dopamine in the degeneration of 5-HT nerve endings in the brain of Dark Agouti rats following 3,4-methylenedioxymethamphetamine (MDMA or 'ecstasy') administration*. *Br J Pharmacol*, 1999. **126**(4): p. 911-24.
355. Hughes, J.C., M. McCabe, and R.J. Evans, *Intracranial haemorrhage associated with ingestion of 'ecstasy'*. *Arch Emerg Med*, 1993. **10**(4): p. 372-4.
356. Kaku, D.A. and D.H. Lowenstein, *Emergence of recreational drug abuse as a major risk factor for stroke in young adults*. *Ann Intern Med*, 1990. **113**(11): p. 821-7.
357. Perez, J.A., Jr., E.L. Arsura, and S. Strategos, *Methamphetamine-related stroke: four cases*. *J Emerg Med*, 1999. **17**(3): p. 469-71.
358. Rothwell, P.M. and R. Grant, *Cerebral venous sinus thrombosis induced by 'ecstasy'*. *J Neurol Neurosurg Psychiatry*, 1993. **56**(9): p. 1035.
359. Gledhill, J.A., et al., *Subarachnoid haemorrhage associated with MDMA abuse [letter]*. *J Neurol Neurosurg Psychiatry*, 1993. **56**(9): p. 1036-7.
360. Manchanda, S. and M.J. Connolly, *Cerebral infarction in association with Ecstasy abuse*. *Postgrad Med J*, 1993. **69**(817): p. 874-5.
361. Selmi, F., et al., *Intracerebral haemorrhage due to amphetamine abuse: report of two cases with underlying arteriovenous malformations*. *Br J Neurosurg*, 1995. **9**(1): p. 93-6.
362. Ho, M.P., J.L. Tsai, and Y.K. Wong, *Subarachnoid hemorrhage and death following coingestion of MDMA with other drugs*. *J Chin Med Assoc*, 2004. **67**(12): p. 640-3.
363. Johnson, J., et al., *Posterior spinal artery aneurysm rupture after 'Ecstasy' abuse*. *J Neurointerv Surg*, 2014.
364. Finley, J.J.t., M.A. Konstam, and J.E. Udelson, *Arginine vasopressin antagonists for the treatment of heart failure and hyponatremia*. *Circulation*, 2008. **118**(4): p. 410-21.
365. Bolignano, D., et al., *Copeptin (CTproAVP), a new tool for understanding the role of vasopressin in pathophysiology*. *Clin Chem Lab Med*, 2014. **52**(10): p. 1447-56.
366. Dowling, G.P., E.T.d. McDonough, and R.O. Bost, *'Eve' and 'Ecstasy'. A report of five deaths associated with the use of MDEA and MDMA*. *Jama*, 1987. **257**(12): p. 1615-7.
367. Milroy, C.M., J.C. Clark, and A.R. Forrest, *Pathology of deaths associated with "ecstasy" and "eve" misuse*. *J Clin Pathol*, 1996. **49**(2): p. 149-53.
368. Montastruc, F., et al., *Valvular heart disease in a patient taking 3,4-methylenedioxymethamphetamine (MDMA, 'Ecstasy')*. *Br J Clin Pharmacol*, 2012. **74**(3): p. 547-8.
369. Baggott, M., L. Jerome, and R. Stuart, *3,4-Methylenedioxymethamphetamine (MDMA); A review of the English-language scientific and medical literature: , in First Edition of*



- Investigator's Brochure for MDMA*. 2001, Multidisciplinary Association for Psychedelic Studies.
370. Brvar, M., et al., *Polydipsia as another mechanism of hyponatremia after 'ecstasy' (3,4-methyldioxymethamphetamine) ingestion*. *Eur J Emerg Med*, 2004. **11**(5): p. 302-4.
371. Henry, J.A. and J.G. Rella, *Medical risks associated with MDMA use*, in *Ecstasy: A Complete Guide*, J. Holland, Editor. 2001, Inner Traditions: Rochester, VT. p. 71-86.
372. Corona, G., et al., *Moderate hyponatremia is associated with increased risk of mortality: evidence from a meta-analysis*. *PLoS One*, 2013. **8**(12): p. e80451.
373. Ayus, J.C., S.G. Achinger, and A. Arief, *Brain cell volume regulation in hyponatremia: role of sex, age, vasopressin, and hypoxia*. *Am J Physiol Renal Physiol*, 2008. **295**(3): p. F619-24.
374. Lien, Y.H. and J.I. Shapiro, *Hyponatremia: clinical diagnosis and management*. *Am J Med*, 2007. **120**(8): p. 653-8.
375. Stoiser, B., et al., *Copeptin, a fragment of the vasopressin precursor, as a novel predictor of outcome in heart failure*. *Eur J Clin Invest*, 2006. **36**(11): p. 771-8.
376. Morgenthaler, N.G., *Copeptin: a biomarker of cardiovascular and renal function*. *Congest Heart Fail*, 2010. **16 Suppl 1**: p. S37-44.
377. Yalta, K., et al., *Copeptin and cardiovascular disease: a review of a novel neurohormone*. *Int J Cardiol*, 2013. **167**(5): p. 1750-9.
378. Rohrig, T.P. and R.W. Prouty, *Tissue distribution of methylenedioxymethamphetamine*. *J Anal Toxicol*, 1992. **16**(1): p. 52-3.
379. Beitia, G., et al., *Ecstasy-induced toxicity in rat liver*. *Liver*, 2000. **20**(1): p. 8-15.
380. Carvalho, M., et al., *Effect of 3,4-methylenedioxymethamphetamine ("ecstasy") on body temperature and liver antioxidant status in mice: influence of ambient temperature*. *Arch Toxicol*, 2002. **76**(3): p. 166-72.
381. Varela-Rey, M., et al., *3,4-methylenedioxymethamphetamine ("Ecstasy") stimulates the expression of alpha1(I) procollagen mRNA in hepatic stellate cells*. *Biochem Biophys Res Commun*, 1999. **259**(3): p. 678-82.
382. Carvalho, M., et al., *Mechanisms underlying the hepatotoxic effects of ecstasy*. *Curr Pharm Biotechnol*, 2010. **11**(5): p. 476-95.
383. Dykhuizen, R.S., et al., *Ecstasy induced hepatitis mimicking viral hepatitis*. *Gut*, 1995. **36**(6): p. 939-41.
384. Ellis, A.J., et al., *Acute liver damage and ecstasy ingestion*. *Gut*, 1996. **38**(3): p. 454-8.
385. Ellis, S.J., *Complications of "ecstasy" misuse*. *Lancet*, 1992. **340**(8821): p. 726.
386. Nadkarni, G.N., et al., *Serotonin syndrome, disseminated intravascular coagulation, and hepatitis after a single ingestion of MDMA in an Asian woman*. *Am J Ther*, 2014. **21**(4): p. e117-9.
387. Andreu, V., et al., *Ecstasy: a common cause of severe acute hepatotoxicity*. *J Hepatol*, 1998. **29**(3): p. 394-7.
388. Carvalho, M., F. Carvalho, and M.L. Bastos, *Is hyperthermia the triggering factor for hepatotoxicity induced by 3,4- methylenedioxymethamphetamine (ecstasy)? An in vitro study using freshly isolated mouse hepatocytes*. *Arch Toxicol*, 2001. **74**(12): p. 789-93.
389. da Silva, D.D., et al., *Mixtures of 3,4-methylenedioxymethamphetamine (ecstasy) and its major human metabolites act additively to induce significant toxicity to liver cells when combined at low, non-cytotoxic concentrations*. *J Appl Toxicol*, 2014. **34**(6): p. 618-27.
390. Cerretani, D., et al., *Acute administration of 3,4-methylenedioxymethamphetamine (MDMA) induces oxidative stress, lipoperoxidation and TNFalpha-mediated apoptosis in rat liver*. *Pharmacol Res*, 2011. **64**(5): p. 517-27.
391. Callahan, B.T., B.J. Cord, and G.A. Ricaurte, *Long-term impairment of anterograde axonal transport along fiber projections originating in the rostral raphe nuclei after treatment with fenfluramine or methylenedioxymethamphetamine*. *Synapse*, 2001. **40**(2): p. 113-21.

392. Hatzidimitriou, G., U.D. McCann, and G.A. Ricaurte, *Altered serotonin innervation patterns in the forebrain of monkeys treated with (+/-)3,4-methylenedioxymethamphetamine seven years previously: factors influencing abnormal recovery*. J Neurosci, 1999. **19**(12): p. 5096-107.
393. Green, A.R., et al., *The pharmacology and clinical pharmacology of 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy")*. Pharmacol Rev, 2003. **55**(3): p. 463-508.
394. O'Callaghan, J.P. and D.B. Miller, *Neurotoxicity profiles of substituted amphetamines in the C57BL/6J mouse*. J Pharmacol Exp Ther, 1994. **270**(2): p. 741-51.
395. Miller, D.B. and J.P. O'Callaghan, *Neurotoxicity of d-amphetamine in the C57BL/6J and CD-1 mouse. Interactions with stress and the adrenal system*. Ann N Y Acad Sci, 1996. **801**: p. 148-67.
396. Molliver, M.E., et al., *Neurotoxicity of MDMA and related compounds: anatomic studies*. Ann N Y Acad Sci, 1990. **600**: p. 649-61; discussion 661-4.
397. Sabol, K.E., et al., *Amphetamine analogs have differential effects on DRL 36-s schedule performance*. Psychopharmacology (Berl), 1995. **121**(1): p. 57-65.
398. Bosch, O.G., et al., *Verbal memory deficits are correlated with prefrontal hypometabolism in (18)FDG PET of recreational MDMA users*. PLoS One, 2013. **8**(4): p. e61234.
399. Taurah, L., C. Chandler, and G. Sanders, *Depression, impulsiveness, sleep, and memory in past and present polydrug users of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy)*. Psychopharmacology (Berl), 2014. **231**(4): p. 737-51.
400. Smithies, V., et al., *Dysfunctional overnight memory consolidation in ecstasy users*. J Psychopharmacol, 2014. **28**(8): p. 751-62.
401. Laws, K.R. and J. Kokkalis, *Ecstasy (MDMA) and memory function: a meta-analytic update*. Hum Psychopharmacol, 2007.
402. Zakzanis, K.K., Z. Campbell, and D. Jovanovski, *The neuropsychology of ecstasy (MDMA) use: a quantitative review*. Hum Psychopharmacol, 2007. **22**(7): p. 427-35.
403. Murphy, P.N., et al., *The effects of 'ecstasy' (MDMA) on visuospatial memory performance: findings from a systematic review with meta-analyses*. Hum Psychopharmacol, 2012. **27**(2): p. 113-38.
404. Wagner, D., et al., *A prospective study of learning, memory, and executive function in new MDMA users*. Addiction, 2013. **108**(1): p. 136-45.
405. Wagner, D., et al., *Learning, Memory, and Executive Function in New MDMA Users: A 2-Year Follow-Up Study*. Front Neurosci, 2015. **9**: p. 445.
406. Price, J.S., P. Shear, and K.M. Lisdahl, *Ecstasy exposure & gender: examining components of verbal memory functioning*. PLoS One, 2014. **9**(12): p. e115645.
407. Kuypers, K.P.C., et al., *Verbal memory impairment in polydrug Ecstasy users: A clinical perspective*. PLoS One, 2016. **11**(2): p. e0149438. .
408. Cole, J.C., et al., *The effects of stereotype threat on cognitive function in ecstasy users*. J Psychopharmacol, 2006. **20**(4): p. 518-25.
409. Wareing, M., et al., *Verbal working memory deficits in current and previous users of MDMA*. Hum Psychopharmacol, 2004. **19**(4): p. 225-34.
410. von Geusau, N.A., et al., *Impaired executive function in male MDMA ("ecstasy") users*. Psychopharmacology (Berl), 2004. **175**(3): p. 331-41.
411. Murphy, P.N., et al., *Executive working memory deficits in abstinent ecstasy/MDMA users: a critical review*. Neuropsychobiology, 2009. **60**(3-4): p. 159-75.
412. Hoshi, R., et al., *Neurocognitive function in current and ex-users of ecstasy in comparison to both matched polydrug-using controls and drug-naive controls*. Psychopharmacology (Berl), 2007. **194**(3): p. 371-9.

413. McGuire, P.K., H. Cope, and T.A. Fahy, *Diversity of psychopathology associated with use of 3,4-methylenedioxymethamphetamine ('Ecstasy')*. Br J Psychiatry, 1994. **165**(3): p. 391-5.
414. Huizink, A.C., et al., *Symptoms of anxiety and depression in childhood and use of MDMA: prospective, population based study*. Bmj, 2006. **332**(7545): p. 825-8.
415. Fonsart, J., et al., *Sprague-Dawley rats display metabolism-mediated sex differences in the acute toxicity of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy)*. Toxicol Appl Pharmacol, 2008.
416. Malpass, A., et al., *Acute toxicity of 3,4-methylenedioxymethamphetamine (MDMA) in Sprague-Dawley and Dark Agouti rats*. Pharmacol Biochem Behav, 1999. **64**(1): p. 29-34.
417. Palenicek, T., et al., *Increased sensitivity to the acute effects of MDMA ("ecstasy") in female rats*. Physiol Behav, 2005. **86**(4): p. 546-53.
418. Davis, W.M., H.T. Hatoum, and I.W. Waters, *Toxicity of MDA (3,4-methylenedioxymethamphetamine) considered for relevance to hazards of MDMA (Ecstasy) abuse*. Alcohol Drug Res, 1987. **7**(3): p. 123-34.
419. Green, A.R., E. O'Shea, and M.I. Colado, *A review of the mechanisms involved in the acute MDMA (ecstasy)-induced hyperthermic response*. Eur J Pharmacol, 2004. **500**(1-3): p. 3-13.
420. Cregg, M.T. and J.A. Tracey, *Ecstasy abuse in Ireland*. Ir Med J, 1993. **86**(4): p. 118-20.
421. Armenian, P., et al., *Multiple MDMA (Ecstasy) overdoses at a rave event: a case series*. J Intensive Care Med, 2013. **28**(4): p. 252-8.
422. CDC, *Ecstasy overdoses at a New Year's Eve rave--Los Angeles, California, 2010*. MMWR Morb Mortal Wkly Rep, 2010. **59**(22): p. 677-81.
423. Gowing, L.R., et al., *The health effects of ecstasy: a literature review*. Drug Alcohol Rev, 2002. **21**(1): p. 53-63.
424. Grunau, B.E., M.O. Wiens, and M. Greidanus, *Dantrolene for the treatment of MDMA toxicity*. CJEM, 2010. **12**(5): p. 457-9.
425. Russell, T., et al., *Ecstasy-induced delayed rhabdomyolysis and neuroleptic malignant syndrome in a patient with a novel variant in the ryanodine receptor type 1 gene*. Anaesthesia, 2012. **67**(9): p. 1021-4.
426. Shah, H.V., G.H. Irvine, and M. Bradley, *Rhabdomyolysis of the masseter muscle: case report*. Br J Oral Maxillofac Surg, 2008. **46**(2): p. 138-40.
427. Davies, O., et al., *Full recovery after severe serotonin syndrome, severe rhabdomyolysis, multi-organ failure and disseminated intravascular coagulopathy from MDMA*. Heart Lung, 2014. **43**(2): p. 117-9.
428. Hall, A.P., et al., *An unusual case of Ecstasy poisoning*. Intensive Care Med, 1996. **22**(7): p. 670-1.
429. Logan, A.S., et al., *Survival following 'Ecstasy' ingestion with a peak temperature of 42 degrees C*. Anaesthesia, 1993. **48**(11): p. 1017-8.
430. Singarajah, C. and N.G. Lavies, *An overdose of ecstasy. A role for dantrolene*. Anaesthesia, 1992. **47**(8): p. 686-7.
431. Bedford Russell, A.R., R.H. Schwartz, and S. Dawling, *Accidental ingestion of 'Ecstasy' (3,4-methylenedioxymethylamphetamine)*. Arch Dis Child, 1992. **67**(9): p. 1114-5.
432. Nimmo, S.M., et al., *Drug-induced hyperthermia*. Anaesthesia, 1993. **48**(10): p. 892-5.
433. Murthy, B.V., R.G. Wilkes, and N.B. Roberts, *Creatine kinase isoform changes following Ecstasy overdose*. Anaesth Intensive Care, 1997. **25**(2): p. 156-9.
434. Ramcharan, S., et al., *Survival after massive ecstasy overdose*. J Toxicol Clin Toxicol, 1998. **36**(7): p. 727-31.
435. Screatton, G.R., et al., *Hyperpyrexia and rhabdomyolysis after MDMA ("ecstasy") abuse*. Lancet, 1992. **339**(8794): p. 677-8.

436. Montgomery, H. and S. Myerson, *3,4-methylenedioxymethamphetamine (MDMA, or "ecstasy") and associated hypoglycemia*. Am J Emerg Med, 1997. **15**(2): p. 218.
437. Williams, A. and R. Unwin, *Prolonged elevation of serum creatine kinase (CK) without renal failure after ingestion of ecstasy*. Nephrol Dial Transplant, 1997. **12**(2): p. 361-2.
438. Hayner, G.N. and H. McKinney, *MDMA. The dark side of ecstasy*. J Psychoactive Drugs, 1986. **18**(4): p. 341-7.
439. Toth, A.R. and T. Varga, *Myocardium and striated muscle damage caused by licit or illicit drugs*. Leg Med (Tokyo), 2009. **11 Suppl 1**: p. S484-7.
440. Vanden Eede, H., et al., *Rhabdomyolysis in MDMA intoxication: a rapid and underestimated killer. "Clean" Ecstasy, a safe party drug?* J Emerg Med, 2012. **42**(6): p. 655-8.
441. Sadeghian, S., et al., *Two ecstasy-induced myocardial infarctions during a three month period in a young man*. Arch Iran Med, 2007. **10**(3): p. 409-12.
442. Diffley, M., et al., *Catecholaminergic polymorphic ventricular tachycardia found in an adolescent after a methylenedioxymethamphetamine and marijuana-induced cardiac arrest*. Crit Care Med, 2012. **40**(7): p. 2223-6.
443. Hoggett, K., D. McCoubrie, and D.M. Fatovich, *Ecstasy-induced acute coronary syndrome: something to rave about*. Emerg Med Australas, 2012. **24**(3): p. 339-42.
444. Moller, M., et al., *Ecstasy-induced myocardial infarction in a teenager: rare complication of a widely used illicit drug*. Clin Res Cardiol, 2010. **99**(12): p. 849-51.
445. Kapoor, A. and K.S. Jackson, *Seizures presenting in pregnancy: eclampsia or something else?* J Obstet Gynaecol, 2013. **33**(6): p. 630-1.
446. Okunoye, G.O. and P. Dutton, *Acute myocardial infarction in pregnancy following unlicensed use of methylenedioxymethamphetamine ('Ecstasy')*. Scott Med J, 2013. **58**(3): p. e4-6.
447. Mizia-Stec, K., et al., *Severe dilated cardiomyopathy as a consequence of Ecstasy intake*. Cardiovasc Pathol, 2008. **17**(4): p. 250-3.
448. Hua, Y.S., et al., *Contraction band necrosis in two ecstasy abusers: a latent lethal lesion associated with ecstasy*. Am J Forensic Med Pathol, 2009. **30**(3): p. 295-7.
449. Sano, R., et al., *A fatal case of myocardial damage due to misuse of the "designer drug" MDMA*. Leg Med (Tokyo), 2009. **11**(6): p. 294-7.
450. Suarez, R.V. and R. Riemersma, *"Ecstasy" and sudden cardiac death*. Am J Forensic Med Pathol, 1988. **9**(4): p. 339-41.
451. Lora-Tamayo, C., T. Tena, and A. Rodriguez, *Amphetamine derivative related deaths*. Forensic Sci Int, 1997. **85**(2): p. 149-57.
452. Bingham, C., et al., *Necrotizing renal vasculopathy resulting in chronic renal failure after ingestion of methamphetamine and 3,4-methylenedioxymethamphetamine ('ecstasy')*. Nephrol Dial Transplant, 1998. **13**(10): p. 2654-5.
453. Ghatol, A. and A. Kazory, *Ecstasy-associated acute severe hyponatremia and cerebral edema: a role for osmotic diuresis?* J Emerg Med, 2012. **42**(6): p. e137-40.
454. Beuerle, J.R. and F. Barrueto, Jr., *Neurogenic bladder and chronic urinary retention associated with MDMA abuse*. J Med Toxicol, 2008. **4**(2): p. 106-8.
455. Bryden, A.A., P.J. Rothwell, and P.H. O'Reilly, *Urinary retention with misuse of "ecstasy"*. Bmj, 1995. **310**(6978): p. 504.
456. Delgado, J.H., et al., *Acute, transient urinary retention from combined ecstasy and methamphetamine use*. J Emerg Med, 2004. **26**(2): p. 173-5.
457. Holden, R. and M.A. Jackson, *Near-fatal hyponatraemic coma due to vasopressin over-secretion after "ecstasy" (3,4-MDMA)*. Lancet, 1996. **347**(9007): p. 1052.
458. Henry, J.A., et al., *Low-dose MDMA ("ecstasy") induces vasopressin secretion*. Lancet, 1998. **351**(9118): p. 1784.
459. Gomez-Balaguer, M., et al., *Syndrome of inappropriate antidiuretic hormone secretion and "designer drugs" (ecstasy)*. J Pediatr Endocrinol Metab, 2000. **13**(4): p. 437-8.

460. Hegadoren, K.M., G.B. Baker, and M. Bourin, *3,4-Methylenedioxy analogues of amphetamine: defining the risks to humans*. *Neurosci Biobehav Rev*, 1999. **23**(4): p. 539-53.
461. Woodrow, G., P. Harnden, and J.H. Turney, *Acute renal failure due to accelerated hypertension following ingestion of 3,4-methylenedioxymethamphetamine ('ecstasy')*. *Nephrol Dial Transplant*, 1995. **10**(3): p. 399-400.
462. Cadier, M.A. and J.A. Clarke, *Ecstasy and Whizz at a rave resulting in a major burn plus complications*. *Burns*, 1993. **19**(3): p. 239-40.
463. Fahal, I.H., et al., *Acute renal failure after ecstasy*. *Bmj*, 1992. **305**(6844): p. 29.
464. Maxwell, D.L., M.I. Polkey, and J.A. Henry, *Hyponatraemia and catatonic stupor after taking "ecstasy"*. *Bmj*, 1993. **307**(6916): p. 1399.
465. Satchell, S.C. and M. Connaughton, *Inappropriate antidiuretic hormone secretion and extreme rises in serum creatinine kinase following MDMA ingestion*. *Br J Hosp Med*, 1994. **51**(9): p. 495.
466. Claffey, C., *A 26-year-old woman with sudden onset cerebral edema*. *J Emerg Nurs*, 2011. **37**(1): p. 55-6.
467. Gritti, P., et al., *Are we aware of ecstasy effects?* *Intern Emerg Med*, 2009. **4**(2): p. 175-7.
468. Parr, M.J., H.M. Low, and P. Botterill, *Hyponatraemia and death after "ecstasy" ingestion*. *Med J Aust*, 1997. **166**(3): p. 136-7.
469. O'Connor, A., et al., *Death from hyponatraemia-induced cerebral oedema associated with MDMA ("Ecstasy") use*. *N Z Med J*, 1999. **112**(1091): p. 255-6.
470. Dar, K.J. and M.E. McBrien, *MDMA induced hyperthermia: report of a fatality and review of current therapy*. *Intensive Care Med*, 1996. **22**(9): p. 995-6.
471. Mikula, T., J. Kozłowska, and A. Wiercinska-Drapalo, *Alcohol and ecstasy (MDMA-3,4-methylenedioxymethamphetamine) overdose as a reason for acute hepatitis with gall bladder inflammation*. *Drug Alcohol Rev*, 2009. **28**(6): p. 685.
472. Payance, A., et al., *Severe chronic hepatitis secondary to prolonged use of ecstasy and cocaine*. *Clin Res Hepatol Gastroenterol*, 2013. **37**(5): p. e109-13.
473. Colak, Y., et al., *Ecstasy induced fatal hepatic failure*. *J Gastrointestin Liver Dis*, 2011. **20**(2): p. 215-6.
474. Chadwick, I.S., et al., *Ecstasy, 3-4 methylenedioxymethamphetamine (MDMA), a fatality associated with coagulopathy and hyperthermia*. *J R Soc Med*, 1991. **84**(6): p. 371.
475. Campkin, N.J. and U.M. Davies, *Treatment of 'ecstasy' overdose with dantrolene*. *Anaesthesia*, 1993. **48**(1): p. 82-3.
476. Fineschi, V. and A. Masti, *Fatal poisoning by MDMA (ecstasy) and MDEA: a case report*. *Int J Legal Med*, 1996. **108**(5): p. 272-5.
477. Wollner, K., et al., *[Death after the intake of amphetamine/ecstasy: two case reports]*. *Arch Kriminol*, 2015. **235**(1-2): p. 53-61.
478. Clark, A.D. and N. Butt, *Ecstasy-induced very severe aplastic anaemia complicated by invasive pulmonary mucormycosis treated with allogeneic peripheral blood progenitor cell transplant*. *Clin Lab Haematol*, 1997. **19**(4): p. 279-81.
479. Marsh, J.C., et al., *Aplastic anaemia following exposure to 3,4-methylenedioxymethamphetamine ('Ecstasy')*. *Br J Haematol*, 1994. **88**(2): p. 281-5.
480. Richman, J. and A. Ferber, *Severe aplastic anemia with hot pockets following daily Ecstasy ingestion*. *Am J Hematol*, 2008. **83**(4): p. 321-2.
481. Gomez-Fernandez, C., et al., *Facial eruption related to snorting Ecstasy*. *Eur J Dermatol*, 2010. **20**(6): p. 853-4.
482. Sauvageau, A., *Death from a possible anaphylactic reaction to ecstasy*. *Clin Toxicol (Phila)*, 2008. **46**(2): p. 156.
483. Kahn, D.E., N. Ferraro, and R.J. Benveniste, *3 cases of primary intracranial hemorrhage associated with "Molly", a purified form of 3,4-methylenedioxymethamphetamine (MDMA)*. *J Neurol Sci*, 2012. **323**(1-2): p. 257-60.

484. De Smet, K., et al., *Bilateral globus pallidus infarcts in ecstasy use*. JBR-BTR, 2011. **94**(2): p. 93.
485. Gardner, H., et al., *Acute hippocampal sclerosis following ecstasy ingestion*. Neurology, 2009. **73**(7): p. 567-9.
486. Nifosi, F., et al., *Hippocampal remodelling after MDMA neurotoxicity: a single case study*. World J Biol Psychiatry, 2009. **10**(4 Pt 3): p. 961-8.
487. Ruis, C., et al., *Cognitive disorders after sporadic ecstasy use? A case report*. Neurocase, 2014.
488. Bruggemann, N., et al., *Acute amnesic syndrome due to MDMA exposure*. J Neurol, 2016.
489. Spatt, J., B. Glawar, and B. Mamoli, *A pure amnesic syndrome after MDMA ("ecstasy") ingestion*. J Neurol Neurosurg Psychiatry, 1997. **62**(4): p. 418-9.
490. Squier, M.V., et al., *Death after ecstasy ingestion: neuropathological findings*. J Neurol Neurosurg Psychiatry, 1995. **58**(6): p. 756.
491. Brand, H.S., S.N. Dun, and A.V. Nieuw Amerongen, *Ecstasy (MDMA) and oral health*. Br Dent J, 2008. **204**(2): p. 77-81.
492. Duxbury, A.J., *Ecstasy--dental implications*. Br Dent J, 1993. **175**(1): p. 38.
493. Nixon, P.J., C.C. Youngson, and A. Beese, *Tooth surface loss: does recreational drug use contribute?* Clin Oral Investig, 2002. **6**(2): p. 128-30.
494. Marchesi, C., M. Tonna, and C. Maggini, *Obsessive-compulsive disorder followed by psychotic episode in long-term ecstasy misuse*. World J Biol Psychiatry, 2009. **10**(4 Pt 2): p. 599-602.
495. Potash, M.N., K.A. Gordon, and K.L. Conrad, *Persistent Psychosis and Medical Complications After a Single Ingestion of MDMA "Ecstasy": A Case Report and Review of the Literature*. Psychiatry (Edgmont), 2009. **6**(7): p. 40-4.
496. Tabatabaei, S.A., M. Soleimani, and A. Khodabandeh, *A case of autoenucleation associated with a contralateral field defect*. Orbit, 2011. **30**(3): p. 165-8.
497. Mutlu, H., et al., *'Ecstasy'(MDMA)-induced pneumomediastinum and epidural pneumatosis*. Diagn Interv Radiol, 2005. **11**(3): p. 150-1.
498. Onwudike, M., *Ecstasy induced retropharyngeal emphysema*. J Accid Emerg Med, 1996. **13**(5): p. 359-61.
499. Pittman, J.A. and J.C. Pounsford, *Spontaneous pneumomediastinum and Ecstasy abuse*. J Accid Emerg Med, 1997. **14**(5): p. 335-6.
500. Quin, G.I., G.M. McCarthy, and D.K. Harries, *Spontaneous pneumomediastinum and ecstasy abuse*. J Accid Emerg Med, 1999. **16**(5): p. 382.
501. Gungadeen, A. and J. Moor, *Extensive subcutaneous emphysema and pneumomediastinum after ecstasy ingestion*. Case Rep Otolaryngol, 2013. **2013**: p. 795867.
502. Clause, A.L., et al., *Spontaneous pneumomediastinum and epidural pneumatosis after oral ecstasy consumption*. Acta Clin Belg, 2014. **69**(2): p. 146-8.
503. Peters, N.F., R. Gosselin, and K.L. Verstraete, *A rare case of diffuse alveolar hemorrhage following oral amphetamine intake*. JBR-BTR, 2014. **97**(1): p. 42-3.
504. Barrett, P.J., *'Ecstasy' misuse--overdose or normal dose?* Anaesthesia, 1993. **48**(1): p. 83.
505. O'Neill, D. and J.K. Dart, *Methylenedioxymphetamine ('Ecstasy') associated keratopathy*. Eye, 1993. **7**(Pt 6): p. 805-6.
506. Schroeder, B. and S. Brieden, *Bilateral sixth nerve palsy associated with MDMA ("ecstasy") abuse*. Am J Ophthalmol, 2000. **129**(3): p. 408-9.
507. Schifano, F., et al., *Death rates from ecstasy (MDMA, MDA) and polydrug use in England and Wales 1996-2002*. Hum Psychopharmacol, 2003. **18**(7): p. 519-24.
508. Abad, S., et al., *MDMA enhances hippocampal-dependent learning and memory under restrictive conditions, and modifies hippocampal spine density*. Psychopharmacology (Berl), 2014. **231**(5): p. 863-74.

509. Fantegrossi, W.E., et al., *Behavioral and neurochemical consequences of long-term intravenous self-administration of MDMA and its enantiomers by rhesus monkeys*. *Neuropsychopharmacology*, 2004. **29**(7): p. 1270-81.
510. Schenk, S., et al., *Development, maintenance and temporal pattern of self-administration maintained by ecstasy (MDMA) in rats*. *Psychopharmacology (Berl)*, 2003. **169**(1): p. 21-7.
511. Trigo, J.M., et al., *A reliable model of intravenous MDMA self-administration in naive mice*. *Psychopharmacology (Berl)*, 2006. **184**(2): p. 212-20.
512. Lile, J.A., J.T. Ross, and M.A. Nader, *A comparison of the reinforcing efficacy of 3,4-methylenedioxymethamphetamine (MDMA, "ecstasy") with cocaine in rhesus monkeys*. *Drug Alcohol Depend*, 2005. **78**(2): p. 135-40.
513. Wang, Z. and W.L. Woolverton, *Estimating the relative reinforcing strength of (+/-)-3,4-methylenedioxymethamphetamine (MDMA) and its isomers in rhesus monkeys: comparison to (+)-methamphetamine*. *Psychopharmacology (Berl)*, 2007. **189**(4): p. 483-8.
514. Robledo, P., et al., *Study of the behavioural responses related to the potential addictive properties of MDMA in mice*. *Naunyn Schmiedebergs Arch Pharmacol*, 2004. **369**(3): p. 338-49.
515. Verheyden, S.L., J.A. Henry, and H.V. Curran, *Acute, sub-acute and long-term subjective consequences of 'ecstasy' (MDMA) consumption in 430 regular users*. *Hum Psychopharmacol*, 2003. **18**(7): p. 507-17.
516. Cottler, L.B., K.S. Leung, and A.B. Abdallah, *Test-re-test reliability of DSM-IV adopted criteria for 3,4-methylenedioxymethamphetamine (MDMA) abuse and dependence: a cross-national study*. *Addiction*, 2009. **104**(10): p. 1679-90.
517. Meyer, J.S., *3,4-methylenedioxymethamphetamine (MDMA): current perspectives*. *Subst Abuse Rehabil*, 2013. **4**: p. 83-99.
518. Topp L.J., H. W.D., and H. J., *Is there a dependence syndrome for ecstasy?* 1997, Randwick, NSW: National Drug & Alcohol Research Centre.
519. Cottler, L.B., et al., *Ecstasy abuse and dependence among adolescents and young adults: applicability and reliability of DSM-IV criteria*. *Hum Psychopharmacol*, 2001. **16**(8): p. 599-606.
520. von Sydow, K., et al., *Use, abuse and dependence of ecstasy and related drugs in adolescents and young adults-a transient phenomenon? Results from a longitudinal community study*. *Drug Alcohol Depend*, 2002. **66**(2): p. 147-59.
521. Degenhardt, L., R. Bruno, and L. Topp, *Is ecstasy a drug of dependence?* *Drug Alcohol Depend*, 2010. **107**(1): p. 1-10.
522. Topp, L., et al., *Ecstasy use in Australia: patterns of use and associated harm*. *Drug Alcohol Depend*, 1999. **55**(1-2): p. 105-15.
523. Davis, A.K. and H. Rosenberg, *The prevalence, intensity, and assessment of craving for MDMA/ecstasy in recreational users*. *J Psychoactive Drugs*, 2014. **46**(2): p. 154-61.
524. De La Garza, R., 2nd, K.R. Fabrizio, and A. Gupta, *Relevance of rodent models of intravenous MDMA self-administration to human MDMA consumption patterns*. *Psychopharmacology (Berl)*, 2007. **189**(4): p. 425-434.
525. Stolaroff, M., *The Secret Chief Revealed: Conversations with a pioneer of the underground therapy movement*. 2004, Sarasota FL: Multidisciplinary Association for Psychedelic Studies.
526. Grob, C., *MDMA research: preliminary investigations with human subjects*. *Int J Drug Policy*, 1998. **9**(2): p. 119-124.
527. Bouso, J.C., et al., *MDMA-assisted psychotherapy using low doses in a small sample of women with chronic posttraumatic stress disorder*. *J Psychoactive Drugs*, 2008. **40**(3): p. 225-36.



528. de la Torre, R., et al., *Pharmacology of MDMA in humans*. Ann N Y Acad Sci, 2000. **914**: p. 225-37.
529. Dumont, G.J. and R.J. Verkes, *A review of acute effects of 3,4-methylenedioxymethamphetamine in healthy volunteers*. J Psychopharmacol, 2006. **20**(2): p. 176-87.
530. Hysek, C.M., et al., *Pharmacokinetic and pharmacodynamic effects of methylphenidate and MDMA administered alone or in combination*. Int J Neuropsychopharmacol, 2014. **17**(3): p. 371-81.
531. Pizarro, N., et al., *Stereochemical analysis of 3,4-methylenedioxymethamphetamine and its main metabolites in human samples including the catechol-type metabolite (3,4-dihydroxymethamphetamine)*. Drug Metab Dispos, 2004. **32**(9): p. 1001-7.
532. de la Torre, R., et al., *Clinical pharmacokinetics of amphetamine and related substances: monitoring in conventional and non-conventional matrices*. Clin Pharmacokinet, 2004. **43**(3): p. 157-85.
533. de Boer, D., et al., *Gas chromatographic/mass spectrometric assay for profiling the enantiomers of 3,4-methylenedioxymethamphetamine and its chiral metabolites using positive chemical ionization ion trap mass spectrometry*. J Mass Spectrom, 1997. **32**(11): p. 1236-46.
534. Helmlin, H.J., et al., *Analysis of 3,4-methylenedioxymethamphetamine (MDMA) and its metabolites in plasma and urine by HPLC-DAD and GC-MS*. J Anal Toxicol, 1996. **20**(6): p. 432-40.
535. Lanz, M., R. Brenneisen, and W. Thormann, *Enantioselective determination of 3,4-methylene-dioxymethamphetamine and two of its metabolites in human urine by cyclodextrin-modified capillary zone electrophoresis*. Electrophoresis, 1997. **18**(6): p. 1035-43.
536. Ortuno, J., et al., *Quantification of 3,4-methylenedioxymetamphetamine and its metabolites in plasma and urine by gas chromatography with nitrogen-phosphorus detection*. J Chromatogr B Biomed Sci Appl, 1999. **723**(1-2): p. 221-32.
537. Helmlin, H.J. and R. Brenneisen, *Determination of psychotropic phenylalkylamine derivatives in biological matrices by high-performance liquid chromatography with photodiode-array detection*. J Chromatogr, 1992. **593**(1-2): p. 87-94.
538. Pizarro, N., et al., *Stereochemical analysis of 3,4-methylenedioxymethamphetamine and its main metabolites by gas chromatography/mass spectrometry*. Rapid Commun Mass Spectrom, 2003. **17**(4): p. 330-6.
539. Pizarro, N., et al., *Determination of MDMA and its metabolites in blood and urine by gas chromatography-mass spectrometry and analysis of enantiomers by capillary electrophoresis*. J Anal Toxicol, 2002. **26**(3): p. 157-65.
540. Segura, M., et al., *3,4-Dihydroxymethamphetamine (HHMA). A Major in Vivo 3,4-methylenedioxymethamphetamine (MDMA) Metabolite in Humans*. Chem Res Toxicol, 2001. **14**(9): p. 1203-1208.
541. Desrosiers, N.A., et al., *Oral fluid and plasma 3,4-methylenedioxymethamphetamine (MDMA) and metabolite correlation after controlled oral MDMA administration*. Anal Bioanal Chem, 2013. **405**(12): p. 4067-76.
542. de la Torre, R., et al., *Human pharmacology of MDMA: pharmacokinetics, metabolism, and disposition*. Ther Drug Monit, 2004. **26**(2): p. 137-44.
543. Abraham, T.T., et al., *Urinary MDMA, MDA, HHMA, and HMA excretion following controlled MDMA administration to humans*. J Anal Toxicol, 2009. **33**(8): p. 439-46.
544. Fallon, J.K., et al., *Stereospecific analysis and enantiomeric disposition of 3, 4-methylenedioxymethamphetamine (Ecstasy) in humans [published erratum appears in Clin Chem 1999 Sep;45(9):1585]*. Clin Chem, 1999. **45**(7): p. 1058-69.
545. Schwaninger, A.E., et al., *Urinary excretion kinetics of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) and its phase I and phase II*

- metabolites in humans following controlled MDMA administration.* Clin Chem, 2011. **57**(12): p. 1748-56.
546. Farre, M., et al., *Human pharmacology of 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) after repeated doses taken 4h apart Human pharmacology of MDMA after repeated doses taken 4h apart.* Eur Neuropsychopharmacol, 2015.
547. Ramamoorthy, S. and R.D. Blakely, *Phosphorylation and sequestration of serotonin transporters differentially modulated by psychostimulants.* Science, 1999. **285**(5428): p. 763-766.
548. Yang, J., et al., *Implications of mechanism-based inhibition of CYP2D6 for the pharmacokinetics and toxicity of MDMA.* J Psychopharmacol, 2006. **20**(6): p. 842-9.
549. O'Mathuna, B., et al., *The consequences of 3,4-methylenedioxymethamphetamine induced CYP2D6 inhibition in humans.* J Clin Psychopharmacol, 2008. **28**(5): p. 523-9.
550. Yubero-Lahoz, S., et al., *Sex Differences in 3,4-Methylenedioxymethamphetamine (MDMA; Ecstasy)-Induced Cytochrome P450 2D6 Inhibition in Humans.* Clin Pharmacokinet, 2011. **50**(5): p. 319-29.
551. Yubero-Lahoz, S., et al., *Changes in CYP1A2 activity in humans after 3,4-methylenedioxymethamphetamine (MDMA, ecstasy) administration using caffeine as a probe drug.* Drug Metab Pharmacokinet, 2012. **27**(6): p. 605-13.
552. Pardo-Lozano, R., et al., *Clinical Pharmacology of 3,4-Methylenedioxymethamphetamine (MDMA, "Ecstasy"): The Influence of Gender and Genetics (CYP2D6, COMT, 5-HTT).* PLoS One, 2012. **7**(10): p. e47599.
553. Gouzoulis-Mayfrank, E., et al., *Psychopathological, neuroendocrine and autonomic effects of 3,4-methylenedioxyethylamphetamine (MDE), psilocybin and d-methamphetamine in healthy volunteers. Results of an experimental double-blind placebo-controlled study.* Psychopharmacology (Berl), 1999. **142**(1): p. 41-50.
554. Davison, D. and A.C. Parrott, *Ecstasy (MDMA) in recreational users: Self-reported psychological and physiological effects.* Human Psychopharmacology Clinical & Experimental, 1997. **12**(3): p. 221-226.
555. Liechti, M.E., et al., *Acute psychological effects of 3,4-methylenedioxymethamphetamine (MDMA, "Ecstasy") are attenuated by the serotonin uptake inhibitor citalopram.* Neuropsychopharmacology, 2000. **22**(5): p. 513-21.
556. Liechti, M.E. and F.X. Vollenweider, *The serotonin uptake inhibitor citalopram reduces acute cardiovascular and vegetative effects of 3,4-methylenedioxymethamphetamine ('Ecstasy') in healthy volunteers.* J Psychopharmacol, 2000. **14**(3): p. 269-74.
557. Pacifici, R., et al., *Paroxetine inhibits acute effects of 3,4-methylenedioxymethamphetamine on the immune system in humans.* J Pharmacol Exp Ther, 2004. **309**(1): p. 285-92.
558. Tancer, M. and C.E. Johanson, *The effects of fluoxetine on the subjective and physiological effects of 3,4-methylenedioxymethamphetamine (MDMA) in humans.* Psychopharmacology (Berl), 2007. **189**(4): p. 565-73.
559. Liechti, M.E., et al., *Psychological and physiological effects of MDMA ("Ecstasy") after pretreatment with the 5-HT(2) antagonist ketanserin in healthy humans.* Neuropsychopharmacology, 2000. **23**(4): p. 396-404.
560. Hysek, C.M., et al., *The norepinephrine transporter inhibitor reboxetine reduces stimulant effects of MDMA ("ecstasy") in humans.* Clin Pharmacol Ther, 2011. **90**(2): p. 246-55.
561. Simmler, L.D., C.M. Hysek, and M.E. Liechti, *Sex differences in the effects of MDMA (ecstasy) on plasma copeptin in healthy subjects.* J Clin Endocrinol Metab, 2011. **96**(9): p. 2844-50.
562. Moron, J.A., et al., *Dopamine uptake through the norepinephrine transporter in brain regions with low levels of the dopamine transporter: evidence from knock-out mouse lines.* J Neurosci, 2002. **22**(2): p. 389-95.

563. Liechti, M.E. and F.X. Vollenweider, *Acute psychological and physiological effects of MDMA ("Ecstasy") after haloperidol pretreatment in healthy humans*. Eur Neuropsychopharmacol, 2000. **10**(4): p. 289-95.
564. Schmid, Y., et al., *Interactions between Bupropion and 3,4-Methylenedioxymethamphetamine in Healthy Subjects*. J Pharmacol Exp Ther, 2015. **353**(1): p. 102-11.
565. Domes, G., et al., *Oxytocin Attenuates Amygdala Responses to Emotional Faces Regardless of Valence*. Biol Psychiatry, 2007.
566. Kirsch, P., et al., *Oxytocin modulates neural circuitry for social cognition and fear in humans*. J Neurosci, 2005. **25**(49): p. 11489-93.
567. Bartz, J.A. and E. Hollander, *The neuroscience of affiliation: forging links between basic and clinical research on neuropeptides and social behavior*. Horm Behav, 2006. **50**(4): p. 518-28.
568. Wirth, M.M. and O.C. Schultheiss, *Effects of affiliation arousal (hope of closeness) and affiliation stress (fear of rejection) on progesterone and cortisol*. Horm Behav, 2006. **50**(5): p. 786-95.
569. Kuypers, K.P., et al., *No Evidence that MDMA-Induced Enhancement of Emotional Empathy Is Related to Peripheral Oxytocin Levels or 5-HT1a Receptor Activation*. PLoS One, 2014. **9**(6): p. e100719.
570. Leng, G. and M. Ludwig, *Intranasal Oxytocin: Myths and Delusions*. Biol Psychiatry, 2016. **79**(3): p. 243-50.
571. Grob, C., *Unpublished data on human study of psychological and physiological effects of MDMA*. 2001.
572. Kuypers, K.P., et al., *Inhibition of MDMA-induced increase in cortisol does not prevent acute impairment of verbal memory*. Br J Pharmacol, 2013. **168**(3): p. 607-17.
573. Hysek, C.M., et al., *MDMA enhances emotional empathy and prosocial behavior*. Soc Cogn Affect Neurosci, 2013.
574. Seibert, J., et al., *Acute effects of 3,4-methylenedioxymethamphetamine and methylphenidate on circulating steroid levels in healthy subjects*. Neuroendocrinology, 2014. **100**(1): p. 17-25.
575. Pacifici, R., et al., *Cell-mediated immune response in MDMA users after repeated dose administration: studies in controlled versus noncontrolled settings*. Ann N Y Acad Sci, 2002. **965**: p. 421-33.
576. Ramaekers, J.G., et al., *Involvement of inferior parietal lobules in prospective memory impairment during acute MDMA (ecstasy) intoxication: an event-related fMRI study*. Neuropsychopharmacology, 2009. **34**(7): p. 1641-8.
577. de Sousa Fernandes Perna, E.B., et al., *Memory and mood during MDMA intoxication, with and without memantine pretreatment*. Neuropharmacology, 2014. **87**: p. 198-205.
578. Pacifici, R., et al., *Immunomodulating activity of MDMA*. Ann N Y Acad Sci, 2000. **914**: p. 215-24.
579. Pacifici, R., et al., *Immunomodulating properties of MDMA alone and in combination with alcohol: a pilot study*. Life Sci, 1999. **65**(26): p. L309-16.
580. Pacifici, R., et al., *Effects of repeated doses of MDMA ("ecstasy") on cell-mediated immune response in humans*. Life Sci, 2001. **69**(24): p. 2931-41.
581. Liechti, M.E., *Effects of MDMA on body temperature in humans*. Temperature, 2014. **1**(3): p. 192-200.
582. Dao, C.K., S.M. Nowinski, and E.M. Mills, *The heat is on: Molecular mechanisms of drug-induced hyperthermia*. Temperature, 2014. **1**(3): p. 183-191.
583. Parrott, A.C. and L. Young, *Saturday night fever in ecstasy/MDMA dance clubbers: Heightened body temperature and associated psychobiological changes*. Temperature, 2014. **1**(3): p. 214-219.

584. Irvine, R.J., et al., *Plasma drug concentrations and physiological measures in 'dance party' participants*. *Neuropsychopharmacology*, 2006. **31**(2): p. 424-30.
585. Cole, J.C., et al., *Preliminary evidence of the cardiovascular effects of polysubstance misuse in nightclubs*. *J Psychopharmacol*, 2005. **19**(1): p. 67-70.
586. Parrott, A.C., et al., *Dance clubbing on MDMA and during abstinence from Ecstasy/MDMA: prospective neuroendocrine and psychobiological changes*. *Neuropsychobiology*, 2008. **57**(4): p. 165-80.
587. Clark, C.M., et al., *Acute effects of MDMA on autonomic cardiac activity and their relation to subjective prosocial and stimulant effects*. *Psychophysiology*, 2015. **52**(3): p. 429-35.
588. Baggott, M.J., et al., *3,4-Methylenedioxymethamphetamine (MDMA, 'Ecstasy') and Prazosin Interactions in Humans.*, in *70th Annual Meeting of the College on Problems of Drug Dependence*. 2008: San Juan, Puerto Rico.
589. Kirkpatrick, M.G., et al., *MDMA effects consistent across laboratories*. *Psychopharmacology (Berl)*, 2014. **231**(19): p. 3899-905.
590. Mithoefer, M.C., et al., *Safety, Efficacy and Durability of MDMA-Assisted Psychotherapy in Subjects with Chronic, Treatment-Resistant Posttraumatic Stress Disorder: Completed and Ongoing Randomized, Controlled, Triple-blind Phase 2 Pilot Studies, in Military medical research across the continuum of care*. 2012: Fort Lauderdale, FL.
591. Phelps, E.A., et al., *Activation of the left amygdala to a cognitive representation of fear*. *Nat Neurosci*, 2001. **4**(4): p. 437-41.
592. Kuypers, K.P., et al., *MDMA intoxication and verbal memory performance: a placebo-controlled pharmac-MRI study*. *J Psychopharmacol*, 2011. **25**(8): p. 1053-61.
593. Chang, L., et al., *Effect of ecstasy [3,4-methylenedioxymethamphetamine (MDMA)] on cerebral blood flow: a co-registered SPECT and MRI study*. *Psychiatry Res*, 2000. **98**(1): p. 15-28.
594. Frei, E., et al., *Localization of MDMA-induced brain activity in healthy volunteers using low resolution brain electromagnetic tomography (LORETA)*. *Hum Brain Mapp*, 2001. **14**(3): p. 152-65.
595. Lamers, C.T., et al., *Dissociable effects of a single dose of ecstasy (MDMA) on psychomotor skills and attentional performance*. *J Psychopharmacol*, 2003. **17**(4): p. 379-87.
596. Tancer, M.E. and C.E. Johanson, *The subjective effects of MDMA and mCPP in moderate MDMA users*. *Drug Alcohol Depend*, 2001. **65**(1): p. 97-101.
597. Huxster, J.K., A. Pirona, and M.J. Morgan, *The sub-acute effects of recreational ecstasy (MDMA) use: a controlled study in humans*. *J Psychopharmacol*, 2006. **20**(2): p. 281-90.
598. Neff, K., *The Development and validation of a scale to measure self-compassion*. *Self and Identity*, 2003. **2**: p. 223-250.
599. Kuypers, K.P. and J.G. Ramaekers, *Transient memory impairment after acute dose of 75mg 3,4-Methylene-dioxymethamphetamine*. *J Psychopharmacol*, 2005. **19**(6): p. 633-9.
600. Ramaekers, J.G. and K.P. Kuypers, *Acute effects of 3,4-methylenedioxymethamphetamine (MDMA) on behavioral measures of impulsivity: alone and in combination with alcohol*. *Neuropsychopharmacology*, 2006. **31**(5): p. 1048-55.
601. Kuypers, K.P., N. Samyn, and J.G. Ramaekers, *MDMA and alcohol effects, combined and alone, on objective and subjective measures of actual driving performance and psychomotor function*. *Psychopharmacology (Berl)*, 2006. **187**(4): p. 467-75.
602. Kuypers, K.P. and J.G. Ramaekers, *Acute dose of MDMA (75 mg) impairs spatial memory for location but leaves contextual processing of visuospatial information unaffected*. *Psychopharmacology (Berl)*, 2007. **189**(4): p. 557-63.
603. Ramaekers, J.G., K.P. Kuypers, and N. Samyn, *Stimulant effects of 3,4-methylenedioxymethamphetamine (MDMA) 75 mg and methylphenidate 20 mg on actual driving during intoxication and withdrawal*. *Addiction*, 2006. **101**(11): p. 1614-21.

604. Spronk, D.B., et al., *The acute effects of MDMA and ethanol administration on electrophysiological correlates of performance monitoring in healthy volunteers*. *Psychopharmacology (Berl)*, 2014. **231**(14): p. 2877-88.
605. Randolph, C., et al., *The Repeatable Battery for the Assessment of Neuropsychological Status (RBANS): preliminary clinical validity*. *J Clin Exp Neuropsychol*, 1998. **20**(3): p. 310-9.
606. Gronwall, D.M., *Paced auditory serial-addition task: a measure of recovery from concussion*. *Percept Mot Skills*, 1977. **44**(2): p. 367-73.
607. Roman, D.D., et al., *Extended norms for the paced auditory serial addition task*. *Clin Neuropsychol*, 1991. **5**(1): p. 33-40.
608. Hoshi, R., J. Bisla, and H.V. Curran, *The acute and sub-acute effects of 'ecstasy' (MDMA) on processing of facial expressions: preliminary findings*. *Drug Alcohol Depend*, 2004. **76**(3): p. 297-304.
609. Frye, C.G., et al., *MDMA decreases the effects of simulated social rejection*. *Pharmacol Biochem Behav*, 2014. **117**: p. 1-6.
610. Kirkpatrick, M.G. and H. de Wit, *MDMA: a social drug in a social context*. *Psychopharmacology (Berl)*, 2015. **232**(6): p. 1155-63.
611. Schmid, Y., et al., *Effects of methylphenidate and MDMA on appraisal of erotic stimuli and intimate relationships*. *Eur Neuropsychopharmacol*, 2015. **25**(1): p. 17-25.
612. MacDonald, E., et al., *A review of safety, side-effects and subjective reactions to intranasal oxytocin in human research*. *Psychoneuroendocrinology*, 2011. **36**(8): p. 1114-26.
613. Bershad, A.K., et al., *Oxytocin receptor gene variation predicts subjective responses to MDMA*. *Soc Neurosci*, 2016.
614. Bosker, W.M., et al., *Dose-related effects of MDMA on psychomotor function and mood before, during, and after a night of sleep loss*. *Psychopharmacology (Berl)*, 2010. **209**(1): p. 69-76.
615. Gradus, J.L., et al., *Posttraumatic stress disorder and completed suicide*. *Am J Epidemiol*, 2010. **171**(6): p. 721-7.
616. Conner, K.R., et al., *Posttraumatic stress disorder and suicide in 5.9 million individuals receiving care in the veterans health administration health system*. *J Affect Disord*, 2014. **166**: p. 1-5.
617. US Food and Drug Administration, *Guidance for industry: suicidal ideation and behavior: prospective assessment of occurrence in clinical trials*. Silver Springs, MD: US Food and Drug Administration Available at: <http://www.fda.gov/Drugs/GuidanceComplianceRegulatoryInformation/Guidances/ucm315156.htm>. Accessed October, 2012. **5**: p. 2012.
618. Stone, T.E., C. Swanson, and M.D. Feldman, *Venlafaxine discontinuation syndrome and suicidal ideation: a case series*. *J Clin Psychopharmacol*, 2007. **27**(1): p. 94-5.
619. Tint, A., P.M. Haddad, and I.M. Anderson, *The effect of rate of antidepressant tapering on the incidence of discontinuation symptoms: a randomised study*. *J Psychopharmacol*, 2008. **22**(3): p. 330-2.
620. Valuck, R.J., H.D. Orton, and A.M. Libby, *Antidepressant discontinuation and risk of suicide attempt: a retrospective, nested case-control study*. *J Clin Psychiatry*, 2009. **70**(8): p. 1069-77.
621. Nilsson, M.E., et al., *Columbia Suicide Severity Rating Scale Scoring and Data Analysis Guide*, in *CSSRS Scoring Version 2.0*. 2013: [http://www.cssrs.columbia.edu/documents/ScoringandDataAnalysisGuide\\_Feb2013.pdf](http://www.cssrs.columbia.edu/documents/ScoringandDataAnalysisGuide_Feb2013.pdf). p. 1-13.
622. Sanger, G.J., *5-hydroxytryptamine and the gastrointestinal tract: where next?* *Trends Pharmacol Sci*, 2008. **29**(9): p. 465-471.

623. Hexel, M. and G. Sonneck, *Somatoform symptoms, anxiety, and depression in the context of traumatic life experiences by comparing participants with and without psychiatric diagnoses*. *Psychopathology*, 2002. **35**(5): p. 303-12.
624. Afari, N., et al., *Psychological trauma and functional somatic syndromes: a systematic review and meta-analysis*. *Psychosom Med*, 2014. **76**(1): p. 2-1.
625. Pietrzak, R.H., et al., *Psychiatric comorbidity of full and partial posttraumatic stress disorder among older adults in the United States: results from wave 2 of the National Epidemiologic Survey on Alcohol and Related Conditions*. *Am J Geriatr Psychiatry*, 2012. **20**(5): p. 380-90.
626. Baker, D.G., et al., *Relationship between posttraumatic stress disorder and self-reported physical symptoms in Persian Gulf War veterans*. *Arch Intern Med*, 1997. **157**(18): p. 2076-8.
627. Hoge, C.W., et al., *Association of posttraumatic stress disorder with somatic symptoms, health care visits, and absenteeism among Iraq war veterans*. *Am J Psychiatry*, 2007. **164**(1): p. 150-3.
628. Blake, D.D., et al., *A clinician rating scale for assessing current and lifetime PTSD: the CAPS-1*. *Behav Ther*, 1990. **13**: p. 187-188.
629. Nagy, L.M., et al., *Open prospective trial of fluoxetine for posttraumatic stress disorder*. *J Clin Psychopharmacol*, 1993. **13**(2): p. 107-13.
630. Weathers, F.W., *Clinician-Administered PTSD Scale (CAPS): Technical Manual*. 2004, Los Angeles, CA: Western Psychological Services.
631. UNDOC. *World Drug Report*. New York, NY: United Nations Office on Drugs and Crime. 2011.
632. Gable RS. Comparison of acute lethal toxicity of commonly abused psychoactive substances. *Addiction (Abingdon, England)*. 2004. **99**(6): p. 686-96.
633. Tiangco DA, Halcomb S, Lattanzio FA, Jr., Hargave BY. 3,4-Methylenedioxymethamphetamine alters left ventricular function and activates nuclear factor-Kappa B (NF-kappaB) in a time and dose dependent manner. *Int J Mol Sci*. 2010. **11**(12): p. 4843-63.
634. Glennon, R.A. and R. Young, Further investigation of the discriminative stimulus properties of MDA. *Pharmacol Biochem Behav*, 1984. **20**(4): p. 501-5.
635. Kamien, J.B., et al., The effects of (+/-)-methylenedioxymethamphetamine and (+/-)-methylenedioxyamphetamine in monkeys trained to discriminate (+)-amphetamine from saline. *Drug Alcohol Depend*, 1986. **18**(2): p. 139-47.
636. Evans, S.M. and C.E. Johanson, Discriminative stimulus properties of (+/-)-3,4-methylenedioxymethamphetamine and (+/-)-3,4-methylenedioxyamphetamine in pigeons. *Drug Alcohol Depend*, 1986. **18**(2): p. 159-64.
637. Oberlender, R. and D.E. Nichols, Drug discrimination studies with MDMA and amphetamine. *Psychopharmacology (Berl)*, 1988. **95**(1): p. 71-6.
638. Callahan, P.M. and J.B. Appel, Differences in the stimulus properties of 3,4-methylenedioxyamphetamine and 3,4-methylenedioxymethamphetamine in animals trained to discriminate hallucinogens from saline. *J Pharmacol Exp Ther*, 1988. **246**(3): p.866-70
639. Glennon, R.A., et al., Discriminative stimulus properties of MDA analogs. *Biol Psychiatry*, 1982. **17**(7): p. 807-14.
640. Goodwin, A.K. and L.E. Baker, A three-choice discrimination procedure dissociates the discriminative stimulus effects of d-amphetamine and (+/-)-MDMA in rats. *Exp Clin Psychopharmacol*, 2000. **8**(3): p. 415-23.
641. Goodwin, A.K., D.M. Pynnonen, and L.E. Baker, Serotonergic-dopaminergic mediation of MDMA's discriminative stimulus effects in a three-choice discrimination. *Pharmacol Biochem Behav*, 2003. **74**(4): p. 987-95.

## 9.0 Appendix

**Table 31: Percentage of Observations of Spontaneously Reported Reactions During Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015**

Dose	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
Sessions	24	19	15	11	14	255	4
<b>Anxiety</b>							
Mild	1 (4%)	---	---	2 (18%)	4 (29%)	48 (19%)	1 (25%)
Moderate	9 (38%)	---	7 (47%)	2 (18%)	1 (7%)	61 (24%)	---
Severe	4 (17%)	---	1 (7%)	---	1 (7%)	13 (5%)	---
Total	14 (58%)	0 (0%)	8 (53%)	4 (36%)	6 (43%)	122 (48%)	1 (25%)
<b>Diarrhea</b>							
Mild	---	---	---	---	---	5 (2%)	---
Moderate	---	---	---	---	---	---	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	5 (2%)	0 (0%)
<b>Disturbance in Attention</b>							
Mild	---	---	3 (20%)	1 (9%)	1 (7%)	28 (11%)	---
Moderate	1 (4%)	---	---	---	---	10 (4%)	---
Severe	---	---	---	---	---	---	---
Total	1 (4%)	0 (0%)	3 (20%)	1 (9%)	1 (7%)	38 (15%)	0 (0%)
<b>Dizziness</b>							
Mild	1 (4%)	---	1 (7%)	1 (9%)	1 (9%)	40 (16%)	2 (50%)
Moderate	1 (4%)	---	---	---	---	19 (7%)	---
Severe	---	---	---	---	---	1 (<1%)	---
Total	2 (8%)	0 (0%)	1 (7%)	1 (9%)	1 (7%)	60 (24%)	2 (50%)
<b>Somnolence</b>							
Mild	---	---	2 (13%)	---	1 (7%)	11 (4%)	---
Moderate	3 (13%)	---	---	---	---	8 (3%)	---
Severe	---	---	---	---	---	---	---
Total	3 (13%)	0 (0%)	2 (13%)	0 (0%)	1 (7%)	19 (7%)	0 (0%)



<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Sessions</b>	24	19	15	11	14	255	4
<b>Dry Mouth</b>							
Mild	---	1 (5%)	2 (13%)	1 (9%)	---	23 (9%)	1 (25%)
Moderate	---	---	1 (7%)	2 (18%)	---	19 (7%)	1 (25%)
Severe	---	---	---	---	---	---	---
<b>Total</b>	0 (0%)	1 (5%)	3 (20%)	3 (27%)	0 (0%)	42 (16%)	2 (50%)
<b>Fatigue</b>							
Mild	3 (13%)	4 (21%)	6 (40%)	---	2 (14%)	32 (13%)	---
Moderate	7 (29%)	---	2 (13%)	3 (27%)	2 (14%)	52 (20%)	---
Severe	---	---	---	---	---	3 (1%)	---
<b>Total</b>	10 (42%)	4 (21%)	8 (53%)	3 (27%)	4 (28%)	87 (34%)	0 (0%)
<b>Headache</b>							
Mild	5 (21%)	1 (5%)	6 (40%)	3 (27%)	10 (71%)	57 (22%)	1 (25%)
Moderate	7 (29%)	---	1 (7%)	1 (9%)	---	37 (15%)	---
Severe	---	---	---	---	---	1 (<1%)	---
<b>Total</b>	12 (50%)	1 (5%)	7 (47%)	4 (36%)	10 (71%)	95 (37%)	1 (25%)
<b>Sensation of Heaviness</b>							
Mild	---	---	---	---	---	5 (2%)	1 (25%)
Moderate	---	---	---	---	---	6 (2%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	11 (4%)	1 (25%)
<b>Disturbed Gait</b>							
Mild	1 (4%)	3 (16%)	---	---	2 (14%)	35 (14%)	2 (50%)
Moderate	---	---	---	---	---	6 (2%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	1 (4%)	3 (16%)	0 (0%)	0 (0%)	2 (14%)	41 (16%)	2 (50%)
<b>Judgment Impaired</b>							
Mild	---	---	---	---	---	1 (<1%)	---
Moderate	---	---	---	---	---	---	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (<1%)	0 (0%)

<b>Dose</b>	<b>0 mg</b> (N=12)	<b>25 mg</b> (N=7)	<b>30 mg</b> (N=7)	<b>40 mg</b> (N=6)	<b>75 mg</b> (N=7)	<b>100-125 mg</b> (N=100)	<b>150 mg</b> (N=3)
<b>Sessions</b>	24	19	15	11	14	255	4
<b>Irritability</b>							
Mild	---	---	1 (7%)	---	---	3 (1%)	---
Moderate	3 (13%)	---	---	---	---	3 (1%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	<b>3 (13%)</b>	<b>0 (0%)</b>	<b>1 (7%)</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>6 (2%)</b>	<b>0 (0%)</b>
<b>Insomnia</b>							
Mild	6 (25%)	1 (5%)	1 (7%)	---	---	21 (8%)	1 (25%)
Moderate	6 (25%)	2 (11%)	---	---	3 (21%)	31 (12%)	---
Severe	---	1 (5%)	---	---	---	6 (2%)	2 (50%)
<b>Total</b>	<b>12 (50%)</b>	<b>4 (21%)</b>	<b>1 (7%)</b>	<b>0 (0%)</b>	<b>3 (21%)</b>	<b>58 (23%)</b>	<b>3 (75%)</b>
<b>Muscle Tightness (jaw)</b>							
Mild	2 (8%)	1 (5%)	---	1 (9%)	4 (29%)	62 (24%)	---
Moderate	3 (13%)	---	---	2 (18%)	1 (7%)	73 (29%)	1 (25%)
Severe	---	---	---	---	---	6 (2%)	1 (25%)
<b>Total</b>	<b>5 (21%)</b>	<b>1 (5%)</b>	<b>0 (0%)</b>	<b>3 (27%)</b>	<b>5 (36%)</b>	<b>141 (55%)</b>	<b>2 (50%)</b>
<b>Decreased Appetite</b>							
Mild	1 (4%)	2 (11%)	3 (20%)	---	4 (29%)	63 (25%)	---
Moderate	1 (4%)	1 (5%)	---	---	---	40 (16%)	1 (25%)
Severe	---	1 (5%)	---	---	---	3 (1%)	---
<b>Total</b>	<b>2 (8%)</b>	<b>4 (21%)</b>	<b>3 (20%)</b>	<b>0 (0%)</b>	<b>4 (29%)</b>	<b>106 (42%)</b>	<b>1 (25%)</b>
<b>Depressed Mood</b>							
Mild	---	---	---	---	---	13 (5%)	---
Moderate	2 (8%)	1 (5%)	1 (7%)	---	1 (7%)	12 (5%)	---
Severe	---	---	---	---	---	2 (1%)	---
<b>Total</b>	<b>2 (8%)</b>	<b>1 (5%)</b>	<b>1 (7%)</b>	<b>0 (0.0%)</b>	<b>1 (7%)</b>	<b>27 (11%)</b>	<b>0 (0.0%)</b>
<b>Muscle Tightness</b>							
Mild	1 (4%)	---	5 (33%)	1 (9%)	1 (7%)	44 (17%)	---
Moderate	2 (8%)	---	2 (13%)	2 (18%)	2 (14%)	25 (10%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	<b>3 (13%)</b>	<b>0 (0%)</b>	<b>7 (47%)</b>	<b>3 (27%)</b>	<b>3 (21%)</b>	<b>69 (27%)</b>	<b>0 (0%)</b>

<b>Dose</b>	<b>0 mg</b> (N=12)	<b>25 mg</b> (N=7)	<b>30 mg</b> (N=7)	<b>40 mg</b> (N=6)	<b>75 mg</b> (N=7)	<b>100-125 mg</b> (N=100)	<b>150 mg</b> (N=3)
<b>Sessions</b>	24	19	15	11	14	255	4
<b>Nausea</b>							
Mild	2 (8%)	2 (11%)	1 (7%)	---	2 (14%)	34 (13%)	1 (25%)
Moderate	1 (4%)	---	2 (13%)	---	---	28 (11%)	---
Severe	---	---	---	---	---	6	---
<b>Total</b>	<b>3 (13%)</b>	<b>2 (11%)</b>	<b>3 (20%)</b>	<b>0 (0%)</b>	<b>2 (14%)</b>	<b>68 (27%)</b>	<b>1 (25%)</b>
<b>Hypersomnia</b>							
Mild	2 (8%)	---	1 (7%)	---	1 (7%)	3 (1%)	1 (25%)
Moderate	1 (4%)	---	---	2 (18%)	---	7 (3%)	---
Severe	---	1 (5%)	---	---	---	---	---
<b>Total</b>	<b>3 (13%)</b>	<b>1 (5%)</b>	<b>1 (7%)</b>	<b>2 (18%)</b>	<b>1 (7%)</b>	<b>10 (4%)</b>	<b>1 (25%)</b>
<b>Nystagmus</b>							
Mild	---	---	---	---	1 (7%)	17 (7%)	1 (25%)
Moderate	---	---	---	---	---	7 (3%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>1 (7%)</b>	<b>24 (9%)</b>	<b>1 (25%)</b>
<b>Paresthesia</b>							
Mild	---	---	1 (7%)	---	1 (7%)	11 (4%)	1 (25%)
Moderate	---	---	---	---	---	5 (2%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	<b>0 (0%)</b>	<b>0 (0%)</b>	<b>1 (7%)</b>	<b>0 (0%)</b>	<b>1 (7%)</b>	<b>16 (6%)</b>	<b>1 (25%)</b>
<b>Hyperhidrosis</b>							
Mild	---	---	2 (13%)	---	3 (21%)	44 (17%)	1 (25%)
Moderate	1 (4%)	---	---	---	---	19 (7%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	<b>1 (4%)</b>	<b>0 (0%)</b>	<b>2 (13%)</b>	<b>0 (0%)</b>	<b>3 (21%)</b>	<b>63 (25%)</b>	<b>1 (25%)</b>
<b>Restlessness</b>							
Mild	---	1 (5%)	5 (33%)	---	3 (21%)	40 (16%)	---
Moderate	2 (8%)	---	1 (7%)	---	2 (4%)	22 (9%)	1 (25%)
Severe	---	---	---	---	---	1 (<1%)	---
<b>Total</b>	<b>2 (8%)</b>	<b>1 (5%)</b>	<b>6 (40%)</b>	<b>0 (0%)</b>	<b>5 (36%)</b>	<b>63 (25%)</b>	<b>1 (25%)</b>

<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Sessions</b>	24	19	15	11	14	255	4
<b>Obsessive Rumination</b>							
Mild	---	1 (5%)	---	---	---	10 (4%)	---
Moderate	1 (4%)	---	2 (13%)	---	---	6 (2%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	1 (4%)	1 (5%)	2 (13%)	0 (0%)	0 (0%)	16 (6%)	0 (0%)
<b>Feeling Cold</b>							
Mild	2 (8%)	1 (5%)	7 (47%)	---	4 (29%)	47 (18%)	---
Moderate	1 (4%)	---	1 (7%)	---	2 (14%)	20 (8%)	---
Severe	---	---	---	---	---	1 (<1%)	---
<b>Total</b>	3 (13%)	1 (5%)	8 (53%)	0 (0%)	6 (43%)	68 (27%)	0 (0%)
<b>Thirst</b>							
Mild	1 (4%)	---	1 (7%)	---	---	29 (11%)	1 (25%)
Moderate	---	---	---	1 (9%)	---	14 (5%)	1 (25%)
Severe	---	---	---	---	---	---	---
<b>Total</b>	1 (4%)	0 (0%)	1 (7%)	1 (9%)	0 (0%)	43 (17%)	2 (50%)
<b>Asthenia</b>							
Mild	1 (4%)	---	---	---	---	7 (3%)	1 (25%)
Moderate	---	---	---	---	---	6 (2%)	---
Severe	---	---	---	---	---	---	---
<b>Total</b>	1 (4%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	13 (5%)	1 (25%)

**Table 32: Spontaneously Reported Reactions on Day 1-7 After All Experimental Sessions in Studies MP-1, MP-2, MP-4, MP-8, MP-9, MP-12, MDA-1, MAA-1, and MP1-E2 as of 01 October 2015**

<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Observations</b>	168	133	105	77	98	1785	21
	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)
<b>Anxiety</b>							
Mild	26 (15%)	4 (3%)	25 (24%)	8 (10%)	5 (5%)	239 (13%)	---
Moderate	35 (21%)	---	10 (10%)	3 (4%)	1 (1%)	172 (10%)	---
Severe	2 (1%)	---	---	---	---	24 (1%)	---
Total	63 (38%)	4 (3%)	35 (33%)	11 (14%)	6 (6%)	435 (24%)	0 (0%)
<b>Diarrhea</b>							
Mild	1 (1%)	---	4 (4%)	---	1 (1%)	15 (1%)	---
Moderate	---	---	---	---	---	4 (0%)	---
Severe	---	---	---	---	---	2 (0%)	---
Total	1 (1%)	0 (0%)	4 (4%)	0 (0%)	1 (1%)	21 (1%)	0 (0%)
<b>Disturbance in Attention</b>							
Mild	11 (7%)	---	2 (2%)	2 (3%)	---	141 (8%)	---
Moderate	11 (7%)	---	2 (2%)	2 (3%)	---	38 (2%)	---
Severe	---	---	---	---	---	3 (<1%)	---
Total	22 (13%)	0 (0%)	4 (4%)	4 (5%)	0 (0%)	182 (10%)	0 (0%)
<b>Dizziness</b>							
Mild	5 (3%)	6 (5%)	---	---	1 (1%)	122 (7%)	---
Moderate	---	1 (1%)	---	---	---	19 (1%)	---
Severe	---	---	---	---	---	2 (<1%)	---
Total	5 (3%)	7 (5%)	0 (0%)	0 (0%)	1 (1%)	143 (8%)	0 (0%)
<b>Somnolence</b>							
Mild	2 (1%)	3 (2%)	1 (1%)	---	---	2 (<1%)	---
Moderate	5 (3%)	---	---	---	---	---	---
Severe	---	---	---	---	---	---	---
Total	7 (4%)	3 (2%)	1 (1%)	0 (0%)	0 (0%)	2 (<1%)	0 (0%)
<b>Dry Mouth</b>							
Mild	---	---	---	1 (1%)	---	23 (1%)	1 (5%)
Moderate	---	---	---	---	---	1 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	0 (0%)	1 (1%)	0 (0%)	24 (1%)	1 (5%)

<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Observations</b>	168	133	105	77	98	1785	21
	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)
<b>Fatigue</b>							
Mild	23 (14%)	16 (12%)	20 (19%)	3 (4%)	27 (28%)	245 (14%)	2 (10%)
Moderate	27 (16%)	8 (6%)	8 (8%)	4 (5%)	1 (1%)	163 (9%)	14 (67%)
Severe	1 (1%)	1 (1%)	---	---	---	13 (1%)	2 (10%)
Total	51 (30%)	25 (19%)	28 (27%)	7 (9%)	28 (29%)	421 (24%)	18 (86%)
<b>Headache</b>							
Mild	9 (5%)	13 (10%)	1 (1%)	3 (4%)	4 (4%)	76 (4%)	---
Moderate	5 (3%)	13 (10%)	---	3 (4%)	---	33 (2%)	---
Severe	---	---	---	---	---	3 (<1%)	---
Total	14 (8%)	26 (20%)	1 (1%)	6 (8%)	4 (4%)	112 (6%)	0 (0%)
<b>Sensation of Heaviness</b>							
Mild	---	1 (1%)	---	---	---	3 (<1%)	---
Moderate	---	---	---	---	---	3 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	1 (1%)	0 (0%)	0 (0%)	0 (0%)	6 (<1%)	0 (0%)
<b>Disturbed Gait</b>							
Mild	1 (1%)	---	---	---	---	10 (1%)	---
Moderate	---	---	---	---	---	3 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	1 (1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	13 (1%)	0 (0%)
<b>Judgment Impaired</b>							
Mild	---	---	1 (1%)	1 (1%)	---	7 (<1%)	---
Moderate	---	---	---	1 (1%)	---	2 (<1%)	---
Severe	---	---	---	---	---	1 (<1%)	---
Total	0 (0%)	0 (0%)	1 (1%)	2 (3%)	0 (0%)	10 (1%)	0 (0%)
<b>Irritability</b>							
Mild	12 (7%)	3 (2%)	5 (5%)	---	4 (4%)	69 (4%)	---
Moderate	9 (5%)	---	3 (3%)	3 (4%)	---	48 (3%)	---
Severe	---	---	---	---	---	2 (<1%)	---
Total	21 (13%)	3 (2%)	8 (8%)	3 (4%)	4 (4%)	119 (7%)	0 (0%)

<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Observations</b>	168	133	105	77	98	1785	21
	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)
<b>Insomnia</b>							
Mild	19 (11%)	17 (13%)	8 (8%)	5 (6%)	6 (6%)	158 (9%)	---
Moderate	29 (17%)	13 (10%)	9 (9%)	5 (6%)	2 (2%)	88 (5%)	---
Severe	1 (1%)	8 (6%)	1 (1%)	5 (6%)	---	8 (<1%)	---
Total	49 (29%)	38 (29%)	18 (17%)	15 (19%)	8 (8%)	254 (14%)	0 (0%)
<b>Muscle Tightness (jaw)</b>							
Mild	2 (1%)	---	---	6 (8%)	2 (2%)	90 (5%)	---
Moderate	1 (1%)	---	---	4 (5%)	---	20 (1%)	---
Severe	---	---	---	---	---	2 (<1%)	---
Total	3 (2%)	0 (0%)	0 (0%)	10 (13%)	2 (2%)	112 (6%)	0 (0%)
<b>Decreased Appetite</b>							
Mild	---	10 (8%)	2 (2%)	1 (1%)	2 (2%)	89 (5%)	---
Moderate	---	2 (2%)	---	1 (1%)	---	67 (4%)	---
Severe	---	3 (2%)	---	---	---	1 (<1%)	---
Total	0 (0%)	15 (11%)	2 (2%)	2 (3%)	2 (2%)	157 (9%)	0 (0%)
<b>Depressed Mood</b>							
Mild	13 (8%)	14 (11%)	4 (4%)	3 (4%)	---	114 (6%)	3 (14%)
Moderate	8 (5%)	11 (8%)	4 (4%)	---	---	101 (6%)	---
Severe	---	---	---	---	---	17 (1%)	---
Total	21 (13%)	25 (19%)	8 (8%)	3 (4%)	0 (0%)	232 (13%)	3 (14%)
<b>Muscle Tightness</b>							
Mild	3 (2%)	1 (1%)	3 (3%)	9 (12%)	6 (6%)	84 (5%)	---
Moderate	6 (4%)	---	---	7 (9%)	1 (1%)	26 (1%)	---
Severe	---	---	---	---	---	5 (<1%)	---
Total	9 (5%)	1 (1%)	3 (3%)	16 (21%)	7 (7%)	115 (6%)	0 (0%)
<b>Nausea</b>							
Mild	9 (5%)	5 (4%)	2 (2%)	1 (1%)	---	66 (4%)	---
Moderate	3 (2%)	2 (2%)	---	1 (1%)	---	28 (2%)	---
Severe	---	7 (5%)	---	---	---	4 (<1%)	---
Total	12 (7%)	14 (11%)	2 (2%)	2 (3%)	0 (0%)	98 (5%)	0 (0%)



<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Observations</b>	168	133	105	77	98	1785	21
	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)
<b>Hypersomnia</b>							
Mild	15 (9%)	8 (6%)	9 (9%)	3 (4%)	17 (17%)	145 (8%)	2 (10%)
Moderate	9 (5%)	7 (5%)	1 (1%)	3 (4%)	---	63 (4%)	---
Severe	---	---	1 (1%)	---	---	---	---
Total	24 (14%)	15 (11%)	11 (10%)	6 (8%)	17 (17%)	208 (12%)	2 (10%)
<b>Nystagmus</b>							
Mild	---	---	---	---	---	1 (<1%)	---
Moderate	---	---	---	---	---	---	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (<1%)	0 (0%)
<b>Paresthesia</b>							
Mild	---	---	---	---	---	4 (<1%)	---
Moderate	---	---	---	---	---	4 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	8 (<1%)	0 (0%)
<b>Hyperhidrosis</b>							
Mild	1 (1%)	---	---	---	1 (1%)	13 (1%)	---
Moderate	---	---	---	1 (1%)	---	1 (<1%)	---
Severe	1 (1%)	---	---	---	---	---	---
Total	2 (1%)	0 (0%)	0 (0%)	1 (1%)	1 (1%)	14 (1%)	0 (0%)
<b>Restlessness</b>							
Mild	---	2 (2%)	1 (1%)	3 (4%)	---	24 (1%)	---
Moderate	---	---	---	2 (3%)	---	24 (1%)	---
Severe	---	---	---	---	---	5 (<1%)	---
Total	0 (0%)	2 (2%)	1 (1%)	5 (6%)	0 (0%)	53 (3%)	0 (0%)
<b>Obsessive Rumination</b>							
Mild	2 (1%)	6 (5%)	2 (2%)	4 (5%)	1 (1%)	37 (2%)	---
Moderate	6 (4%)	1 (1%)	---	2 (3%)	---	38 (2%)	---
Severe	---	---	---	---	---	5 (<1%)	---
Total	8 (5%)	7 (5%)	2 (2%)	6 (8%)	1 (1%)	80 (4%)	0 (0%)

<b>Dose</b>	0 mg (N=12)	25 mg (N=7)	30 mg (N=7)	40 mg (N=6)	75 mg (N=7)	100-125 mg (N=100)	150 mg (N=3)
<b>Observations</b>	168	133	105	77	98	1785	21
	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)	Counts (%)
<b>Feeling Cold</b>							
Mild	---	2 (2%)	2 (2%)	---	---	26 (1%)	---
Moderate	---	---	---	---	---	7 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	2 (2%)	2 (2%)	0 (0%)	0 (0%)	33 (2%)	0 (0%)
<b>Thirst</b>							
Mild	---	---	---	1 (1%)	---	7 (<1%)	---
Moderate	---	---	---	---	---	1 (<1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	0 (0%)	1 (1%)	0 (0%)	8 (<1%)	0 (0%)
<b>Asthenia</b>							
Mild	---	---	1 (1%)	---	---	20 (1%)	---
Moderate	---	---	---	1 (1%)	---	11 (1%)	---
Severe	---	---	---	---	---	---	---
Total	0 (0%)	0 (0%)	1 (1%)	1 (1%)	0 (0%)	31 (2%)	0 (0%)